

3. Absence of non-vertebrogenic pathology per clinical assessment or radiology studies that could explain the source of the patient's pain, including but not limited to fracture, tumor, infection, or significant deformity, AND
4. Evidence of Type 1 or Type 2 Modic changes on MRI, such as inflammation, edema, vertebral endplate changes, disruption and fissuring of the endplate, vascularized fibrous tissues within the adjacent marrow, hypotensive signals (Type 1 Modic change), and changes to the vertebral body marrow including replacement of normal bone marrow by fat, and hypertensive signals (Type 2 Modic change), in 1 or more vertebrae from L3-S1.

* Pain assessment and a disability scale must be obtained at baseline to be used for functional assessment.

** Non-surgical management may include but is not limited to:

- Avoidance of activities that aggravate pain;
- Trial of Chiropractic manipulation;
- Trial of Physical Therapy;
- Cognitive support and recovery reassurance;
- Injection therapy – epidural and/or facet;
- Spine biomechanics education;
- Specific lumbar exercise program;
- Home use of heat/cold modalities;
- Low impact aerobic exercise as tolerated;
- Pharmacotherapy (e.g., non-narcotic analgesics, NSAIDs, muscle relaxants, neuroleptics, and narcotics).

Patients must have undergone careful screening, evaluation, and diagnosis by a multidisciplinary team prior to thermal destruction of the intraosseous BVN (such screening must include psychological, as well as physical evaluation). Documentation of the history and careful screening must be available in the patient chart if requested.

Limitations

Services that are not reasonable and necessary cannot be covered by Medicare in the following:

1. Skeletally immature patients (≤ 18 years old);
2. Severe cardiac or pulmonary compromise;
3. Active systemic infection or local infection at the intended treatment level;
4. Bleeding diathesis;
5. Pregnancy;
6. Primary radicular pain into the lower extremities (defined as nerve pain following a dermatomal distribution and that correlates with nerve compression on imaging);
7. Previous lumbar/lumbosacral spine surgery at the intended treatment level (with the exception of discectomy/laminectomy if performed >6 months prior to BVN nerve ablation and radicular pain resolved);