

# Mathematics I- MATH F111

Saranya G. Nair  
Department of Mathematics

BITS Pilani

September 25, 2024



If you are traveling along a curve in space, the Cartesian  $\vec{i}, \vec{j}$ , and  $\vec{k}$  coordinate system for representing the vectors describing your motion may not be very relevant to you.

If you are traveling along a curve in space, the Cartesian  $\vec{i}, \vec{j}$ , and  $\vec{k}$  coordinate system for representing the vectors describing your motion may not be very relevant to you.

Instead, the vectors that represent your forward direction (the unit tangent vector  $\mathbf{T}$ ), the direction in which your path is turning (the unit normal vector  $\mathbf{N}$ ), and the tendency of your motion to “twist” out of the plane created by these vectors in the direction perpendicular to this plane (defined by the unit binormal vector  $\mathbf{B} = \mathbf{T} \times \mathbf{N}$ ) are likely to be more important.



If you are traveling along a curve in space, the Cartesian  $\vec{i}, \vec{j}$ , and  $\vec{k}$  coordinate system for representing the vectors describing your motion may not be very relevant to you.

Instead, the vectors that represent your forward direction (the unit tangent vector  $\mathbf{T}$ ), the direction in which your path is turning (the unit normal vector  $\mathbf{N}$ ), and the tendency of your motion to “twist” out of the plane created by these vectors in the direction perpendicular to this plane (defined by the unit binormal vector  $\mathbf{B} = \mathbf{T} \times \mathbf{N}$ ) are likely to be more important.

Expressing the acceleration vector along the curve as a linear combination of this  $\mathbf{TNB}$  frame of mutually orthogonal unit vectors traveling with the motion can reveal much about the nature of the path and motion along it.

## Definition

The binormal vector of a curve in space is  $\mathbf{B} = \mathbf{T} \times \mathbf{N}$  which is a unit vector that is orthogonal to both  $\mathbf{T}$  and  $\mathbf{N}$ . Together  $\mathbf{T}$ ,  $\mathbf{N}$ , and  $\mathbf{B}$  define a moving righthanded vector frame that plays a significant role in calculating the paths of particles moving through space. It is called the Frenet ("fre-nay") frame (after Jean-Frédéric Frenet, 1816–1900), or the **TNB** frame.



# Tangential and Normal Components of Acceleration

When an object is accelerated by gravity or brakes, we often need to know how much of the acceleration acts in the direction of motion, which is the direction of the tangent vector  $\mathbf{T}$ .

# Tangential and Normal Components of Acceleration

When an object is accelerated by gravity or brakes, we often need to know how much of the acceleration acts in the direction of motion, which is the direction of the tangent vector  $\mathbf{T}$ . We can calculate this using the Chain Rule to rewrite  $\vec{v}$  as

$$\vec{v} = \frac{d\vec{r}}{dt} = \frac{d\vec{r}}{ds} \frac{ds}{dt} = \mathbf{T} \frac{ds}{dt}.$$

# Tangential and Normal Components of Acceleration

When an object is accelerated by gravity or brakes, we often need to know how much of the acceleration acts in the direction of motion, which is the direction of the tangent vector  $\mathbf{T}$ . We can calculate this using the Chain Rule to rewrite  $\vec{v}$  as

$$\vec{v} = \frac{d\vec{r}}{dt} = \frac{d\vec{r}}{ds} \frac{ds}{dt} = \mathbf{T} \frac{ds}{dt}.$$

Then we differentiate both ends of this string of equalities to get



# Tangential and Normal Components of Acceleration

When an object is accelerated by gravity or brakes, we often need to know how much of the acceleration acts in the direction of motion, which is the direction of the tangent vector  $\mathbf{T}$ . We can calculate this using the Chain Rule to rewrite  $\vec{v}$  as

$$\vec{v} = \frac{d\vec{r}}{dt} = \frac{d\vec{r}}{ds} \frac{ds}{dt} = \mathbf{T} \frac{ds}{dt}.$$

Then we differentiate both ends of this string of equalities to get

$$\begin{aligned}\vec{a} &= \frac{d\vec{v}}{dt} = \frac{d}{dt} \left( \mathbf{T} \frac{ds}{dt} \right) = \frac{d^2s}{dt^2} \mathbf{T} + \frac{ds}{dt} \frac{d\mathbf{T}}{dt} \\ &= \frac{d^2s}{dt^2} \mathbf{T} + \frac{ds}{dt} \left( \frac{d\mathbf{T}}{ds} \frac{ds}{dt} \right) = \frac{d^2s}{dt^2} \mathbf{T} + \frac{ds}{dt} \left( \kappa \mathbf{N} \frac{ds}{dt} \right) \\ &= \frac{d^2s}{dt^2} \mathbf{T} + \kappa \left( \frac{ds}{dt} \right)^2 \mathbf{N}.\end{aligned}$$

## Definition

If the acceleration vector is written as  $\vec{a} = a_T \mathbf{T} + a_N \mathbf{N}$ , then

$a_T = \frac{d^2s}{dt^2} = \frac{d}{dt}|\vec{v}|$  and  $a_N = \kappa \left(\frac{ds}{dt}\right)^2 = \kappa|\vec{v}|^2$  are the tangential and normal scalar components of acceleration.

## Definition


If the acceleration vector is written as  $\vec{a} = a_T \mathbf{T} + a_N \mathbf{N}$ , then  $a_T = \frac{d^2 s}{dt^2} = \frac{d}{dt} |\vec{v}|$  and  $a_N = \kappa \left( \frac{ds}{dt} \right)^2 = \kappa |\vec{v}|^2$  are the tangential and normal scalar components of acceleration.

- Note that the binormal vector  $\mathbf{B}$  does not appear in above equation. No matter how the path of the moving object we are watching may appear to twist and turn in space, the acceleration  $\vec{a}$  always lies in the plane of  $\mathbf{T}$  and  $\mathbf{N}$  orthogonal to  $\mathbf{B}$ .


## Definition

If the acceleration vector is written as  $\vec{a} = a_T \mathbf{T} + a_N \mathbf{N}$ , then  $a_T = \frac{d^2 s}{dt^2} = \frac{d}{dt} |\vec{v}|$  and  $a_N = \kappa \left( \frac{ds}{dt} \right)^2 = \kappa |\vec{v}|^2$  are the tangential and normal scalar components of acceleration.

- Note that the binormal vector  $\mathbf{B}$  does not appear in above equation. No matter how the path of the moving object we are watching may appear to twist and turn in space, the acceleration  $\vec{a}$  always lies in the plane of  $\mathbf{T}$  and  $\mathbf{N}$  orthogonal to  $\mathbf{B}$ .
- The equation also tells us exactly how much of the acceleration takes place tangent to the motion and how much takes place normal to the motion.



By definition, acceleration  $\vec{a}$  is the rate of change of velocity  $\vec{v}$ , and in general, both the length and direction of  $\vec{v}$  change as an object moves along its path.



By definition, acceleration  $\vec{a}$  is the rate of change of velocity  $\vec{v}$ , and in general, both the length and direction of  $\vec{v}$  change as an object moves along its path.

- The tangential component of acceleration  $a_T$  measures the rate of change of the length of  $\vec{v}$  (that is, the change in the speed).
- The normal component of acceleration  $a_N$  measures the rate of change of the direction of  $\vec{v}$ .

The normal scalar component of the acceleration is the curvature times the square of the speed. This explains why you have to hold on when your car makes a sharp (large  $\kappa$ ), high-speed (large  $|\vec{v}|$ ) turn. If you double the speed of your car, you will experience four times the normal component of acceleration for the same curvature.

The normal scalar component of the acceleration is the curvature times the square of the speed. This explains why you have to hold on when your car makes a sharp (large  $\kappa$ ), high-speed (large  $|\vec{v}|$ ) turn. If you double the speed of your car, you will experience four times the normal component of acceleration for the same curvature.

If an object moves in a circle at a constant speed,  $\frac{d^2s}{dt^2}$  is zero and all the acceleration points along  $\mathbf{N}$  toward the circle's center. If the object is speeding up or slowing down,  $\vec{a}$  has a nonzero tangential component.



To calculate  $a_N$ , we usually use the formula  $a_N = \sqrt{|\vec{a}|^2 - a_T^2}$ , which comes from solving the equation  $|\vec{a}|^2 = \vec{a} \cdot \vec{a} = a_T^2 + a_N^2$ . Using this we can find  $a_N$  without having to calculate  $\kappa$  first.

To calculate  $a_N$ , we usually use the formula  $a_N = \sqrt{|\vec{a}^2| - a_T^2}$ , which comes from solving the equation  $|\vec{a}^2| = \vec{a} \cdot \vec{a} = a_T^2 + a_N^2$ . Using this we can find  $a_N$  without having to calculate  $\kappa$  first.

$$a_N = \sqrt{|\vec{a}^2| - a_T^2}.$$

To calculate  $a_N$ , we usually use the formula  $a_N = \sqrt{|\vec{a}|^2 - a_T^2}$ , which comes from solving the equation  $|\vec{a}|^2 = \vec{a} \cdot \vec{a} = a_T^2 + a_N^2$ . Using this we can find  $a_N$  without having to calculate  $\kappa$  first.

$$a_N = \sqrt{|\vec{a}|^2 - a_T^2}.$$

Problem 1: Without finding  $\vec{T}$  and  $\vec{N}$ , write the acceleration of the motion

$$r(t) = (\cos t + t \sin t)\vec{i} + (\sin t - t \cos t)\vec{j}, t > 0$$

in the form

$$a = a_T \vec{T} + a_N \vec{N}.$$

To calculate  $a_N$ , we usually use the formula  $a_N = \sqrt{|\vec{a}|^2 - a_T^2}$ , which comes from solving the equation  $|\vec{a}|^2 = \vec{a} \cdot \vec{a} = a_T^2 + a_N^2$ . Using this we can find  $a_N$  without having to calculate  $\kappa$  first.

$$a_N = \sqrt{|\vec{a}|^2 - a_T^2}.$$

Problem 1: Without finding  $\vec{T}$  and  $\vec{N}$ , write the acceleration of the motion

$$r(t) = (\cos t + t \sin t)\vec{i} + (\sin t - t \cos t)\vec{j}, t > 0$$

in the form

$$a = a_T \vec{T} + a_N \vec{N}.$$

Sol:  $a_T = 1, a_N = t$ .

# Torsion

How does  $\frac{d\mathbf{B}}{ds}$  behave in relation to  $\mathbf{T}$ ,  $\mathbf{N}$ , and  $\mathbf{B}$ ?

$$\frac{d\mathbf{B}}{ds} = \frac{d}{ds}(\mathbf{T} \times \mathbf{N}) = \frac{d\mathbf{T}}{ds} \times \mathbf{N} + \mathbf{T} \times \frac{d\mathbf{N}}{ds} = \mathbf{T} \times \frac{d\mathbf{N}}{ds}.$$

# Torsion

How does  $\frac{d\mathbf{B}}{ds}$  behave in relation to  $\mathbf{T}$ ,  $\mathbf{N}$ , and  $\mathbf{B}$ ?

$$\frac{d\mathbf{B}}{ds} = \frac{d}{ds}(\mathbf{T} \times \mathbf{N}) = \frac{d\mathbf{T}}{ds} \times \mathbf{N} + \mathbf{T} \times \frac{d\mathbf{N}}{ds} = \mathbf{T} \times \frac{d\mathbf{N}}{ds}.$$

From this we see that  $\frac{d\mathbf{B}}{ds}$  is orthogonal to  $\mathbf{T}$ .

# Torsion

How does  $\frac{d\mathbf{B}}{ds}$  behave in relation to  $\mathbf{T}$ ,  $\mathbf{N}$ , and  $\mathbf{B}$ ?

$$\frac{d\mathbf{B}}{ds} = \frac{d}{ds}(\mathbf{T} \times \mathbf{N}) = \frac{d\mathbf{T}}{ds} \times \mathbf{N} + \mathbf{T} \times \frac{d\mathbf{N}}{ds} = \mathbf{T} \times \frac{d\mathbf{N}}{ds}.$$

From this we see that  $\frac{d\mathbf{B}}{ds}$  is orthogonal to  $\mathbf{T}$ . Since  $\mathbf{B}(\mathbf{t})$  has constant length, so  $\mathbf{B}(t)$  and  $\frac{d\mathbf{B}}{dt}$  are orthogonal and

# Torsion

How does  $\frac{d\mathbf{B}}{ds}$  behave in relation to  $\mathbf{T}$ ,  $\mathbf{N}$ , and  $\mathbf{B}$ ?

$$\frac{d\mathbf{B}}{ds} = \frac{d}{ds}(\mathbf{T} \times \mathbf{N}) = \frac{d\mathbf{T}}{ds} \times \mathbf{N} + \mathbf{T} \times \frac{d\mathbf{N}}{ds} = \mathbf{T} \times \frac{d\mathbf{N}}{ds}.$$

From this we see that  $\frac{d\mathbf{B}}{ds}$  is orthogonal to  $\mathbf{T}$ . Since  $\mathbf{B}(\mathbf{t})$  has constant length, so  $\mathbf{B}(t)$  and  $\frac{d\mathbf{B}}{dt}$  are orthogonal and

$$\frac{d\mathbf{B}}{ds} = \frac{d\mathbf{B}}{dt} \frac{dt}{ds} = \frac{d\mathbf{B}}{dt} \frac{1}{|v(t)|}.$$



# Torsion

How does  $\frac{d\mathbf{B}}{ds}$  behave in relation to  $\mathbf{T}$ ,  $\mathbf{N}$ , and  $\mathbf{B}$ ?

$$\frac{d\mathbf{B}}{ds} = \frac{d}{ds}(\mathbf{T} \times \mathbf{N}) = \frac{d\mathbf{T}}{ds} \times \mathbf{N} + \mathbf{T} \times \frac{d\mathbf{N}}{ds} = \mathbf{T} \times \frac{d\mathbf{N}}{ds}.$$

From this we see that  $\frac{d\mathbf{B}}{ds}$  is orthogonal to  $\mathbf{T}$ . Since  $\mathbf{B}(\mathbf{t})$  has constant length, so  $\mathbf{B}(t)$  and  $\frac{d\mathbf{B}}{dt}$  are orthogonal and

$$\frac{d\mathbf{B}}{ds} = \frac{d\mathbf{B}}{dt} \frac{dt}{ds} = \frac{d\mathbf{B}}{dt} \frac{1}{|v(t)|}.$$

Thus  $\frac{d\mathbf{B}}{ds}$  is also orthogonal to  $\mathbf{B}$ , it follows that  $\frac{d\mathbf{B}}{ds}$  is orthogonal to the plane of  $\mathbf{B}$  and  $\mathbf{T}$ .

# Torsion

How does  $\frac{d\mathbf{B}}{ds}$  behave in relation to  $\mathbf{T}$ ,  $\mathbf{N}$ , and  $\mathbf{B}$ ?

$$\frac{d\mathbf{B}}{ds} = \frac{d}{ds}(\mathbf{T} \times \mathbf{N}) = \frac{d\mathbf{T}}{ds} \times \mathbf{N} + \mathbf{T} \times \frac{d\mathbf{N}}{ds} = \mathbf{T} \times \frac{d\mathbf{N}}{ds}.$$

From this we see that  $\frac{d\mathbf{B}}{ds}$  is orthogonal to  $\mathbf{T}$ . Since  $\mathbf{B}(\mathbf{t})$  has constant length, so  $\mathbf{B}(t)$  and  $\frac{d\mathbf{B}}{dt}$  are orthogonal and

$$\frac{d\mathbf{B}}{ds} = \frac{d\mathbf{B}}{dt} \frac{dt}{ds} = \frac{d\mathbf{B}}{dt} \frac{1}{|v(t)|}.$$

Thus  $\frac{d\mathbf{B}}{ds}$  is also orthogonal to  $\mathbf{B}$ , it follows that  $\frac{d\mathbf{B}}{ds}$  is orthogonal to the plane of  $\mathbf{B}$  and  $\mathbf{T}$ . In other words,  $\frac{d\mathbf{B}}{ds}$  is parallel to  $\mathbf{N}$ , so  $\frac{d\mathbf{B}}{ds}$  is a scalar multiple of  $\mathbf{N}$ .

# Torsion

How does  $\frac{d\mathbf{B}}{ds}$  behave in relation to  $\mathbf{T}$ ,  $\mathbf{N}$ , and  $\mathbf{B}$ ?

$$\frac{d\mathbf{B}}{ds} = \frac{d}{ds}(\mathbf{T} \times \mathbf{N}) = \frac{d\mathbf{T}}{ds} \times \mathbf{N} + \mathbf{T} \times \frac{d\mathbf{N}}{ds} = \mathbf{T} \times \frac{d\mathbf{N}}{ds}.$$

From this we see that  $\frac{d\mathbf{B}}{ds}$  is orthogonal to  $\mathbf{T}$ . Since  $\mathbf{B}(\mathbf{t})$  has constant length, so  $\mathbf{B}(t)$  and  $\frac{d\mathbf{B}}{dt}$  are orthogonal and

$$\frac{d\mathbf{B}}{ds} = \frac{d\mathbf{B}}{dt} \frac{dt}{ds} = \frac{d\mathbf{B}}{dt} \frac{1}{|v(t)|}.$$

Thus  $\frac{d\mathbf{B}}{ds}$  is also orthogonal to  $\mathbf{B}$ , it follows that  $\frac{d\mathbf{B}}{ds}$  is orthogonal to the plane of  $\mathbf{B}$  and  $\mathbf{T}$ . In other words,  $\frac{d\mathbf{B}}{ds}$  is parallel to  $\mathbf{N}$ , so  $\frac{d\mathbf{B}}{ds}$  is a scalar multiple of  $\mathbf{N}$ . i.e

$$\frac{d\mathbf{B}}{ds} = -\tau \mathbf{N}.$$

The scalar  $\tau$  is called the torsion along the curve. Notice that


$$\frac{d\mathbf{B}}{ds} \cdot \mathbf{N} = -\tau \mathbf{N} \cdot \mathbf{N} = -\tau.$$

### Definition

Let  $\mathbf{B} = \mathbf{T} \times \mathbf{N}$ . The torsion function of a smooth curve is

$$\tau = -\frac{d\mathbf{B}}{ds} \cdot \mathbf{N}.$$

Unlike the curvature  $\kappa$ , which is never negative, the torsion  $\tau$  may be positive, negative, or zero.



**FIGURE 13.28** The names of the three planes determined by  $\mathbf{T}$ ,  $\mathbf{N}$ , and  $\mathbf{B}$ .

The curvature  $\kappa = \left| \frac{d\mathbf{T}}{ds} \right|$  can be thought of as the rate at which the normal plane turns as the point  $P$  moves along its path. Similarly, the torsion  $\tau$  is the rate at which the osculating plane turns about  $\mathbf{T}$  as  $P$  moves along the curve. Torsion measures how the curve twists.

### Vector Formula for Curvature

$$\kappa = \frac{|\mathbf{v} \times \mathbf{a}|}{|\mathbf{v}|^3}$$

### Formula for Torsion

$$\tau = \frac{\begin{vmatrix} \dot{x} & \dot{y} & \dot{z} \\ \ddot{x} & \ddot{y} & \ddot{z} \\ \dddot{x} & \dddot{y} & \dddot{z} \end{vmatrix}}{|\mathbf{v} \times \mathbf{a}|^2} \quad (\text{if } \mathbf{v} \times \mathbf{a} \neq \mathbf{0})$$

Find the curvature and torsion for the helix

$$\mathbf{r}(t) = (a \cos t)\vec{i} + (a \sin t)\vec{j} + bt\vec{k}, \quad a, b \geq 0, \quad a^2 + b^2 \neq 0$$

using the above formula.

Derive the vector formula for curvature

$$\kappa = \frac{|\mathbf{v} \times \mathbf{a}|}{|\mathbf{v}|^3}.$$

Derive

$$\tau = -\frac{1}{|\mathbf{v}|} \left( \frac{d\mathbf{B}}{dt} \cdot \mathbf{N} \right).$$