

# Double Integrals - Area and Polar Form

Devika S

Department of Mathematics  
BITS Pilani, K K Birla Goa Campus

November 4, 2024



# Examples - Double Integrals

Example 1. Reverse the order of the integration

$$\int_0^1 \int_2^{4-2x} dy dx.$$

# Examples - Double Integrals

Example 1. Reverse the order of the integration

$$\int_0^1 \int_2^{4-2x} dy dx.$$

**Solution:** First we sketch the region and find the limits of  $x$  as functions of  $y$ :

# Examples - Double Integrals

Example 1. Reverse the order of the integration

$$\int_0^1 \int_2^{4-2x} dy dx.$$

**Solution:** First we sketch the region and find the limits of  $x$  as functions of  $y$ : The limits of  $y$  are  $2 \leq y \leq 4$ .

# Examples - Double Integrals

Example 1. Reverse the order of the integration

$$\int_0^1 \int_2^{4-2x} dy dx.$$

**Solution:** First we sketch the region and find the limits of  $x$  as functions of  $y$ : The limits of  $y$  are  $2 \leq y \leq 4$ . Then the limits of  $x$  are bounded by the lines  $x = 0$  and  $x = \frac{1}{2}(4 - y)$ .

# Examples - Double Integrals

**Example 1.** Reverse the order of the integration

$$\int_0^1 \int_2^{4-2x} dy dx.$$

**Solution:** First we sketch the region and find the limits of  $x$  as functions of  $y$ : The limits of  $y$  are  $2 \leq y \leq 4$ . Then the limits of  $x$  are bounded by the lines  $x = 0$  and  $x = \frac{1}{2}(4 - y)$ .

So the reversed integral is  $\int_2^4 \int_0^{\frac{1}{2}(4-y)} dx dy$ .

# Examples - Double Integrals

**Example 1.** Reverse the order of the integration

$$\int_0^1 \int_2^{4-2x} dy dx.$$

**Solution:** First we sketch the region and find the limits of  $x$  as functions of  $y$ : The limits of  $y$  are  $2 \leq y \leq 4$ . Then the limits of  $x$  are bounded by the lines  $x = 0$  and  $x = \frac{1}{2}(4 - y)$ .

So the reversed integral is  $\int_2^4 \int_0^{\frac{1}{2}(4-y)} dx dy$ .

**Example 2.** Reverse the order of the integral in

$$\int_0^{\frac{3}{2}} \int_0^{9-4x^2} 16x dy dx.$$

# Examples - Double Integrals

**Example 1.** Reverse the order of the integration

$$\int_0^1 \int_2^{4-2x} dy dx.$$

**Solution:** First we sketch the region and find the limits of  $x$  as functions of  $y$ : The limits of  $y$  are  $2 \leq y \leq 4$ . Then the limits of  $x$  are bounded by the lines  $x = 0$  and  $x = \frac{1}{2}(4 - y)$ .

So the reversed integral is  $\int_2^4 \int_0^{\frac{1}{2}(4-y)} dx dy$ .

**Example 2.** Reverse the order of the integral in

$$\int_0^{\frac{3}{2}} \int_0^{9-4x^2} 16x dy dx.$$

**Solution:** The reversed integral is  $\int_0^9 \int_0^{\frac{\sqrt{9-y}}{2}} 16x dx dy$ .



# Integral over an unbounded region

Example 3. Compute the integral

$$\int_1^{\infty} \int_{e^{-x}}^1 \frac{1}{x^3 y} dy dx.$$

# Integral over an unbounded region

**Example 3.** Compute the integral

$$\int_1^{\infty} \int_{e^{-x}}^1 \frac{1}{x^3 y} dy dx.$$

**Solution:** Just as in the single variable case we define this integral as the limit

$$\lim_{t \rightarrow \infty} \int_1^t \int_{e^{-x}}^1 \frac{1}{x^3 y} dy dx.$$

# Integral over an unbounded region

**Example 3.** Compute the integral

$$\int_1^{\infty} \int_{e^{-x}}^1 \frac{1}{x^3 y} dy dx.$$

**Solution:** Just as in the single variable case we define this integral as the limit

$$\lim_{t \rightarrow \infty} \int_1^t \int_{e^{-x}}^1 \frac{1}{x^3 y} dy dx.$$

And the integral comes to

$$\int_1^t \frac{1}{x^2} dx = \left[ -\frac{1}{x} \right]_1^t = -\frac{1}{t} + 1.$$

# Integral over an unbounded region

**Example 3.** Compute the integral

$$\int_1^{\infty} \int_{e^{-x}}^1 \frac{1}{x^3 y} dy dx.$$

**Solution:** Just as in the single variable case we define this integral as the limit

$$\lim_{t \rightarrow \infty} \int_1^t \int_{e^{-x}}^1 \frac{1}{x^3 y} dy dx.$$

And the integral comes to

$$\int_1^t \frac{1}{x^2} dx = \left[ -\frac{1}{x} \right]_1^t = -\frac{1}{t} + 1.$$

As we pass through the limit we get 1.

# Examples

1. Sketch the region of integration, reverse the order of integration, and evaluate the integrals:

①  $\int_0^3 \int_{\sqrt{x/3}}^1 e^{y^3} dy dx \quad (e - 1)$


②  $\int_0^1 \int_y^1 x^2 e^{xy} dx dy \quad ((e - 2)/2)$

2. Sketch the region of integration and evaluate  $\iint_R (y - 2x^2) dA$  where  $R$  is the region bounded by  $|x| + |y| = 1$ .  $(-2/3)$

3. Compute the integral:  $\int_0^\infty \int_0^\infty x e^{-(x+2y)} dx dy \quad (1/2)$


# Examples

4. Evaluate  $\iint_R (3 - 6xy) dA$  where  $D$  is the region shown below:



(36 - Split the integral up and do the actual integration over separate sub regions.)

5. Evaluate  $\iint_D e^{y^4} dA$  where  $D$  is the region shown below:



$$\left(\frac{e-1}{2}\right)$$

## Examples

6. Find the volume of the solid bounded by the planes  $x = 0$ ,  $y = 0$ ,  $z = 0$ , and  $2x + 3y + z = 6$ . (6 cubic units - the solid is a tetrahedron with the base on the  $xy$  plane and a height  $z = 6 - 2x - 3y$ . The base is the region bounded by the lines,  $x = 0$ ,  $y = 0$ , and  $2x + 3y = 6$  where  $z = 0$ .)

7. Find the volume of the first octant part of the solid bounded by the cylinders  $x^2 + y^2 = 1$  and  $y^2 + z^2 = 1$ .

$$\left( \int_0^1 \int_0^{\sqrt{1-y^2}} \sqrt{1-y^2} \, dx dy = 2/3 \right)$$

8. Find the region  $R$  on the  $xy$  plane that maximizes the integral

$$\iint_R (4 - x^2 - y^2) dy dx.$$

(The region must be  $x^2 + y^2 \leq 4$  - justify using the monotonicity property of double integral).

# Area of Bounded Regions

## Area

Area of a closed, bounded plane region  $R$  is

$$\text{Area of } R = \iint_R dA.$$



# Area of Bounded Regions

## Area

Area of a closed, bounded plane region  $R$  is

$$\text{Area of } R = \iint_R dA.$$

## Average

Average value of  $f$  over  $R$  is

$$\frac{1}{\text{Area of } R} \iint_R f dA.$$

## Examples

**Example 1:** Find the average value of the function  $f(x, y) = 7xy^2$  on the region  $R$  bounded by the line  $x = y$  and the curve  $x = \sqrt{y}$ .

# Examples

**Example 1:** Find the average value of the function  $f(x, y) = 7xy^2$  on the region  $R$  bounded by the line  $x = y$  and the curve  $x = \sqrt{y}$ .

$$\text{Area of } R = \iint_R dA = \int_0^1 \int_y^{\sqrt{y}} dx dy = 1/6.$$

## Examples

**Example 1:** Find the average value of the function  $f(x, y) = 7xy^2$  on the region  $R$  bounded by the line  $x = y$  and the curve  $x = \sqrt{y}$ .

$$\text{Area of } R = \iint_R dA = \int_0^1 \int_y^{\sqrt{y}} dx dy = 1/6.$$

Hence, average is

$$\frac{1}{1/6} \iint_R f dA = \frac{1}{1/6} \int_0^1 \int_y^{\sqrt{y}} 7xy^2 dx dy = 21/20.$$

## Examples

**Example 1:** Find the average value of the function  $f(x, y) = 7xy^2$  on the region  $R$  bounded by the line  $x = y$  and the curve  $x = \sqrt{y}$ .

$$\text{Area of } R = \iint_R dA = \int_0^1 \int_y^{\sqrt{y}} dx dy = 1/6.$$

Hence, average is

$$\frac{1}{1/6} \iint_R f dA = \frac{1}{1/6} \int_0^1 \int_y^{\sqrt{y}} 7xy^2 dx dy = 21/20.$$

**Example 2.** Sketch the region bounded by the curve  $y = e^x$ , and the lines  $y = 0$ ,  $x = 0$ , and  $x = \ln 2$ . Then express the region's area as an iterated double integral and evaluate the integral. (1)

**Example 3.** Find the average height of the paraboloid  $z = x^2 + y^2$  over the square  $0 \leq x \leq 2$ ,  $0 \leq y \leq 2$ . (8/3)


# Integral of Function of Polar Coordinates

Suppose that a function  $f(r, \theta)$  is defined in polar coordinates over the region  $R$  which lies between the angles  $\alpha \leq \theta \leq \beta$  and the radius bounded as  $g_1(\theta) \leq r \leq g_2(\theta)$  where  $g_1(\theta), g_2(\theta)$  are continuous functions.

# Integral of Function of Polar Coordinates

Suppose that a function  $f(r, \theta)$  is defined in polar coordinates over the region  $R$  which lies between the angles  $\alpha \leq \theta \leq \beta$  and the radius bounded as  $g_1(\theta) \leq r \leq g_2(\theta)$  where  $g_1(\theta), g_2(\theta)$  are continuous functions.

Suppose also that  $0 \leq g_1(\theta) \leq g_2(\theta) \leq a$  for every value of  $\theta$  between  $\alpha$  and  $\beta$ . Then  $R$  lies in a fan shaped region  $Q$  defined by the inequality  $0 \leq r \leq a$  and  $\alpha \leq \theta \leq \beta$ .



# Integral of Function of Polar Coordinates

We cover  $Q$  by a grid of circular arcs and rays. The arcs are cut from circles centered at the origin, with radii  $\Delta r, 2\Delta r, \dots, m\Delta r$ , where  $\Delta r = a/m$ . The rays are given by

$$\theta = \alpha, \theta = \alpha + \Delta\theta, \dots, \theta = \alpha + m'\Delta\theta = \beta,$$

where  $\Delta\theta = (\beta - \alpha)/m'$ .



# Integral of Function of Polar Coordinates

We cover  $Q$  by a grid of circular arcs and rays. The arcs are cut from circles centered at the origin, with radii  $\Delta r, 2\Delta r, \dots, m\Delta r$ , where  $\Delta r = a/m$ . The rays are given by

$$\theta = \alpha, \theta = \alpha + \Delta\theta, \dots, \theta = \alpha + m'\Delta\theta = \beta,$$

where  $\Delta\theta = (\beta - \alpha)/m'$ .

The arcs and rays partition  $Q$  into small patches called “polar rectangles”. We number the polar rectangles that lies inside  $R$ , calling their areas  $\Delta A_1, \Delta A_2, \dots, \Delta A_n$ . We let  $(r_k, \theta_k)$  be any point in the polar rectangles whose area is  $\Delta A_k$ . We then form the sum

$$S_n = \sum_{k=1}^n f(r_k, \theta_k) \Delta A_k.$$

# Integral of Function of Polar Coordinates

We cover  $Q$  by a grid of circular arcs and rays. The arcs are cut from circles centered at the origin, with radii  $\Delta r, 2\Delta r, \dots, m\Delta r$ , where  $\Delta r = a/m$ . The rays are given by

$$\theta = \alpha, \theta = \alpha + \Delta\theta, \dots, \theta = \alpha + m'\Delta\theta = \beta,$$

where  $\Delta\theta = (\beta - \alpha)/m'$ .

The arcs and rays partition  $Q$  into small patches called “polar rectangles”. We number the polar rectangles that lies inside  $R$ , calling their areas  $\Delta A_1, \Delta A_2, \dots, \Delta A_n$ . We let  $(r_k, \theta_k)$  be any point in the polar rectangles whose area is  $\Delta A_k$ . We then form the sum

$$S_n = \sum_{k=1}^n f(r_k, \theta_k) \Delta A_k.$$

If  $f$  is continuous on  $R$ , this sum will approach a limit as we refine the grid to make  $\Delta r$  and  $\Delta\theta$  go to zero. The limit is called the double integral of  $f$  over  $R$ .

# Integral of Function of Polar Coordinates

In symbol

$$\iint_R f(r, \theta) dA = \lim_{n \rightarrow \infty} S_n.$$

# Integral of Function of Polar Coordinates

In symbol

$$\iint_R f(r, \theta) dA = \lim_{n \rightarrow \infty} S_n.$$

We note that  $\Delta A_k = r_k \Delta r \Delta \theta$ , therefore

$$S_n = \sum_{k=1}^n f(r_k, \theta_k) r_k \Delta r \Delta \theta \text{ and}$$

$$\iint_R f(r, \theta) dA = \lim_{n \rightarrow \infty} S_n = \iint_R f(r, \theta) r \, dr \, d\theta.$$

# Integral of Function of Polar Coordinates

In symbol

$$\iint_R f(r, \theta) dA = \lim_{n \rightarrow \infty} S_n.$$

We note that  $\Delta A_k = r_k \Delta r \Delta \theta$ , therefore

$$S_n = \sum_{k=1}^n f(r_k, \theta_k) r_k \Delta r \Delta \theta \text{ and}$$

$$\iint_R f(r, \theta) dA = \lim_{n \rightarrow \infty} S_n = \iint_R f(r, \theta) r \, dr \, d\theta.$$

A version of Fubini's theorem says that

$$\iint_R f(r, \theta) dA = \int_{\theta=\alpha}^{\theta=\beta} \int_{r=g_1(\theta)}^{r=g_2(\theta)} f(r, \theta) r \, dr \, d\theta.$$

# How to find the limits of integrals in polar coordinates

- 1 First sketch the region given and label the bounding curves.

# How to find the limits of integrals in polar coordinates

- 1 First sketch the region given and label the bounding curves.
- 2 Imagine a ray from the origin entering and leaving the region, the point of entrance as a function of  $\theta$  is the lower limit and the point of departure as a function of  $\theta$  is the upper limit.

# How to find the limits of integrals in polar coordinates

- 1 First sketch the region given and label the bounding curves.
- 2 Imagine a ray from the origin entering and leaving the region, the point of entrance as a function of  $\theta$  is the lower limit and the point of departure as a function of  $\theta$  is the upper limit.
- 3 Find the extreme angles of the points of the region, the smallest is the lower limit, the largest is the upper limit of the integral.



# How to find the limits of integrals in polar coordinates

- 1 First sketch the region given and label the bounding curves.
- 2 Imagine a ray from the origin entering and leaving the region, the point of entrance as a function of  $\theta$  is the lower limit and the point of departure as a function of  $\theta$  is the upper limit.
- 3 Find the extreme angles of the points of the region, the smallest is the lower limit, the largest is the upper limit of the integral.

## Changing Integrals from Cartesian to Polar Coordinates:

- 1 First substitute  $x = r \cos \theta, y = r \sin \theta$ .

# How to find the limits of integrals in polar coordinates

- 1 First sketch the region given and label the bounding curves.
- 2 Imagine a ray from the origin entering and leaving the region, the point of entrance as a function of  $\theta$  is the lower limit and the point of departure as a function of  $\theta$  is the upper limit.
- 3 Find the extreme angles of the points of the region, the smallest is the lower limit, the largest is the upper limit of the integral.

## Changing Integrals from Cartesian to Polar Coordinates:

- 1 First substitute  $x = r \cos \theta, y = r \sin \theta$ .
- 2 Then substitute  $dydx$  with  $r drd\theta$ .

# How to find the limits of integrals in polar coordinates

- 1 First sketch the region given and label the bounding curves.
- 2 Imagine a ray from the origin entering and leaving the region, the point of entrance as a function of  $\theta$  is the lower limit and the point of departure as a function of  $\theta$  is the upper limit.
- 3 Find the extreme angles of the points of the region, the smallest is the lower limit, the largest is the upper limit of the integral.

## Changing Integrals from Cartesian to Polar Coordinates:

- 1 First substitute  $x = r \cos \theta, y = r \sin \theta$ .
- 2 Then substitute  $dydx$  with  $r drd\theta$ .
- 3 Sketch the region  $R$  of integration and then find the polar limits of the region from the sketch using the above method.

# Examples

**Example 1.** Find the integral by changing to polar coordinates:

$$\int_0^6 \int_0^y x dx dy.$$

## Examples

**Example 1.** Find the integral by changing to polar coordinates:

$$\int_0^6 \int_0^y x dx dy.$$

**Solution:** First let us sketch the region of integration. In the  $xy$ -plane, this region is the triangular region bounded by the line  $y = x$ , the line  $y = 6$ , and the  $x$ -axis.

# Examples

**Example 1.** Find the integral by changing to polar coordinates:

$$\int_0^6 \int_0^y x dx dy.$$

**Solution:** First let us sketch the region of integration. In the  $xy$ -plane, this region is the triangular region bounded by the line  $y = x$ , the line  $y = 6$ , and the  $x$ -axis.

The angle  $\theta$  is in the domain  $\frac{\pi}{4} \leq \theta \leq \frac{\pi}{2}$ . The radius is bounded below by 0 and above by  $\frac{6}{\sin \theta}$ .

## Examples

**Example 1.** Find the integral by changing to polar coordinates:

$$\int_0^6 \int_0^y x dx dy.$$

**Solution:** First let us sketch the region of integration. In the  $xy$ -plane, this region is the triangular region bounded by the line  $y = x$ , the line  $y = 6$ , and the  $x$ -axis.

The angle  $\theta$  is in the domain  $\frac{\pi}{4} \leq \theta \leq \frac{\pi}{2}$ . The radius is bounded below by 0 and above by  $\frac{6}{\sin \theta}$ . So the integral becomes:

$$\begin{aligned} \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \int_0^{\frac{6}{\sin \theta}} r^2 \cos \theta dr d\theta &= 72 \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \frac{\cos \theta d\theta}{\sin^3 \theta} = 72 \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \cot \theta \operatorname{cosec}^2 \theta d\theta \\ &= -36 \left[ \cot^2 \theta \right]_{\frac{\pi}{4}}^{\frac{\pi}{2}} = 36. \end{aligned}$$

# Examples

**Example 2.** Find the integral by changing to the polar coordinates:

$$\int_0^\infty \int_0^\infty e^{-(x^2+y^2)} dx dy.$$



# Examples

**Example 2.** Find the integral by changing to the polar coordinates:

$$\int_0^{\infty} \int_0^{\infty} e^{-(x^2+y^2)} dx dy.$$

**Solution:** The region is clearly the first quadrant. So the limits of the angle is  $0 \leq \theta \leq \frac{\pi}{2}$ . The limits of the radius is clearly between 0 and  $\infty$ .

# Examples

**Example 2.** Find the integral by changing to the polar coordinates:

$$\int_0^{\infty} \int_0^{\infty} e^{-(x^2+y^2)} dx dy.$$

**Solution:** The region is clearly the first quadrant. So the limits of the angle is  $0 \leq \theta \leq \frac{\pi}{2}$ . The limits of the radius is clearly between 0 and  $\infty$ . So the integral becomes:

$$\begin{aligned} \int_0^{\frac{\pi}{2}} \int_0^{\infty} e^{-r^2} r dr d\theta &= 1/2 \int_0^{\frac{\pi}{2}} \lim_{t \rightarrow \infty} \left[ -e^{-r^2} \right]_0^t d\theta \\ &= \frac{\pi}{4}. \end{aligned}$$

# Examples

**Example 2.** Find the integral by changing to the polar coordinates:

$$\int_0^{\infty} \int_0^{\infty} e^{-(x^2+y^2)} dx dy.$$

**Solution:** The region is clearly the first quadrant. So the limits of the angle is  $0 \leq \theta \leq \frac{\pi}{2}$ . The limits of the radius is clearly between 0 and  $\infty$ . So the integral becomes:

$$\begin{aligned} \int_0^{\frac{\pi}{2}} \int_0^{\infty} e^{-r^2} r dr d\theta &= 1/2 \int_0^{\frac{\pi}{2}} \lim_{t \rightarrow \infty} \left[ -e^{-r^2} \right]_0^t d\theta \\ &= \frac{\pi}{4}. \end{aligned}$$

Try the integral in Cartesian coordinate form.

# Examples

**Example 2.** Find the integral by changing to the polar coordinates:

$$\int_0^{\infty} \int_0^{\infty} e^{-(x^2+y^2)} dx dy.$$

**Solution:** The region is clearly the first quadrant. So the limits of the angle is  $0 \leq \theta \leq \frac{\pi}{2}$ . The limits of the radius is clearly between 0 and  $\infty$ . So the integral becomes:

$$\begin{aligned} \int_0^{\frac{\pi}{2}} \int_0^{\infty} e^{-r^2} r dr d\theta &= 1/2 \int_0^{\frac{\pi}{2}} \lim_{t \rightarrow \infty} \left[ -e^{-r^2} \right]_0^t d\theta \\ &= \frac{\pi}{4}. \end{aligned}$$

Try the integral in Cartesian coordinate form. This shows the power of polar coordinates in dealing with certain types of functions.

# Examples

**Example 2.** Find the integral by changing to the polar coordinates:

$$\int_0^{\infty} \int_0^{\infty} e^{-(x^2+y^2)} dx dy.$$

**Solution:** The region is clearly the first quadrant. So the limits of the angle is  $0 \leq \theta \leq \frac{\pi}{2}$ . The limits of the radius is clearly between 0 and  $\infty$ . So the integral becomes:

$$\begin{aligned} \int_0^{\frac{\pi}{2}} \int_0^{\infty} e^{-r^2} r dr d\theta &= 1/2 \int_0^{\frac{\pi}{2}} \lim_{t \rightarrow \infty} \left[ -e^{-r^2} \right]_0^t d\theta \\ &= \frac{\pi}{4}. \end{aligned}$$

Try the integral in Cartesian coordinate form. This shows the power of polar coordinates in dealing with certain types of functions. Namely the **radially symmetric functions**.

# Area in Polar Coordinates

## Area in Polar

The area of a region  $R$  in polar coordinates is defined as

$$\iint_R r dr d\theta.$$

# Area in Polar Coordinates

## Area in Polar

The area of a region  $R$  in polar coordinates is defined as

$$\iint_R r dr d\theta.$$

**Example 1.** Find the area of one leaf of the rose  $r = 12 \cos 3\theta$ .

# Area in Polar Coordinates


## Area in Polar

The area of a region  $R$  in polar coordinates is defined as

$$\iint_R r dr d\theta.$$

**Example 1.** Find the area of one leaf of the rose  $r = 12 \cos 3\theta$ .

**Solution:**





# Examples

We get  $-\frac{\pi}{6} \leq \theta \leq \frac{\pi}{6}$ .

# Examples

We get  $-\frac{\pi}{6} \leq \theta \leq \frac{\pi}{6}$ . And clearly the limits of the radius is  $0 \leq r \leq 12 \cos 3\theta$ .

# Examples

We get  $-\frac{\pi}{6} \leq \theta \leq \frac{\pi}{6}$ . And clearly the limits of the radius is  $0 \leq r \leq 12 \cos 3\theta$ .

So the area is given by the integral:

$$\begin{aligned} \int_{-\frac{\pi}{6}}^{\frac{\pi}{6}} \int_0^{12 \cos 3\theta} r dr d\theta &= 2 \frac{12^2}{2} \int_0^{\frac{\pi}{6}} \cos^2 3\theta d\theta \\ &= 2 \frac{12^2}{4} \int_0^{\frac{\pi}{6}} (1 + \cos 6\theta) d\theta = 12\pi. \end{aligned}$$

# Examples

We get  $-\frac{\pi}{6} \leq \theta \leq \frac{\pi}{6}$ . And clearly the limits of the radius is  $0 \leq r \leq 12 \cos 3\theta$ .

So the area is given by the integral:

$$\begin{aligned} \int_{-\frac{\pi}{6}}^{\frac{\pi}{6}} \int_0^{12 \cos 3\theta} r dr d\theta &= 2 \frac{12^2}{2} \int_0^{\frac{\pi}{6}} \cos^2 3\theta d\theta \\ &= 2 \frac{12^2}{4} \int_0^{\frac{\pi}{6}} (1 + \cos 6\theta) d\theta = 12\pi. \end{aligned}$$