

# MATHEMATICS-I

ANUSHAYA MOHAPATRA

Department of Mathematics  
BITS PILANI K K Birla Goa Campus, Goa

September 5, 2024

# Lecture 12

## Power series

# Power series

- A **power series about**  $x = a$  is a series of the form

$$\sum_{n=0}^{\infty} c_n(x-a)^n = c_0 + c_1(x-a) + c_2(x-a)^2 + \cdots. \quad (0.1)$$

# Power series

- A **power series about**  $x = a$  is a series of the form

$$\sum_{n=0}^{\infty} c_n(x-a)^n = c_0 + c_1(x-a) + c_2(x-a)^2 + \cdots. \quad (0.1)$$

Here the **center**  $a$  and the **coefficients**

$c_0, c_1, \dots, c_n, \dots$  are constants.

- A **power series about**  $x = 0$  is a series of the form

$$\sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 x + c_2 x^2 + c_3 x^3 + \cdots. \quad (0.2)$$

# Power series

- A **power series about**  $x = a$  is a series of the form

$$\sum_{n=0}^{\infty} c_n(x-a)^n = c_0 + c_1(x-a) + c_2(x-a)^2 + \cdots. \quad (0.1)$$

Here the **center**  $a$  and the **coefficients**

$c_0, c_1, \dots, c_n, \dots$  are constants.

- A **power series about**  $x = 0$  is a series of the form

$$\sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 x + c_2 x^2 + c_3 x^3 + \cdots. \quad (0.2)$$

# Examples

- Geometric series:  $\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + x^3 + \dots$ .

# Examples

- Geometric series:  $\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + x^3 + \dots$ .

The series converges to  $\frac{1}{1-x}$  for all  $-1 < x < 1$  and diverges if  $|x| > 1$ . At  $x = \pm 1$ , the series clearly diverges.

# Examples

- Geometric series:  $\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + x^3 + \dots$ .

The series converges to  $\frac{1}{1-x}$  for all  $-1 < x < 1$  and diverges if  $|x| > 1$ . At  $x = \pm 1$ , the series clearly diverges.

- $\sum_{n=0}^{\infty} \left(-\frac{1}{2}\right)^n (x-2)^n = 1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 - \dots$ .



## Examples

- Geometric series:  $\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + x^3 + \dots$ .

The series converges to  $\frac{1}{1-x}$  for all  $-1 < x < 1$  and diverges if  $|x| > 1$ . At  $x = \pm 1$ , the series clearly diverges.

- $\sum_{n=0}^{\infty} \left(-\frac{1}{2}\right)^n (x-2)^n = 1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 - \dots$ .

The series converges when  $|x-2| < 2$ . In that case, the series converges to  $f(x) = \frac{1}{1 - \frac{-1}{2}(x-2)} = \frac{2}{x}$ .

# Examples

$$\bullet \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots .$$

## Examples

- $$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots .$$

Apply Ratio test:  $\left| \frac{u_{n+1}}{u_n} \right| = \frac{|x|}{1 + 1/n} \rightarrow |x|.$

## Examples

- $$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots .$$

Apply Ratio test:  $\left| \frac{u_{n+1}}{u_n} \right| = \frac{|x|}{1 + 1/n} \rightarrow |x|.$

- By ratio test, if  $|x| < 1$  the the series converges.

## Examples

- $$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots .$$

Apply Ratio test:  $\left| \frac{u_{n+1}}{u_n} \right| = \frac{|x|}{1 + 1/n} \rightarrow |x|.$

- By ratio test, if  $|x| < 1$  the the series converges.
- If  $|x| > 1$ , then  $\{(-1)^{n-1} \frac{x^n}{n}\}$  is divergent and hence the series diverges.

## Examples

- $$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots .$$

Apply Ratio test:  $\left| \frac{u_{n+1}}{u_n} \right| = \frac{|x|}{1 + 1/n} \rightarrow |x|.$

- By ratio test, if  $|x| < 1$  the the series converges.
- If  $|x| > 1$ , then  $\{(-1)^{n-1} \frac{x^n}{n}\}$  is divergent and hence the series diverges.
- For  $x = 1$ , the series is an alternating harmonic series and hence it converges.

## Examples

- $$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$$

Apply Ratio test:  $\left| \frac{u_{n+1}}{u_n} \right| = \frac{|x|}{1 + 1/n} \rightarrow |x|.$

- By ratio test, if  $|x| < 1$  the the series converges.
- If  $|x| > 1$ , then  $\{(-1)^{n-1} \frac{x^n}{n}\}$  is divergent and hence the series diverges.
- For  $x = 1$ , the series is an alternating harmonic series and hence it converges.
- For  $x = -1$  the series becomes negative of harmonic series hence it diverges.

# Examples

Find the values of  $x$  where the following series converges and diverges.

• (a)  $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{2n-1}}{2n-1}$ , (b)  $\sum_{n=0}^{\infty} \frac{(n+1)(x-2)^n}{(2n+1)!}$

(c)  $\sum_{n=0}^{\infty} n! x^n$ .



# Convergence of Power Series

## Theorem 0.1.

*If the power series  $\sum_{n=0}^{\infty} a_n x^n$  converges at  $x = c \neq 0$ , then it converges absolutely for all  $x$  with  $|x| < |c|$ . If the series diverges at  $x = d$ , then it diverges for all  $x$  with  $|x| > |d|$ .*

# Convergence of Power Series

## Theorem 0.1.

*If the power series  $\sum_{n=0}^{\infty} a_n x^n$  converges at  $x = c \neq 0$ , then it converges absolutely for all  $x$  with  $|x| < |c|$ . If the series diverges at  $x = d$ , then it diverges for all  $x$  with  $|x| > |d|$ .*

**Proof:** Suppose that the series converges for  $x = c$  that is  $\sum_{n=0}^{\infty} a_n c^n$  converges.

# Convergence of Power Series

## Theorem 0.1.

*If the power series  $\sum_{n=0}^{\infty} a_n x^n$  converges at  $x = c \neq 0$ , then it converges absolutely for all  $x$  with  $|x| < |c|$ . If the series diverges at  $x = d$ , then it diverges for all  $x$  with  $|x| > |d|$ .*

**Proof:** Suppose that the series converges for  $x = c$  that is  $\sum_{n=0}^{\infty} a_n c^n$  converges.

It implies  $\lim_{n \rightarrow \infty} a_n c^n = 0$  by  $n$ -th term test. Hence there exists a positive integer  $N$  such that  $|a_n c^n| < 1$  for all  $n > N$ .

# Convergence of Power Series

We have that  $|a_n| < \frac{1}{|c|^n}$  for  $n > N$ .

# Convergence of Power Series

We have that  $|a_n| < \frac{1}{|c|^n}$  for  $n > N$ .

Now for any  $|x| < |c|$ , we see that

$$|a_n x^n| = |a_n| |x|^n < \frac{|x|^n}{|c|^n} = |x/c|^n \text{ for } n > N.$$

## Convergence of Power Series

We have that  $|a_n| < \frac{1}{|c|^n}$  for  $n > N$ .

Now for any  $|x| < |c|$ , we see that

$$|a_n x^n| = |a_n| |x|^n < \frac{|x|^n}{|c|^n} = |x/c|^n \text{ for } n > N.$$

Since  $|x/c| < 1$ , the geometric series  $\sum_{n=0}^{\infty} |x/c|^n$  converges, and hence by comparison test, the series  $\sum_{n=0}^{\infty} |a_n x^n|$  converges.

## Convergence of Power Series

We have that  $|a_n| < \frac{1}{|c|^n}$  for  $n > N$ .

Now for any  $|x| < |c|$ , we see that

$$|a_n x^n| = |a_n| |x|^n < \frac{|x|^n}{|c|^n} = |x/c|^n \text{ for } n > N.$$

Since  $|x/c| < 1$ , the geometric series  $\sum_{n=0}^{\infty} |x/c|^n$  converges, and hence by comparison test, the series  $\sum_{n=0}^{\infty} |a_n x^n|$  converges.

Therefore the  $\sum_{n=0}^{\infty} a_n x^n$  converges absolutely for  $|x| < |c|$ .

# Convergence of Power Series

Now suppose that the series  $\sum_{n=0}^{\infty} a_n x^n$  diverges for  $x = d$ . We have to prove that the series  $\sum_n a_n x^n$  diverges for  $|x| > |d|$ .



## Convergence of Power Series

Now suppose that the series  $\sum_{n=0}^{\infty} a_n x^n$  diverges for  $x = d$ . We have to prove that the series  $\sum_n a_n x^n$  diverges for  $|x| > |d|$ .

Proof by contradiction:

# Convergence of Power Series

Now suppose that the series  $\sum_{n=0}^{\infty} a_n x^n$  diverges for  $x = d$ . We have to prove that the series  $\sum_n a_n x^n$  diverges for  $|x| > |d|$ .

**Proof by contradiction:** Suppose that for any  $x$  with  $|x| > |d|$ , the series  $\sum_{n=0}^{\infty} a_n x^n$  is convergent. .

## Convergence of Power Series

Now suppose that the series  $\sum_{n=0}^{\infty} a_n x^n$  diverges for  $x = d$ . We have to prove that the series  $\sum_n a_n x^n$  diverges for  $|x| > |d|$ .

**Proof by contradiction:** Suppose that for any  $x$  with  $|x| > |d|$ , the series  $\sum_{n=0}^{\infty} a_n x^n$  is convergent. .

Then by first part, the series  $\sum_{n=0}^{\infty} a_n d^n$  is convergent which contradicts the hypothesis.

## Convergence of Power Series

Now suppose that the series  $\sum_{n=0}^{\infty} a_n x^n$  diverges for  $x = d$ . We have to prove that the series  $\sum_n a_n x^n$  diverges for  $|x| > |d|$ .

**Proof by contradiction:** Suppose that for any  $x$  with  $|x| > |d|$ , the series  $\sum_{n=0}^{\infty} a_n x^n$  is convergent. .

Then by first part, the series  $\sum_{n=0}^{\infty} a_n d^n$  is convergent which contradicts the hypothesis.

Therefore, the series  $\sum_n a_n x^n$  is not convergent (diverges) for any  $|x| > |d|$ .

# Convergence of Power Series

- 1 The above theorem deals with convergence of power series of the form  $\sum_{n=0}^{\infty} a_n x^n$ . For the series of the form  $\sum_{n=0}^{\infty} a_n (x - a)^n$ , we can replace  $(x - a)$  by  $t$  and apply the results to the series  $\sum_{n=0}^{\infty} a_n t^n$ .

# Convergence of Power Series

- 1 The above theorem deals with convergence of power series of the form  $\sum_{n=0}^{\infty} a_n x^n$ . For the series of the form  $\sum_{n=0}^{\infty} a_n (x - a)^n$ , we can replace  $(x - a)$  by  $t$  and apply the results to the series  $\sum_{n=0}^{\infty} a_n t^n$ .
- 2 If the power series  $\sum_{n=0}^{\infty} a_n (x - a)^n$  converges for some  $x = c$  then it converges absolutely for all  $x$  such that  $|x - a| < |c - a|$ .

# Convergence of Power Series

- 1 The above theorem deals with convergence of power series of the form  $\sum_{n=0}^{\infty} a_n x^n$ . For the series of the form  $\sum_{n=0}^{\infty} a_n (x - a)^n$ , we can replace  $(x - a)$  by  $t$  and apply the results to the series  $\sum_{n=0}^{\infty} a_n t^n$ .
- 2 If the power series  $\sum_{n=0}^{\infty} a_n (x - a)^n$  converges for some  $x = c$  then it converges absolutely for all  $x$  such that  $|x - a| < |c - a|$ .
- 3 If the power series  $\sum_{n=0}^{\infty} a_n (x - a)^n$  diverges for some  $x = d$  then it diverges for all  $x$  such that  $|x - a| > |x - d|$ .