

SIMPLE HARMONIC MOTION

SUPERPOSITIONS

SIMPLE HARMONIC MOTION

SUPERPOSITIONS

- 1 **One-dimensional Superposition**
 - Equal frequency
- 2 **Multiple vibrations of same frequency**
 - Beats
- 3 **Perpendicular Vibrations: Lissajous Figures**

Superposed Vibrations in One dimension

- Most physical situations involve combined vibrations

Superposed Vibrations in One dimension


- Most physical situations involve combined vibrations
- As long as the system is **linear**
i.e. displacement \propto force,
harmonic vibrations can simply be added, mathematically!

Superposed Vibrations in One dimension

- Most physical situations involve combined vibrations
- As long as the system is **linear**
i.e. displacement \propto force,
harmonic vibrations can simply be added, mathematically!

SUPERPOSITION


Two SHM's: same frequency, different phase



$$x_1 = A_1 \cos(\omega t + \phi_1);$$

$$x_2 = A_2 \cos(\omega t + \phi_2)$$

Two SHM's: same frequency, different phase




$$x_1 = A_1 \cos(\omega t + \phi_1);$$

$$x_2 = A_2 \cos(\omega t + \phi_2)$$

Adding, we get the wave

$$x_1 + x_2 = R \cos(\omega t + \theta)$$

Two SHM's: same frequency, different phase




$$x_1 = A_1 \cos(\omega t + \phi_1);$$

$$x_2 = A_2 \cos(\omega t + \phi_2)$$

Adding, we get the wave


$$x_1 + x_2 = R \cos(\omega t + \theta)$$



Resultant wave obtained by simply adding the component waves point by point!

Calculation of R and θ


Rotating vector picture



Adding the vectors \vec{r}_1 and \vec{r}_2 ,

Calculation of R and θ

Rotating vector picture




Adding the vectors \vec{r}_1 and \vec{r}_2 ,

$$R^2 = A_1^2 + A_2^2 + 2A_1A_2 \cos \delta.$$

Calculation of R and θ

Rotating vector picture




Adding the vectors \vec{r}_1 and \vec{r}_2 ,

$$R^2 = A_1^2 + A_2^2 + 2A_1A_2 \cos \delta.$$

$$\tan \theta = \frac{A_1 \sin \phi_1 + A_2 \sin \phi_2}{A_1 \cos \phi_1 + A_2 \cos \phi_2},$$

Calculation of R and θ

Rotating vector picture



Adding the vectors \vec{r}_1 and \vec{r}_2 ,

$$R^2 = A_1^2 + A_2^2 + 2A_1A_2 \cos \delta.$$

$$\tan \theta = \frac{A_1 \sin \phi_1 + A_2 \sin \phi_2}{A_1 \cos \phi_1 + A_2 \cos \phi_2},$$


When $A_2 = A_1 = A$,

$$\theta = \frac{\delta}{2},$$

$$R = 2A \cos(\delta/2).$$

Calculation of R and θ

Rotating vector picture



Adding the vectors \vec{r}_1 and \vec{r}_2 ,

$$R^2 = A_1^2 + A_2^2 + 2A_1A_2 \cos \delta.$$

$$\tan \theta = \frac{A_1 \sin \phi_1 + A_2 \sin \phi_2}{A_1 \cos \phi_1 + A_2 \cos \phi_2},$$

When $A_2 = A_1 = A$,

$$\theta = \frac{\delta}{2},$$

$$R = 2A \cos(\delta/2).$$

Eg: Interference

Calculation of R and θ

Complex Exponential Method

Calculation of R and θ

Complex Exponential Method

$$z_1 = A_1 e^{i(\omega t + \phi_1)}$$

Calculation of R and θ

Complex Exponential Method

$$z_1 = A_1 e^{i(\omega t + \phi_1)}$$

$$z_2 = A_2 e^{i(\omega t + \phi_2)}$$

Calculation of R and θ

Complex Exponential Method

$$z_1 = A_1 e^{i(\omega t + \phi_1)}$$

$$z_2 = A_2 e^{i(\omega t + \phi_2)}$$

$$z = z_1 + z_2 = A_1 e^{i(\omega t + \phi_1)} + A_2 e^{i(\omega t + \phi_2)}$$

Calculation of R and θ

Complex Exponential Method

$$z_1 = A_1 e^{i(\omega t + \phi_1)}$$

$$z_2 = A_2 e^{i(\omega t + \phi_2)}$$

$$z = z_1 + z_2 = A_1 e^{i(\omega t + \phi_1)} + A_2 e^{i(\omega t + \phi_2)}$$

$$\text{or, } R e^{i\omega t + \theta} = e^{i(\omega t + \phi_1)} (A_1 + A_2 e^{i\delta})$$

Calculation of R and θ

Complex Exponential Method

$$z_1 = A_1 e^{i(\omega t + \phi_1)}$$

$$z_2 = A_2 e^{i(\omega t + \phi_2)}$$

$$z = z_1 + z_2 = A_1 e^{i(\omega t + \phi_1)} + A_2 e^{i(\omega t + \phi_2)}$$

$$\text{or, } R e^{i\omega t + \theta} = e^{i(\omega t + \phi_1)} (A_1 + A_2 e^{i\delta})$$

$$\text{where } R = \sqrt{(Re(z))^2 + (Im(z))^2},$$

Calculation of R and θ

Complex Exponential Method

$$z_1 = A_1 e^{i(\omega t + \phi_1)}$$

$$z_2 = A_2 e^{i(\omega t + \phi_2)}$$

$$z = z_1 + z_2 = A_1 e^{i(\omega t + \phi_1)} + A_2 e^{i(\omega t + \phi_2)}$$

$$\text{or, } Re^{i\omega t + \theta} = e^{i(\omega t + \phi_1)} (A_1 + A_2 e^{i\delta})$$

$$\text{where } R = \sqrt{(Re(z))^2 + (Im(z))^2},$$

$$\tan \theta = \frac{Im(z)}{Re(z)}$$

Calculation of R and θ

Complex Exponential Method

$$z_1 = A_1 e^{i(\omega t + \phi_1)}$$

$$z_2 = A_2 e^{i(\omega t + \phi_2)}$$

$$z = z_1 + z_2 = A_1 e^{i(\omega t + \phi_1)} + A_2 e^{i(\omega t + \phi_2)}$$

$$\text{or, } Re^{i\omega t + \theta} = e^{i(\omega t + \phi_1)} (A_1 + A_2 e^{i\delta})$$

$$\text{where } R = \sqrt{(Re(z))^2 + (Im(z))^2},$$

$$\tan \theta = \frac{Im(z)}{Re(z)}$$

Ex.: Show that you get the same formulae as from the geometric picture.

Multiple vibrations of same frequency

and same relative phase

$$x_1 = a \cos(\omega t),$$



Multiple vibrations of same frequency

and same relative phase

$$x_1 = a \cos(\omega t),$$

$$x_2 = a \cos(\omega t + \delta),$$



Multiple vibrations of same frequency

and same relative phase

$$x_1 = a \cos(\omega t),$$

$$x_2 = a \cos(\omega t + \delta),$$

$$x_3 = a \cos(\omega t + 2\delta),$$



Multiple vibrations of same frequency

and same relative phase


$$x_1 = a \cos(\omega t),$$

$$x_2 = a \cos(\omega t + \delta),$$

$$x_3 = a \cos(\omega t + 2\delta),$$


$$\vdots$$

$$x_n = a \cos(\omega t + (n - 1)\delta).$$



Multiple vibrations of same frequency

and same relative phase



$$x_1 = a \cos(\omega t),$$

$$x_2 = a \cos(\omega t + \delta),$$


$$x_3 = a \cos(\omega t + 2\delta),$$

$$\vdots$$

$$x_n = a \cos(\omega t + (n - 1)\delta).$$

Multiple vibrations of same frequency

and same relative phase



$$x_1 = a \cos(\omega t),$$

$$x_2 = a \cos(\omega t + \delta),$$


$$x_3 = a \cos(\omega t + 2\delta),$$

$$\vdots$$

$$x_n = a \cos(\omega t + (n - 1)\delta).$$

Multiple vibrations of same frequency

and same relative phase



$$x_1 = a \cos(\omega t),$$

$$x_2 = a \cos(\omega t + \delta),$$

$$x_3 = a \cos(\omega t + 2\delta),$$


$$\vdots$$

$$x_n = a \cos(\omega t + (n - 1)\delta).$$

$$X = A \cos(\omega t + \alpha)$$

Multiple vibrations of same frequency

and same relative phase



$$x_1 = a \cos(\omega t),$$

$$x_2 = a \cos(\omega t + \delta),$$

$$x_3 = a \cos(\omega t + 2\delta),$$

$$\vdots$$

$$x_n = a \cos(\omega t + (n-1)\delta).$$

$$X = A \cos(\omega t + \alpha)$$


What are A and α ?

Multiple vibrations of same frequency

and same relative phase

Geometric method:


Construct a circle superscribing the polygon, of radius R centered at O .



$$X = A \cos(\omega t + \alpha).$$

Multiple vibrations of same frequency

and same relative phase



$$X = A \cos(\omega t + \alpha).$$


Geometric method:

Construct a circle superscribing the polygon, of radius R centered at O .

Observe that $\sin\left(\frac{n\delta}{2}\right) = \frac{A}{2R}, \quad \sin\left(\frac{\delta}{2}\right) = \frac{a}{2R}.$

Multiple vibrations of same frequency

and same relative phase



$$X = A \cos(\omega t + \alpha).$$

Geometric method:


Construct a circle superscribing the polygon, of radius R centered at O .

Observe that $\sin\left(\frac{n\delta}{2}\right) = \frac{A}{2R}$, $\sin\left(\frac{\delta}{2}\right) = \frac{a}{2R}$.

$$A = 2R \sin\left(\frac{n\delta}{2}\right),$$

Multiple vibrations of same frequency

and same relative phase



$$X = A \cos(\omega t + \alpha).$$

Geometric method:


Construct a circle superscribing the polygon, of radius R centered at O .

Observe that $\sin\left(\frac{n\delta}{2}\right) = \frac{A}{2R}$, $\sin\left(\frac{\delta}{2}\right) = \frac{a}{2R}$.

$$A = 2R \sin\left(\frac{n\delta}{2}\right), \quad a = 2R \sin\left(\frac{\delta}{2}\right),$$

Multiple vibrations of same frequency

and same relative phase



$$X = A \cos(\omega t + \alpha).$$

Geometric method:


Construct a circle superscribing the polygon, of radius R centered at O .

Observe that $\sin\left(\frac{n\delta}{2}\right) = \frac{A}{2R}$, $\sin\left(\frac{\delta}{2}\right) = \frac{a}{2R}$.

$$A = 2R \sin\left(\frac{n\delta}{2}\right), \quad a = 2R \sin\left(\frac{\delta}{2}\right),$$

$$\alpha = \angle OPS - \angle OPQ.$$

and same relative phase



Construct a circle superscribing the polygon, of radius R centered at O .

$$\alpha = \angle OPS - \angle OPQ.$$

$$A = a \frac{\sin(n\delta/2)}{\sin(\delta/2)}, \quad \alpha = (n-1) \frac{\delta}{2}.$$

Multiple vibrations of same frequency

and same relative phase

Using Complex Exponentials: $Z(t) = X(t) + iY(t) = Ae^{i(\omega t + \alpha)}$

Multiple vibrations of same frequency

and same relative phase

Using Complex Exponentials: $Z(t) = X(t) + iY(t) = Ae^{i(\omega t + \alpha)}$

$$Z = ae^{i\omega t} + ae^{i\omega t + \delta} + ae^{i\omega t + 2\delta} \dots + ae^{i\omega t + (n-1)\delta}.$$

Multiple vibrations of same frequency

and same relative phase

Using Complex Exponentials: $Z(t) = X(t) + iY(t) = Ae^{i(\omega t + \alpha)}$

$$\begin{aligned} Z &= ae^{i\omega t} + ae^{i\omega t + \delta} + ae^{i\omega t + 2\delta} \dots + ae^{i\omega t + (n-1)\delta} \\ &= ae^{i\omega t}(1 + f + f^2 + \dots + f^{n-1}) \quad \text{where } f = e^{i\delta}. \end{aligned}$$

Multiple vibrations of same frequency

and same relative phase

Using Complex Exponentials: $Z(t) = X(t) + iY(t) = Ae^{i(\omega t + \alpha)}$

$$\begin{aligned} Z &= ae^{i\omega t} + ae^{i\omega t + \delta} + ae^{i\omega t + 2\delta} \dots + ae^{i\omega t + (n-1)\delta} \\ &= ae^{i\omega t}(1 + f + f^2 + \dots + f^{n-1}) \quad \text{where } f = e^{i\delta}. \end{aligned}$$

Sum of geometric progression:

Multiple vibrations of same frequency

and same relative phase

Using Complex Exponentials: $Z(t) = X(t) + iY(t) = Ae^{i(\omega t + \alpha)}$

$$\begin{aligned} Z &= ae^{i\omega t} + ae^{i\omega t + \delta} + ae^{i\omega t + 2\delta} \dots + ae^{i\omega t + (n-1)\delta} \\ &= ae^{i\omega t}(1 + f + f^2 + \dots + f^{n-1}) \quad \text{where } f = e^{i\delta}. \end{aligned}$$

Sum of geometric progression:

$$S(f) = 1 + f + f^2 + \dots + f^{n-1}$$

Multiple vibrations of same frequency

and same relative phase

Using Complex Exponentials: $Z(t) = X(t) + iY(t) = Ae^{i(\omega t + \alpha)}$

$$\begin{aligned} Z &= ae^{i\omega t} + ae^{i\omega t + \delta} + ae^{i\omega t + 2\delta} \dots + ae^{i\omega t + (n-1)\delta} \\ &= ae^{i\omega t}(1 + f + f^2 + \dots + f^{n-1}) \quad \text{where } f = e^{i\delta}. \end{aligned}$$

Sum of geometric progression:

$$\begin{aligned} S(f) &= 1 + f + f^2 + \dots + f^{n-1} \\ fS(f) &= f + f^2 + \dots + f^n. \end{aligned}$$

Multiple vibrations of same frequency

and same relative phase

Using Complex Exponentials: $Z(t) = X(t) + iY(t) = Ae^{i(\omega t + \alpha)}$

$$\begin{aligned} Z &= ae^{i\omega t} + ae^{i\omega t + \delta} + ae^{i\omega t + 2\delta} \dots + ae^{i\omega t + (n-1)\delta} \\ &= ae^{i\omega t}(1 + f + f^2 + \dots + f^{n-1}) \quad \text{where } f = e^{i\delta}. \end{aligned}$$

Sum of geometric progression:

$$S(f) = 1 + f + f^2 + \dots + f^{n-1}$$

$$fS(f) = f + f^2 + \dots + f^n.$$

Subtracting, $S(f) = \frac{1 - f^n}{1 - f}$

Multiple vibrations of same frequency

and same relative phase

Using Complex Exponentials: $Z(t) = X(t) + iY(t) = Ae^{i(\omega t + \alpha)}$

$$\begin{aligned} Z &= ae^{i\omega t} + ae^{i\omega t + \delta} + ae^{i\omega t + 2\delta} \dots + ae^{i\omega t + (n-1)\delta} \\ &= ae^{i\omega t}(1 + f + f^2 + \dots + f^{n-1}) \quad \text{where } f = e^{i\delta}. \end{aligned}$$

Sum of geometric progression:

$$S(f) = 1 + f + f^2 + \dots + f^{n-1}$$

$$fS(f) = f + f^2 + \dots + f^n.$$

$$\text{Subtracting, } S(f) = \frac{1 - f^n}{1 - f} = \frac{1 - e^{in\delta}}{1 - e^{i\delta}}$$

Multiple vibrations of same frequency

and same relative phase

Using Complex Exponentials: $Z(t) = X(t) + iY(t) = Ae^{i(\omega t + \alpha)}$

$$Z = ae^{i\omega t} + ae^{i\omega t + \delta} + ae^{i\omega t + 2\delta} \dots + ae^{i\omega t + (n-1)\delta}.$$
$$= ae^{i\omega t}(1 + f + f^2 + \dots + f^{n-1}) \quad \text{where } f = e^{i\delta}.$$

Sum of geometric progression:

$$S(f) = 1 + f + f^2 + \dots + f^{n-1}$$

$$fS(f) = f + f^2 + \dots + f^n.$$

$$\text{Subtracting, } S(f) = \frac{1 - f^n}{1 - f} = \frac{1 - e^{in\delta}}{1 - e^{i\delta}} = \frac{e^{in\delta/2}(e^{-in\delta/2} - e^{in\delta/2})}{e^{i\delta/2}(e^{-i\delta/2} - e^{i\delta/2})}$$

Multiple vibrations of same frequency

and same relative phase

Using Complex Exponentials: $Z(t) = X(t) + iY(t) = Ae^{i(\omega t + \alpha)}$

$$Z = ae^{i\omega t} + ae^{i\omega t + \delta} + ae^{i\omega t + 2\delta} \dots + ae^{i\omega t + (n-1)\delta}.$$
$$= ae^{i\omega t}(1 + f + f^2 + \dots + f^{n-1}) \quad \text{where } f = e^{i\delta}.$$

Sum of geometric progression:

$$S(f) = 1 + f + f^2 + \dots + f^{n-1}$$

$$fS(f) = f + f^2 + \dots + f^n.$$

$$\text{Subtracting, } S(f) = \frac{1 - f^n}{1 - f} = \frac{1 - e^{in\delta}}{1 - e^{i\delta}} = \frac{e^{in\delta/2}(e^{-in\delta/2} - e^{in\delta/2})}{e^{i\delta/2}(e^{-i\delta/2} - e^{i\delta/2})} = e^{i(n-1)\delta/2} \frac{\sin(n\delta/2)}{\sin(\delta/2)}.$$

Multiple vibrations of same frequency

and same relative phase

When n is very large and δ very small, $\alpha \sim n\delta/2$,

$$X = na \frac{\sin \alpha}{\alpha} \cos(\omega t + n\delta/2)$$

Amplitude depends

on δ as a

“sinc” function.

Multiple vibrations of same frequency

and same relative phase

When n is very large and δ very small, $\alpha \sim n\delta/2$,


$$X = na \frac{\sin \alpha}{\alpha} \cos(\omega t + n\delta/2)$$

Amplitude depends


on δ as a

“sinc” function.

This is the pattern for diffraction from a thin slit!




Addition of Different Frequencies




$$\omega_1 = 2$$

$$x_1 = A \cos(\omega_1 t), \quad x_2 = A \cos(\omega_2 t);$$



$$\omega_2 = 3.5$$

Addition of Different Frequencies




$$\omega_1 = 2$$


$$x_1 = A \cos(\omega_1 t), \quad x_2 = A \cos(\omega_2 t);$$

Resultant wave is a complicated function of time:


$$x = 2A \cos\left(\frac{\omega_1 - \omega_2}{2}t\right) \cos\left(\frac{\omega_1 + \omega_2}{2}t\right).$$



$$\omega_2 = 3.5$$



Addition of Different Frequencies




$$\omega_1 = 2$$

$$x_1 = A \cos(\omega_1 t), \quad x_2 = A \cos(\omega_2 t);$$

Resultant wave is a complicated function of time:

$$x = 2A \cos\left(\frac{\omega_1 - \omega_2}{2}t\right) \cos\left(\frac{\omega_1 + \omega_2}{2}t\right).$$




$$\omega_2 = 6$$

Periodic with period T **only** if T_1 and T_2 are commensurate:


$$\text{if } n_1 T_1 = n_2 T_2 = T$$

for integer n_1 and n_2 .



$$x_1 + x_2$$

Addition of Different Frequencies




$$\omega_1 = 2$$

$$x_1 = A \cos(\omega_1 t), \quad x_2 = A \cos(\omega_2 t);$$

Resultant wave is a complicated function of time:

$$x = 2A \cos\left(\frac{\omega_1 - \omega_2}{2}t\right) \cos\left(\frac{\omega_1 + \omega_2}{2}t\right).$$




$$\omega_2 = 6$$

Periodic with period T **only** if T_1 and T_2 are commensurate:

$$\text{if } n_1 T_1 = n_2 T_2 = T$$


for integer n_1 and n_2 .




$$x_1 + x_2$$

Here, $T = T_1 = 3T_2$

When the frequencies are nearly equal...Beats!




$$\omega_1 = 5$$




$$\omega_2 = 5.5$$

$$x = x_1 + x_2 = A(\cos \omega_1 t + \cos \omega_2 t) = 2A \cos \left(\frac{\omega_1 - \omega_2}{2} t \right) \cos \left(\frac{\omega_1 + \omega_2}{2} t \right)$$


When the frequencies are nearly equal...Beats!



$$\omega_1 = 5$$



$$\omega_2 = 5.5$$




$$x_1 + x_2$$


$$x = x_1 + x_2 = A(\cos \omega_1 t + \cos \omega_2 t) = 2A \cos \left(\frac{\omega_1 - \omega_2}{2} t \right) \cos \left(\frac{\omega_1 + \omega_2}{2} t \right)$$

If $\omega_1 \sim \omega_2$, this is an oscillation of frequency $\omega_1 \sim \omega_2 \sim \frac{\omega_1 + \omega_2}{2}$ (average),


When the frequencies are nearly equal...Beats!



$$\omega_1 = 5$$



$$\omega_2 = 5.5$$




$$x_1 + x_2, \omega_b = 0.25$$

$$x = x_1 + x_2 = A(\cos \omega_1 t + \cos \omega_2 t) = 2A \cos \left(\frac{\omega_1 - \omega_2}{2} t \right) \cos \left(\frac{\omega_1 + \omega_2}{2} t \right)$$


If $\omega_1 \sim \omega_2$, this is an oscillation of frequency $\omega_1 \sim \omega_2 \sim \frac{\omega_1 + \omega_2}{2}$ (average),

with an envelop of frequency $\frac{|\omega_1 - \omega_2|}{2}$.


When the frequencies are nearly equal...Beats!



$$\omega_1 = 5$$



$$\omega_2 = 5.5$$



$$x_1 + x_2, \omega_b = 0.25$$

$$x = x_1 + x_2 = A(\cos \omega_1 t + \cos \omega_2 t) = 2A \cos \left(\frac{\omega_1 - \omega_2}{2} t \right) \cos \left(\frac{\omega_1 + \omega_2}{2} t \right)$$

If $\omega_1 \sim \omega_2$, this is an oscillation of frequency $\omega_1 \sim \omega_2 \sim \frac{\omega_1 + \omega_2}{2}$ (average),

with an envelop of frequency $\frac{|\omega_1 - \omega_2|}{2}$.

Amplitude oscillates with frequency $\Delta\omega = |\omega_1 - \omega_2|$: beat frequency.

Perpendicular Vibrations: Lissajous Figures

1. In phase:

$$x = A_1 \sin(\omega t),$$


$$y = A_2 \sin(\omega t),$$

Perpendicular Vibrations: Lissajous Figures

1. In phase:

$$x = A_1 \sin(\omega t),$$

$$y = A_2 \sin(\omega t),$$




Perpendicular Vibrations: Lissajous Figures

1. In phase:

$$x = A_1 \sin(\omega t),$$

$$y = A_2 \sin(\omega t),$$




Perpendicular Vibrations: Lissajous Figures

1. In phase:

$$x = A_1 \sin(\omega t),$$

$$y = A_2 \sin(\omega t),$$




Perpendicular Vibrations: Lissajous Figures

1. In phase:

$$x = A_1 \sin(\omega t),$$

$$y = A_2 \sin(\omega t),$$




Perpendicular Vibrations: Lissajous Figures

1. In phase:

$$x = A_1 \sin(\omega t),$$

$$y = A_2 \sin(\omega t),$$



Perpendicular Vibrations: Lissajous Figures


1. In phase:

$$x = A_1 \sin(\omega t),$$

$$y = A_2 \sin(\omega t),$$

Resultant motion is a straight line with slope

$$\frac{y}{x} = \frac{A_2}{A_1}.$$



Perpendicular Vibrations: Lissajous Figures

1. In phase:

$$x = A_1 \sin(\omega t),$$

$$y = A_2 \sin(\omega t),$$

Resultant motion is a
straight line with slope

$$\frac{y}{x} = \frac{A_2}{A_1}.$$

Perpendicular Vibrations: same frequency


2. Out of phase by $\pi/2$:

$$x = A_1 \cos(\omega t)$$

$$y = A_2 \sin(\omega t)$$

Resultant motion traces
the ellipse

$$\frac{x^2}{A_1^2} + \frac{y^2}{A_2^2} = 1.$$



Perpendicular Vibrations: same frequency

2. Out of phase by $\pi/2$:

$$x = A_1 \cos(\omega t)$$

$$y = A_2 \sin(\omega t)$$

Resultant motion traces
the ellipse

$$\frac{x^2}{A_1^2} + \frac{y^2}{A_2^2} = 1.$$

Perpendicular Vibrations: same frequency

In general,

$$x = A_1 \cos(\omega t), \quad y = A_2 \cos(\omega t + \delta).$$

Perpendicular Vibrations: same frequency

In general,

$$x = A_1 \cos(\omega t), \quad y = A_2 \cos(\omega t + \delta).$$

Let $\frac{x}{A_1} = \tilde{x}$ and $\frac{y}{A_2} = \tilde{y}$,

we get $\tilde{x}^2 + \tilde{y}^2 - 2\tilde{x}\tilde{y} \cos \delta = \sin^2 \delta$,

Perpendicular Vibrations: same frequency

In general,

$$x = A_1 \cos(\omega t), \quad y = A_2 \cos(\omega t + \delta).$$

Let $\frac{x}{A_1} = \tilde{x}$ and $\frac{y}{A_2} = \tilde{y}$,

we get $\tilde{x}^2 + \tilde{y}^2 - 2\tilde{x}\tilde{y}\cos\delta = \sin^2\delta$, —an ellipse

Perpendicular Vibrations: same frequency

In general,

$$x = A_1 \cos(\omega t), \quad y = A_2 \cos(\omega t + \delta).$$

Let $\frac{x}{A_1} = \tilde{x}$ and $\frac{y}{A_2} = \tilde{y}$,

we get $\tilde{x}^2 + \tilde{y}^2 - 2\tilde{x}\tilde{y}\cos\delta = \sin^2\delta$, —an ellipse which degenerates into a straight line for $\delta = 0, \pi$.

Perpendicular Vibrations: same frequency

In general,

$$x = A_1 \cos(\omega t), \quad y = A_2 \cos(\omega t + \delta).$$


Let $\frac{x}{A_1} = \tilde{x}$ and $\frac{y}{A_2} = \tilde{y}$,

we get $\tilde{x}^2 + \tilde{y}^2 - 2\tilde{x}\tilde{y}\cos\delta = \sin^2\delta$, —an ellipse which degenerates into a straight line for $\delta = 0, \pi$.

If $A_1 = A_2$, you get a circle for $\delta = \pi/2$.

Lissajous figures: same frequency


Changing δ from 0 through 2π .




$$\delta = 0$$




$$\delta = \pi/4$$




$$\delta = \pi/2$$




$$\delta = 3\pi/4$$




$$\delta = \pi$$




$$\delta = 4\pi/4$$



$$\delta = 3\pi/2$$



$$\delta = 7\pi/4$$



$$\delta = 2\pi$$

Perpendicular vibrations with different frequencies

$$\omega_1 = 2, \omega_2 = 3$$

Perpendicular vibrations with different frequencies

$$\omega_1 = 2, \omega_2 = 3$$

Perpendicular vibrations with different frequencies

For different
frequency ratios

