

Functions of Several variables

Saranya G. Nair
Department of Mathematics

BITS Pilani

September 25, 2024



Definition: Function of Several Variables

The cartesian product \mathbb{R}^n is $\{(r_1, r_2, \dots, r_n) : r_1, r_2, \dots, r_n \in \mathbb{R}\}$.

Definition

Let D be a non-empty subset of \mathbb{R}^n . Then a map $f : D \rightarrow \mathbb{R}$ is called a **function of several variables**.

Definition: Function of Several Variables

The cartesian product \mathbb{R}^n is $\{(r_1, r_2, \dots, r_n) : r_1, r_2, \dots, r_n \in \mathbb{R}\}$.

Definition

Let D be a non-empty subset of \mathbb{R}^n . Then a map $f : D \rightarrow \mathbb{R}$ is called a **function of several variables**.

Remark

In this case, we often denote $w = f(x_1, x_2, \dots, x_n)$.

*Observe that w is the **dependent variable** of f and f is called a function of the n **independent variables** x_1 to x_n .*

*We often call x_1, x_2, \dots, x_n as function's **input variables** and w as function's **output variable**.*

Example: Function of Several Variables

Let $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ be defined by

$$f(x_1, x_2, x_3) = \sqrt{x_1^2 + x_2^2 + x_3^2}.$$

Then $f(0, 3, 4) = 5$ and $f(1, 1, 2) = \sqrt{6}$.

Finding Domain and Ranges Function of Several Variables

Example

① $z = \sqrt{y - x^2}$

② $z = \frac{1}{xy}$

③ $z = \sin xy$

④ $w = \sqrt{x^2 + y^2 + z^2}$

⑤ $w = \frac{1}{x^2 + y^2 + z^2}$

⑥ $w = xy \ln z$

Definitions

Definition (Open disk)

Let $r > 0$ be a real number and (x_0, y_0) be a point of \mathbb{R}^2 . Then the set

$$\{(x, y) \in \mathbb{R}^2 : \sqrt{(x - x_0)^2 + (y - y_0)^2} < r\}$$

is called the **open disk** or **open ball** of radius r at (x_0, y_0) and is denoted by $B_r(x_0, y_0)$.

Open and Closed

Interior Point: A point (x_0, y_0) of a region R in the xy plane is called an interior point of R if it is the center of a disk of positive radius that lies entirely in R .

Open and Closed


Interior Point: A point (x_0, y_0) of a region R in the xy plane is called an interior point of R if it is the center of a disk of positive radius that lies entirely in R .

Boundary Point: A point (x_0, y_0) is a boundary point of R if every disk centered at (x_0, y_0) contains points that lie outside of R as well as points that lie in R . (The boundary point itself need not belong to R .)

Open and Closed

Interior Point: A point (x_0, y_0) of a region R in the xy plane is called an interior point of R if it is the center of a disk of positive radius that lies entirely in R .

Boundary Point: A point (x_0, y_0) is a boundary point of R if every disk centered at (x_0, y_0) contains points that lie outside of R as well as points that lie in R . (The boundary point itself need not belong to R .)



The set of all interior points of a region is called **interior** of the region.
The set of all boundary points of a region is called **boundary** of the region.

A region is **open** if every point in that region is an interior point.

A region is **open** if every point in that region is an interior point. A region is **closed** if it contains all its boundary points.

A region is **open** if every point in that region is an interior point. A region is **closed** if it contains all its boundary points.

Describe each of the following sets, open or closed.

① $\{(x, y) : x^2 + y^2 < 1\}$:

A region is **open** if every point in that region is an interior point. A region is **closed** if it contains all its boundary points.

Describe each of the following sets, open or closed.

① $\{(x, y) : x^2 + y^2 < 1\}$: Open

② $\{(x, y) : x^2 + y^2 = 1\}$:

A region is **open** if every point in that region is an interior point. A region is **closed** if it contains all its boundary points.

Describe each of the following sets, open or closed.

- ① $\{(x, y) : x^2 + y^2 < 1\}$: Open
- ② $\{(x, y) : x^2 + y^2 = 1\}$: Closed
- ③ $\{(x, y) : x^2 + y^2 \leq 1\}$:

A region is **open** if every point in that region is an interior point. A region is **closed** if it contains all its boundary points.

Describe each of the following sets, open or closed.

- ① $\{(x, y) : x^2 + y^2 < 1\}$: Open
- ② $\{(x, y) : x^2 + y^2 = 1\}$: Closed
- ③ $\{(x, y) : x^2 + y^2 \leq 1\}$: Closed
- ④ $\{(x, y) : x^2 + y^2 > 1\}$:

A region is **open** if every point in that region is an interior point. A region is **closed** if it contains all its boundary points.

Describe each of the following sets, open or closed.

- ① $\{(x, y) : x^2 + y^2 < 1\}$: Open
- ② $\{(x, y) : x^2 + y^2 = 1\}$: Closed
- ③ $\{(x, y) : x^2 + y^2 \leq 1\}$: Closed
- ④ $\{(x, y) : x^2 + y^2 > 1\}$: Open
- ⑤ Empty set:

A region is **open** if every point in that region is an interior point. A region is **closed** if it contains all its boundary points.

Describe each of the following sets, open or closed.

- ① $\{(x, y) : x^2 + y^2 < 1\}$: Open
- ② $\{(x, y) : x^2 + y^2 = 1\}$: Closed
- ③ $\{(x, y) : x^2 + y^2 \leq 1\}$: Closed
- ④ $\{(x, y) : x^2 + y^2 > 1\}$: Open
- ⑤ Empty set: Both Open and Closed
- ⑥ Entire plane, \mathbb{R}^2 :

A region is **open** if every point in that region is an interior point. A region is **closed** if it contains all its boundary points.

Describe each of the following sets, open or closed.

- ① $\{(x, y) : x^2 + y^2 < 1\}$: Open
- ② $\{(x, y) : x^2 + y^2 = 1\}$: Closed
- ③ $\{(x, y) : x^2 + y^2 \leq 1\}$: Closed
- ④ $\{(x, y) : x^2 + y^2 > 1\}$: Open
- ⑤ Empty set: Both Open and Closed
- ⑥ Entire plane, \mathbb{R}^2 : Both Open and Closed

A region is **open** if every point in that region is an interior point. A region is **closed** if it contains all its boundary points.

Describe each of the following sets, open or closed.

- ① $\{(x, y) : x^2 + y^2 < 1\}$: Open
- ② $\{(x, y) : x^2 + y^2 = 1\}$: Closed
- ③ $\{(x, y) : x^2 + y^2 \leq 1\}$: Closed
- ④ $\{(x, y) : x^2 + y^2 > 1\}$: Open
- ⑤ Empty set: Both Open and Closed
- ⑥ Entire plane, \mathbb{R}^2 : Both Open and Closed

Definition

A region in the plane is bounded if it lies inside a disk of finite radius.

A region is **open** if every point in that region is an interior point. A region is **closed** if it contains all its boundary points.

Describe each of the following sets, open or closed.

- ① $\{(x, y) : x^2 + y^2 < 1\}$: Open
- ② $\{(x, y) : x^2 + y^2 = 1\}$: Closed
- ③ $\{(x, y) : x^2 + y^2 \leq 1\}$: Closed
- ④ $\{(x, y) : x^2 + y^2 > 1\}$: Open
- ⑤ Empty set: Both Open and Closed
- ⑥ Entire plane, \mathbb{R}^2 : Both Open and Closed

Definition

A region in the plane is bounded if it lies inside a disk of finite radius. A region is unbounded if it is not bounded.

Find and sketch the domain for each function.

① $f(x, y) = \sqrt{y - x^2}$

② $f(x, y) = \sqrt{y - x - 2}$

③ $f(x, y) = \frac{1}{\ln(4 - x^2 - y^2)}$

④ $f(x, y) = \sqrt{(x^2 - 4)(y^2 - 9)}$

For the following functions (a) find the function's domain, (b) find the function's range, (c) describe the function's level curves, (d) find the boundary of the function's domain, (e) determine if the domain is an open region, a closed region, or neither, and (f) decide if the domain is bounded or unbounded.

① $f(x, y) = y - x$

② $f(x, y) = \sqrt{y - x}$

③ $f(x, y) = \frac{1}{\sqrt{16 - x^2 - y^2}}$

④ $f(x, y) = e^{-(x^2 + y^2)}$

Level Curves

The set of all points $(x, y, f(x, y))$ in space, for (x, y) in the domain of f , is called the graph of f . The graph of f is also called the surface $z = f(x, y)$.

Level Curves

The set of all points $(x, y, f(x, y))$ in space, for (x, y) in the domain of f , is called the graph of f . The graph of f is also called the surface $z = f(x, y)$.

Definition


The set of points in the plane where a function $f(x, y)$ has a constant value $f(x, y) = c$ is called a level curve of f .

Level Curves

The set of all points $(x, y, f(x, y))$ in space, for (x, y) in the domain of f , is called the graph of f . The graph of f is also called the surface $z = f(x, y)$.



Definition

The set of points in the plane where a function $f(x, y)$ has a constant value $f(x, y) = c$ is called a level curve of f .





Consider $f(x, y) = x^2 + y^2$. The graph $(x, y, f(x, y))$ and level curves $f(x, y) = k$ for $k = 0, 1, 2, 3, 4, 5$ are

Consider $f(x, y) = x^2 + y^2$. The graph $(x, y, f(x, y))$ and level curves $f(x, y) = k$ for $k = 0, 1, 2, 3, 4, 5$ are




Level Curves



Sketch some level curves of the function $f(x, y) = 4x^2 + y^2$.

Sketch some level curves of the function $f(x, y) = 4x^2 + y^2$.





Example


Graph $f(x, y) = 1 - x^2 - y^2$ and plot the level curves $f(x, y) = 0$, $f(x, y) = 1$, $f(x, y) = 2$ and $f(x, y) = -2$ in the domain of f in the plane.

Example


Graph $f(x, y) = 1 - x^2 - y^2$ and plot the level curves $f(x, y) = 0$, $f(x, y) = 1$, $f(x, y) = 2$ and $f(x, y) = -2$ in the domain of f in the plane.




Some examples






(a) $z = \sin x + 2 \sin y$



(b) $z = (4x^2 + y^2)e^{-x^2-y^2}$



(c) $z = xye^{-y^2}$



Definition (Level Surface)

Let f be a real valued function with domain $B \subseteq \mathbb{R}^3$; then for a real constant c , the set of all points (x, y, z) of B satisfying $f(x, y, z) = c$ is called the **level surface**.

Example


Describe the level surfaces of the function

$$f(x, y, z) = \sqrt{x^2 + y^2 + z^2}.$$

Example

Describe the level surfaces of the function

$$f(x, y, z) = \sqrt{x^2 + y^2 + z^2}.$$



Find and sketch the level curves $f(x, y) = c$ on the same set of coordinate axes for the given values of c .

① $f(x, y) = x + y - 1, \quad c = -3, -2, -1, 0, 1, 2, 3$

② $f(x, y) = x^2 + y^2, \quad c = 0, 1, 4, 9, 16, 25$

③ $f(x, y) = xy, \quad c = -9, -4, -1, 0, 1, 4, 9$

Open and Closed sets in \mathbb{R}^3

The definitions of interior, boundary, open, closed and unbounded for regions in space are similar to those of regions in plane. To accomodate the extra dimension, we use solid balls of positive radius instead of disks. An open ball B in \mathbb{R}^3 centered at (x_0, y_0, z_0) with radius r is the set of all points (x, y, z) such that

$$\sqrt{(x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2} < r.$$