

MATH F111- Mathematics I

Saranya G. Nair
Department of Mathematics

BITS Pilani

August 12, 2024



Properties of limits

Theorem

Uniqueness of Limits. *A sequence in \mathbb{R} can have at most one limit.*

Properties of limits

Theorem

Uniqueness of Limits. *A sequence in \mathbb{R} can have at most one limit.*

Proof. Let (a_n) be a real sequence and suppose that ℓ_1 and ℓ_2 are both limits for (a_n) and let $\ell_1 \neq \ell_2$.

- Let $\epsilon := |\ell_1 - \ell_2|/2$. Since $\ell_1 \neq \ell_2, \epsilon > 0$.
- Since ℓ_1 is a limit of the sequence, for the chosen $\epsilon, \exists N_1 \in \mathbb{N}$ such that

$$|a_n - \ell_1| < \epsilon, \text{ for all } n \geq N_1.$$

- Since ℓ_2 is a limit of the sequence, for the chosen $\epsilon, \exists N_2 \in \mathbb{N}$ such that

$$|a_n - \ell_2| < \epsilon, \text{ for all } n \geq N_2.$$

Let $N = \max\{N_1, N_2\}$. Then:

$$|a_n - \ell_1| < \epsilon \quad \text{and} \quad |a_n - \ell_2| < \epsilon \quad \text{for all } n \geq N,$$

and hence,

$$|\ell_1 - \ell_2| = |(a_N - \ell_2) - (a_N - \ell_1)| \leq |a_N - \ell_1| + |a_N - \ell_2| < \epsilon + \epsilon = |\ell_1 - \ell_2|$$

which is a contradiction. Hence, $\ell_1 = \ell_2$. ■

Theorem

A convergent sequence is bounded.

Theorem

A convergent sequence is bounded.

Suppose $a_n \rightarrow \ell$. Let $\epsilon := 1$. There is $N \in \mathbb{N}$ such that

$$|a_n - \ell| < 1 \text{ for all } n \geq N.$$

Hence

$$|a_n| \leq |a_n - \ell| + |\ell| < 1 + |\ell| \text{ for all } n \geq N.$$

Theorem

A convergent sequence is bounded.

Suppose $a_n \rightarrow \ell$. Let $\epsilon := 1$. There is $N \in \mathbb{N}$ such that

$$|a_n - \ell| < 1 \text{ for all } n \geq N.$$

Hence

$$|a_n| \leq |a_n - \ell| + |\ell| < 1 + |\ell| \text{ for all } n \geq N.$$

- Thus it remains to find a bound for a_1, a_2, \dots, a_{N-1} . Choose $\beta = \max\{|a_1|, |a_2|, \dots, |a_{N-1}|\}$. Then $|a_n| \leq \beta$, for all $1 \leq n \leq N-1$.
- Define $\alpha := \max\{|a_1|, \dots, |a_{N-1}|, |\ell| + 1\}$. Then $|a_n| \leq \alpha$ for all $n \in \mathbb{N}$. Hence (a_n) is bounded.

If a_n is convergent, then a_n is bounded. Equivalently, if a_n is not bounded, then a_n is not convergent. This result can be used to show if a sequence is not bounded.

- The sequence $\{(-1)^n n : n \in \mathbb{N}\}$ divergent since it is not bounded.
- A bounded sequence need not be convergent. For example, the sequence $\{(-1)^n : n \in \mathbb{N}\}$ is bounded but not convergent.

Theorem

Limit theorems. *Let $\{a_n\}$ and $\{b_n\}$ be two convergent sequences that converge to A and B respectively. Then:*

- $\lim(a_n \pm b_n) = A \pm B.$

- $\lim(a_n b_n) = AB.$

In particular, $\lim(ca_n) = cA$ for $c \in \mathbb{R}.$

- $\lim \frac{a_n}{b_n} = \frac{A}{B},$ provided (b_n) is a sequence of non-zero real numbers and $B \neq 0.$

Theorem

Limit theorems. Let $\{a_n\}$ and $\{b_n\}$ be two convergent sequences that converge to A and B respectively. Then:

- $\lim(a_n \pm b_n) = A \pm B.$

- $\lim(a_n b_n) = AB.$

In particular, $\lim(ca_n) = cA$ for $c \in \mathbb{R}.$

- $\lim \frac{a_n}{b_n} = \frac{A}{B},$ provided (b_n) is a sequence of non-zero real numbers and $B \neq 0.$

- $\lim_{n \rightarrow \infty} \frac{-1}{n} = -\lim_{n \rightarrow \infty} \frac{1}{n} = 0.$

Theorem

Limit theorems. Let $\{a_n\}$ and $\{b_n\}$ be two convergent sequences that converge to A and B respectively. Then:

- $\lim(a_n \pm b_n) = A \pm B$.
 - $\lim(a_n b_n) = AB$.
In particular, $\lim(ca_n) = cA$ for $c \in \mathbb{R}$.
 - $\lim \frac{a_n}{b_n} = \frac{A}{B}$, provided (b_n) is a sequence of non-zero real numbers and $B \neq 0$.
-
- $\lim_{n \rightarrow \infty} \frac{-1}{n} = -\lim_{n \rightarrow \infty} \frac{1}{n} = 0$.
 - $\lim_{n \rightarrow \infty} \frac{n+1}{n} = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right) = 1$.

Theorem

Limit theorems. Let $\{a_n\}$ and $\{b_n\}$ be two convergent sequences that converge to A and B respectively. Then:

- $\lim(a_n \pm b_n) = A \pm B.$

- $\lim(a_n b_n) = AB.$

In particular, $\lim(ca_n) = cA$ for $c \in \mathbb{R}.$

- $\lim \frac{a_n}{b_n} = \frac{A}{B},$ provided (b_n) is a sequence of non-zero real numbers and $B \neq 0.$

- $\lim_{n \rightarrow \infty} \frac{-1}{n} = -\lim_{n \rightarrow \infty} \frac{1}{n} = 0.$

- $\lim_{n \rightarrow \infty} \frac{n+1}{n} = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right) = 1.$

- $\lim_{n \rightarrow \infty} \frac{4-7n^6}{n^6+3} = \lim_{n \rightarrow \infty} \frac{\left(\frac{4}{n^6}-7\right)}{1+\frac{3}{n^6}} = \frac{0-7}{1+0} = -7.$

Theorem

Let (x_n) be a convergent sequence of real numbers and there exists a positive integer m such that $x_n \geq 0$ for all $n \geq m$. Then $\lim x_n \geq 0$.

Theorem

Let (x_n) be a convergent sequence of real numbers and there exists a positive integer m such that $x_n \geq 0$ for all $n \geq m$. Then $\lim x_n \geq 0$.

Corollary

If (x_n) and (y_n) are convergent sequences of real numbers and if there is a positive integer m such that $x_n \leq y_n$ for all $n \geq m$, then $\lim x_n \leq \lim y_n$.

Theorem

Let (x_n) be a convergent sequence of real numbers and there exists a positive integer m such that $x_n \geq 0$ for all $n \geq m$. Then $\lim x_n \geq 0$.

Corollary

If (x_n) and (y_n) are convergent sequences of real numbers and if there is a positive integer m such that $x_n \leq y_n$ for all $n \geq m$, then $\lim x_n \leq \lim y_n$.

Proof. Let $z_n := y_n - x_n$.

Theorem

Let (x_n) be a convergent sequence of real numbers and there exists a positive integer m such that $x_n \geq 0$ for all $n \geq m$. Then $\lim x_n \geq 0$.

Corollary

If (x_n) and (y_n) are convergent sequences of real numbers and if there is a positive integer m such that $x_n \leq y_n$ for all $n \geq m$, then $\lim x_n \leq \lim y_n$.

Proof. Let $z_n := y_n - x_n$. Then (z_n) is convergent sequence of real numbers such that $z_n \geq 0$ for all $n \geq m$. It then follows from the preceding theorem that

$$\lim z_n = \lim(y_n - x_n) = \lim y_n - \lim x_n \geq 0.$$

Theorem

Sandwich Theorem. *Let $(a_n), (b_n), (c_n)$ be three sequences of real numbers and there is a natural number m such that*

$$a_n \leq b_n \leq c_n \text{ for all } n \geq m.$$

If $\lim a_n = \lim c_n = \ell$, then (b_n) is convergent and $\lim b_n = \ell$.

Theorem

Sandwich Theorem. *Let $(a_n), (b_n), (c_n)$ be three sequences of real numbers and there is a natural number m such that*

$$a_n \leq b_n \leq c_n \text{ for all } n \geq m.$$

If $\lim a_n = \lim c_n = \ell$, then (b_n) is convergent and $\lim b_n = \ell$.

Examples:

(i) $\lim_{n \rightarrow \infty} \frac{\cos n}{n}$

Theorem

Sandwich Theorem. Let $(a_n), (b_n), (c_n)$ be three sequences of real numbers and there is a natural number m such that

$$a_n \leq b_n \leq c_n \text{ for all } n \geq m.$$

If $\lim a_n = \lim c_n = \ell$, then (b_n) is convergent and $\lim b_n = \ell$.

Examples:

(i) $\lim_{n \rightarrow \infty} \frac{\cos n}{n}$

$-1 \leq \cos n \leq 1$. Therefore $-\frac{1}{n} \leq \frac{\cos n}{n} \leq \frac{1}{n}$ and $\lim_{n \rightarrow \infty} \frac{\cos n}{n} = 0$.

Theorem

Sandwich Theorem. Let $(a_n), (b_n), (c_n)$ be three sequences of real numbers and there is a natural number m such that

$$a_n \leq b_n \leq c_n \text{ for all } n \geq m.$$

If $\lim a_n = \lim c_n = \ell$, then (b_n) is convergent and $\lim b_n = \ell$.

Examples:

(i) $\lim_{n \rightarrow \infty} \frac{\cos n}{n}$
 $-1 \leq \cos n \leq 1$. Therefore $-\frac{1}{n} \leq \frac{\cos n}{n} \leq \frac{1}{n}$ and $\lim_{n \rightarrow \infty} \frac{\cos n}{n} = 0$.

(ii) $\lim_{n \rightarrow \infty} \frac{1}{2^n} = 0$
as $0 \leq \frac{1}{2^n} \leq \frac{1}{n}$

Theorem

Sandwich Theorem. Let $(a_n), (b_n), (c_n)$ be three sequences of real numbers and there is a natural number m such that

$$a_n \leq b_n \leq c_n \text{ for all } n \geq m.$$

If $\lim a_n = \lim c_n = \ell$, then (b_n) is convergent and $\lim b_n = \ell$.

Examples:

(i) $\lim_{n \rightarrow \infty} \frac{\cos n}{n}$

$-1 \leq \cos n \leq 1$. Therefore $-\frac{1}{n} \leq \frac{\cos n}{n} \leq \frac{1}{n}$ and $\lim_{n \rightarrow \infty} \frac{\cos n}{n} = 0$.

(ii) $\lim_{n \rightarrow \infty} \frac{1}{2^n} = 0$

as $0 \leq \frac{1}{2^n} \leq \frac{1}{n}$

(iii) $\lim_{n \rightarrow \infty} (-1)^n \frac{1}{n} = 0$.

Examples (continued)

(iv) Let $a_n := \frac{n^3 + 3n^2 + 1}{n^4 + 8n^2 + 2}$ for $n \in \mathbb{N}$. Then $a_n \rightarrow 0$,

(iii) Let $a_n := \frac{1}{n} \sin\left(\frac{1}{n}\right)$ for $n \in \mathbb{N}$. Then $a_n \rightarrow 0$,

Examples (continued)

(iv) Let $a_n := \frac{n^3 + 3n^2 + 1}{n^4 + 8n^2 + 2}$ for $n \in \mathbb{N}$. Then $a_n \rightarrow 0$,

$$\text{since } 0 \leq a_n \leq \frac{n^3 + 3n^2 + 1}{n^4} \leq \frac{1}{n} + \frac{3}{n^2} + \frac{1}{n^4} \rightarrow 0.$$

(iii) Let $a_n := \frac{1}{n} \sin\left(\frac{1}{n}\right)$ for $n \in \mathbb{N}$. Then $a_n \rightarrow 0$,

Examples (continued)

(iv) Let $a_n := \frac{n^3 + 3n^2 + 1}{n^4 + 8n^2 + 2}$ for $n \in \mathbb{N}$. Then $a_n \rightarrow 0$,

$$\text{since } 0 \leq a_n \leq \frac{n^3 + 3n^2 + 1}{n^4} \leq \frac{1}{n} + \frac{3}{n^2} + \frac{1}{n^4} \rightarrow 0.$$

(iii) Let $a_n := \frac{1}{n} \sin\left(\frac{1}{n}\right)$ for $n \in \mathbb{N}$. Then $a_n \rightarrow 0$,

Examples (continued)

(iv) Let $a_n := \frac{n^3 + 3n^2 + 1}{n^4 + 8n^2 + 2}$ for $n \in \mathbb{N}$. Then $a_n \rightarrow 0$,

$$\text{since } 0 \leq a_n \leq \frac{n^3 + 3n^2 + 1}{n^4} \leq \frac{1}{n} + \frac{3}{n^2} + \frac{1}{n^4} \rightarrow 0.$$

(iii) Let $a_n := \frac{1}{n} \sin\left(\frac{1}{n}\right)$ for $n \in \mathbb{N}$. Then $a_n \rightarrow 0$,

$$\text{since } |a_n| \leq \frac{1}{n} \rightarrow 0.$$