

# Triple Integrals in Rectangular and Cylindrical Coordinates

Devika S

Department of Mathematics  
BITS Pilani, K K Birla Goa Campus

November 13, 2024



## Examples - Triple Integrals

**Example 1:** Let  $D$  be the region bounded by the paraboloids  $z = 8 - x^2 - y^2$  and  $z = x^2 + y^2$ . Write six different triple iterated integrals for the volume of  $D$ . Evaluate one of the integrals.

## Examples - Triple Integrals

**Example 1:** Let  $D$  be the region bounded by the paraboloids  $z = 8 - x^2 - y^2$  and  $z = x^2 + y^2$ . Write six different triple iterated integrals for the volume of  $D$ . Evaluate one of the integrals.

**Solution:** Let us first sketch the region and set up the integral.

## Examples - Triple Integrals

**Example 1:** Let  $D$  be the region bounded by the paraboloids  $z = 8 - x^2 - y^2$  and  $z = x^2 + y^2$ . Write six different triple iterated integrals for the volume of  $D$ . Evaluate one of the integrals.

**Solution:** Let us first sketch the region and set up the integral.

Case 1:  $dzdydx$

## Examples - Triple Integrals

**Example 1:** Let  $D$  be the region bounded by the paraboloids  $z = 8 - x^2 - y^2$  and  $z = x^2 + y^2$ . Write six different triple iterated integrals for the volume of  $D$ . Evaluate one of the integrals.

**Solution:** Let us first sketch the region and set up the integral.

**Case 1:**  $dzdydx$  In this case,

$$V = \int_{-2}^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} \int_{x^2+y^2}^{8-x^2-y^2} dzdydx = 16\pi.$$

# Examples - Triple Integrals

**Example 1:** Let  $D$  be the region bounded by the paraboloids  $z = 8 - x^2 - y^2$  and  $z = x^2 + y^2$ . Write six different triple iterated integrals for the volume of  $D$ . Evaluate one of the integrals.

**Solution:** Let us first sketch the region and set up the integral.

**Case 1:**  $dzdydx$  In this case,

$$V = \int_{-2}^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} \int_{x^2+y^2}^{8-x^2-y^2} dzdydx = 16\pi.$$

**Case 2:**  $dzdx dy$

# Examples - Triple Integrals

**Example 1:** Let  $D$  be the region bounded by the paraboloids  $z = 8 - x^2 - y^2$  and  $z = x^2 + y^2$ . Write six different triple iterated integrals for the volume of  $D$ . Evaluate one of the integrals.

**Solution:** Let us first sketch the region and set up the integral.

**Case 1:**  $dzdydx$  In this case,

$$V = \int_{-2}^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} \int_{x^2+y^2}^{8-x^2-y^2} dzdydx = 16\pi.$$

**Case 2:**  $dzdxdy$

$$V = \int_{-2}^2 \int_{-\sqrt{4-y^2}}^{\sqrt{4-y^2}} \int_{x^2+y^2}^{8-x^2-y^2} dzdxdy.$$

# Examples

Case 3:  $dydzdx$



# Examples

## Case 3: $dydzdx$

$$V = \int_{-2}^2 \int_{x^2}^4 \int_{-\sqrt{z-x^2}}^{\sqrt{z-x^2}} dy dz dx + \int_{-2}^2 \int_4^{8-x^2} \int_{-\sqrt{8-z-x^2}}^{\sqrt{8-z-x^2}} dy dz dx.$$

# Examples

Case 3:  $dydzdx$

$$V = \int_{-2}^2 \int_{x^2}^4 \int_{-\sqrt{z-x^2}}^{\sqrt{z-x^2}} dy dz dx + \int_{-2}^2 \int_4^{8-x^2} \int_{-\sqrt{8-z-x^2}}^{\sqrt{8-z-x^2}} dy dz dx.$$

Case 4:  $dydx dz$

# Examples

## Case 3: $dydzdx$

$$V = \int_{-2}^2 \int_{x^2}^4 \int_{-\sqrt{z-x^2}}^{\sqrt{z-x^2}} dydzdx + \int_{-2}^2 \int_4^{8-x^2} \int_{-\sqrt{8-z-x^2}}^{\sqrt{8-z-x^2}} dydzdx.$$

## Case 4: $dydx dz$

$$V = \int_0^4 \int_{-\sqrt{z}}^{\sqrt{z}} \int_{-\sqrt{z-x^2}}^{\sqrt{z-x^2}} dydx dz + \int_4^8 \int_{-\sqrt{8-z}}^{\sqrt{8-z}} \int_{-\sqrt{8-z-x^2}}^{\sqrt{8-z-x^2}} dydx dz.$$

# Examples

Case :5  $dx dz dy$

# Examples

Case :5  $dx dz dy$

$$V = \int_{-2}^2 \int_{y^2}^4 \int_{-\sqrt{z-y^2}}^{\sqrt{z-y^2}} dx dz dy + \int_{-2}^2 \int_4^{8-y^2} \int_{-\sqrt{8-z-y^2}}^{\sqrt{8-z-y^2}} dx dz dy.$$

# Examples

Case 5:  $dx dz dy$

$$V = \int_{-2}^2 \int_{y^2}^4 \int_{-\sqrt{z-y^2}}^{\sqrt{z-y^2}} dx dz dy + \int_{-2}^2 \int_4^{8-y^2} \int_{-\sqrt{8-z-y^2}}^{\sqrt{8-z-y^2}} dx dz dy.$$

Case 6:  $dx dy dz$

# Examples

Case 5:  $dx dz dy$

$$V = \int_{-2}^2 \int_{y^2}^4 \int_{-\sqrt{z-y^2}}^{\sqrt{z-y^2}} dx dz dy + \int_{-2}^2 \int_4^{8-y^2} \int_{-\sqrt{8-z-y^2}}^{\sqrt{8-z-y^2}} dx dz dy.$$

Case 6:  $dx dy dz$

$$V = \int_0^4 \int_{-\sqrt{z}}^{\sqrt{z}} \int_{-\sqrt{z-y^2}}^{\sqrt{z-y^2}} dx dy dz + \int_4^8 \int_{-\sqrt{8-z}}^{\sqrt{8-z}} \int_{-\sqrt{8-z-y^2}}^{\sqrt{8-z-y^2}} dx dy dz.$$

# Examples

**Example 2:** Rewrite the following integral as an equivalent iterated integral in the order  $dy \, dz \, dx$  and  $dx \, dy \, dz$ .

$$\int_{-1}^1 \int_{x^2}^1 \int_0^{1-y} F(x, y, z) \, dz \, dy \, dx.$$



## Examples

**Example 2:** Rewrite the following integral as an equivalent iterated integral in the order  $dy \, dz \, dx$  and  $dx \, dy \, dz$ .

$$\int_{-1}^1 \int_{x^2}^1 \int_0^{1-y} F(x, y, z) \, dz \, dy \, dx.$$

**Solution:** First we need to sketch the region of integration:

$$-1 \leq x \leq 1; \, x^2 \leq y \leq 1; \, 0 \leq z \leq 1 - y.$$


# Examples

**Example 2:** Rewrite the following integral as an equivalent iterated integral in the order  $dy \, dz \, dx$  and  $dx \, dy \, dz$ .

$$\int_{-1}^1 \int_{x^2}^1 \int_0^{1-y} F(x, y, z) \, dz \, dy \, dx.$$

**Solution:** First we need to sketch the region of integration:

$$-1 \leq x \leq 1; \quad x^2 \leq y \leq 1; \quad 0 \leq z \leq 1 - y.$$



# Examples

The integral in the order  $dy \, dz \, dx$ :

$$\int_{-1}^1 \int_0^{1-x^2} \int_{x^2}^{1-z} dy \, dz \, dx.$$

# Examples

The integral in the order  $dy \, dz \, dx$ :

$$\int_{-1}^1 \int_0^{1-x^2} \int_{x^2}^{1-z} dy \, dz \, dx.$$

The integral in the order  $dx \, dy \, dz$ :

$$\int_0^1 \int_0^{1-z} \int_{-\sqrt{y}}^{\sqrt{y}} dx \, dy \, dz.$$

# Cylindrical Coordinates

The three dimensional analogues of the polar coordinates substitution are the **cylindrical** and **spherical coordinates** systems.

# Cylindrical Coordinates

The three dimensional analogues of the polar coordinates substitution are the **cylindrical** and **spherical coordinates** systems.

## Cylindrical Coordinates

Cylindrical coordinates represent a point  $P$  in space by ordered triples  $(r, \theta, z)$  in which

- $r$  and  $\theta$  are polar coordinates for the vertical projection of  $P$  on the  $xy$ -plane
- $z$  is the rectangular vertical coordinate.

# Cylindrical Coordinates

The three dimensional analogues of the polar coordinates substitution are the **cylindrical** and **spherical coordinates** systems.

## Cylindrical Coordinates


Cylindrical coordinates represent a point  $P$  in space by ordered triples  $(r, \theta, z)$  in which

- $r$  and  $\theta$  are polar coordinates for the vertical projection of  $P$  on the  $xy$ -plane
- $z$  is the rectangular vertical coordinate.

Equations relating rectangular  $(x, y, z)$  and cylindrical  $(r, \theta, z)$  coordinates

$$x = r \cos \theta, \quad y = r \sin \theta, \quad z = z,$$
$$r^2 = x^2 + y^2, \quad \tan \theta = y/x.$$

# Cylindrical Coordinates






# Examples

**Example 1.** Plot the point with cylindrical coordinates  $(2, 2\pi/3, 1)$  and find its rectangular coordinates.

# Examples

**Example 1.** Plot the point with cylindrical coordinates  $(2, 2\pi/3, 1)$  and find its rectangular coordinates.


**Solution:**



# Examples

**Example 1.** Plot the point with cylindrical coordinates  $(2, 2\pi/3, 1)$  and find its rectangular coordinates.

**Solution:**



$$x = 2 \cos(2\pi/3) = 2(-1/2) = -1$$

$$y = 2 \sin(2\pi/3) = 2(\sqrt{3}/2) = \sqrt{3}$$

$$z = 1$$

Rectangular coordinates:  
 $(-1, \sqrt{3}, 1)$

# Examples

**Example 2:** Find cylindrical coordinates of the point with rectangular coordinates  $(3, -3, -7)$ .

# Examples

**Example 2:** Find cylindrical coordinates of the point with rectangular coordinates  $(3, -3, -7)$ .

**Solution:**

$$r = \sqrt{3^2 + (-3)^2} = 3\sqrt{2}$$

$$\tan \theta = -1 \implies \theta = \frac{7\pi}{4} + 2n\pi$$

$$z = -7$$

Therefore one set of cylindrical coordinates is  $(3\sqrt{2}, \frac{7\pi}{4}, -7)$ . Another is  $(3\sqrt{2}, \frac{-\pi}{4}, -7)$ . As with polar coordinates, there are infinitely many choices.

Surface whose equation in cylindrical coordinates is  $r = c$   
and  $z = r$

Surface whose equation in cylindrical coordinates is  $r = c$  and  $z = r$




Figure:  $r = c$ , a cylinder

Surface whose equation in cylindrical coordinates is  $r = c$  and  $z = r$




Figure:  $r = c$ , a cylinder





Figure:  $z = r$ , a cone



# Constant-coordinate equations in cylindrical coordinates



## How to find the limits in cylindrical coordinates

When computing triple integrals over a region  $D$  in cylindrical coordinates, we partition the region into  $n$  small cylindrical wedges, rather than into rectangular boxes. In cylindrical coordinates the volume of the wedge is approximated by the product

$$\Delta V = \Delta z r \Delta r \Delta \theta.$$

# How to find the limits in cylindrical coordinates

When computing triple integrals over a region  $D$  in cylindrical coordinates, we partition the region into  $n$  small cylindrical wedges, rather than into rectangular boxes. In cylindrical coordinates the volume of the wedge is approximated by the product

$$\Delta V = \Delta z r \Delta r \Delta \theta.$$


To find the limits for the triple integral  $\iiint_D f dV$ :

- First sketch the solid  $D$  in three space.
- Find the  $z$  limits of the region.
- Find the projection of the solid  $R$  on the  $XY$  plane, from the projected region find the coordinates  $r, \theta$  as one would do for polar coordinates.

Once the limits are found then substitute  $dV$  or  $dz dy dx$  with  $rdz dr d\theta$  to convert the integral.


# Step 1

**Sketch** the region  $D$  along with its projection  $R$  on the  $xy$ -plane. Label the surrounding faces and curves that bound  $D$  and  $R$ .




## Step 2

Draw a line  $M$  through a typical point  $(r, \theta)$  of  $R$  parallel to the  $z$  axis.



## Step 3

Draw a ray  $L$  through  $(r, \theta)$  from the origin.



# Examples

The integral is

$$\iiint_D F(x, y, z) dV = \int_{\alpha}^{\beta} \int_{h_1(\theta)}^{h_2(\theta)} \int_{g_1(r, \theta)}^{g_2(r, \theta)} F(r \cos \theta, r \sin \theta, z) r dz dr d\theta.$$

# Examples

The integral is

$$\iiint_D F(x, y, z) dV = \int_{\alpha}^{\beta} \int_{h_1(\theta)}^{h_2(\theta)} \int_{g_1(r, \theta)}^{g_2(r, \theta)} F(r \cos \theta, r \sin \theta, z) r dz dr d\theta.$$

or

$$\iiint_D f(r, \theta, z) dV = \int_{\alpha}^{\beta} \int_{h_1(\theta)}^{h_2(\theta)} \int_{g_1(r, \theta)}^{g_2(r, \theta)} f(r, \theta, z) r dz dr d\theta.$$



# Examples

The integral is

$$\iiint_D F(x, y, z) dV = \int_{\alpha}^{\beta} \int_{h_1(\theta)}^{h_2(\theta)} \int_{g_1(r, \theta)}^{g_2(r, \theta)} F(r \cos \theta, r \sin \theta, z) r dz dr d\theta.$$

or

$$\iiint_D f(r, \theta, z) dV = \int_{\alpha}^{\beta} \int_{h_1(\theta)}^{h_2(\theta)} \int_{g_1(r, \theta)}^{g_2(r, \theta)} f(r, \theta, z) r dz dr d\theta.$$

**Example 1.** The domain  $D$  is the solid right cylinder whose base is the region in the  $xy$  plane that lies inside the cardioid  $r = 1 + \cos \theta$  and outside the circle  $r = 1$  and whose top lies on the plane  $z = 4$ . Find the volume of the solid.

# Examples

The integral is

$$\iiint_D F(x, y, z) dV = \int_{\alpha}^{\beta} \int_{h_1(\theta)}^{h_2(\theta)} \int_{g_1(r, \theta)}^{g_2(r, \theta)} F(r \cos \theta, r \sin \theta, z) r dz dr d\theta.$$


or

$$\iiint_D f(r, \theta, z) dV = \int_{\alpha}^{\beta} \int_{h_1(\theta)}^{h_2(\theta)} \int_{g_1(r, \theta)}^{g_2(r, \theta)} f(r, \theta, z) r dz dr d\theta.$$


**Example 1.** The domain  $D$  is the solid right cylinder whose base is the region in the  $xy$  plane that lies inside the cardioid  $r = 1 + \cos \theta$  and outside the circle  $r = 1$  and whose top lies on the plane  $z = 4$ . Find the volume of the solid.

**Solution:** First we sketch the solid.

# Examples




# Examples



The limits are

$$0 \leq z \leq 4$$

# Examples




The limits are

$$0 \leq z \leq 4$$

$$1 \leq r \leq 1 + \cos \theta$$

# Examples




The limits are

$$0 \leq z \leq 4$$

$$1 \leq r \leq 1 + \cos \theta$$

$$-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$$

# Examples



The limits are

$$0 \leq z \leq 4$$

$$1 \leq r \leq 1 + \cos \theta$$

$$-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$$

So the integral turns out to be:

$$\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_1^{1+\cos \theta} \int_0^4 r dz dr d\theta = 8 + \pi.$$