

Differentials & Extreme Values

Devika S

Department of Mathematics
BITS Pilani, K K Birla Goa Campus

October 25, 2024



ANNOUNCEMENT:

An additional class will be held this **Saturday (26 October 2024)** from **12:00 PM to 1:00 PM** in **LT3**.

Recall

Let f be a function with derivatives of all orders throughout some interval containing x_0 as an interior point. The **Taylor's series expansion** of $f(x)$ at $x = x_0$ is

Recall

Let f be a function with derivatives of all orders throughout some interval containing x_0 as an interior point. The **Taylor's series expansion** of $f(x)$ at $x = x_0$ is

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2}(x - x_0)^2 + \cdots$$

Recall

Let f be a function with derivatives of all orders throughout some interval containing x_0 as an interior point. The **Taylor's series expansion** of $f(x)$ at $x = x_0$ is

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2}(x - x_0)^2 + \cdots$$

The **linearization** of a differentiable function $f(x, y)$ at (x_0, y_0) is

$$L(x, y) = f(x_0, y_0) + f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0).$$

The approximation $f(x, y) \approx L(x, y)$ is called the **standard linear approximation** of the function f at (x_0, y_0) .

Recall

Let f be a function with derivatives of all orders throughout some interval containing x_0 as an interior point. The **Taylor's series expansion** of $f(x)$ at $x = x_0$ is

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2}(x - x_0)^2 + \cdots$$

The **linearization** of a differentiable function $f(x, y)$ at (x_0, y_0) is

$$L(x, y) = f(x_0, y_0) + f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0).$$

The approximation $f(x, y) \approx L(x, y)$ is called the **standard linear approximation** of the function f at (x_0, y_0) .

Linearization of $f(x, y)$ is a tangent-plane approximation. **Why?**

Recall

Let f be a function with derivatives of all orders throughout some interval containing x_0 as an interior point. The **Taylor's series expansion** of $f(x)$ at $x = x_0$ is

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2}(x - x_0)^2 + \cdots$$

The **linearization** of a differentiable function $f(x, y)$ at (x_0, y_0) is

$$L(x, y) = f(x_0, y_0) + f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0).$$

The approximation $f(x, y) \approx L(x, y)$ is called the **standard linear approximation** of the function f at (x_0, y_0) .

Linearization of $f(x, y)$ is a tangent-plane approximation. Why?

Write down the equation of tangent plane of the surface $z = f(x, y)$ and compare it with $L(x, y)$ of $f(x, y)$.

Error in the Standard Linear Approximation

The error $E(x, y) = f(x, y) - L(x, y)$ in the standard linear approximation of a function f can be estimated in the following way.

Error in the Standard Linear Approximation

The error $E(x, y) = f(x, y) - L(x, y)$ in the standard linear approximation of a function f can be estimated in the following way.

Error in the standard linear approximation

If f has continuous second partial derivatives throughout an open set containing a rectangle R centered at (x_0, y_0)

($R := \{(x, y) : |x - x_0| \leq h, |y - y_0| \leq k\}$) and if M is an upper bound for the values of $|f_{xx}|$, $|f_{xy}|$ and $|f_{yy}|$ in the rectangle R (that is, $M = \max\{|f_{xx}|, |f_{xy}|, |f_{yy}|\}$ on R), then for any $(x, y) \in R$,

$$|E(x, y)| \leq \frac{1}{2}M (|x - x_0| + |y - y_0|)^2.$$

Example

Find the linearization of $f(x, y) = \ln x + \ln y$ at $(1, 1)$ and the error of linearisation in the rectangle $R : |x - 1| \leq 0.2, |y - 1| \leq 0.2$.

Example

Find the linearization of $f(x, y) = \ln x + \ln y$ at $(1, 1)$ and the error of linearisation in the rectangle $R : |x - 1| \leq 0.2, |y - 1| \leq 0.2$.

Solution. Note that f is differentiable at $(1, 1)$.

$\nabla f|_{(1,1)} = (1/x, 1/y)|_{(1,1)} = (1, 1)$ and $L(x, y) = x + y - 2$.

Example

Find the linearization of $f(x, y) = \ln x + \ln y$ at $(1, 1)$ and the error of linearisation in the rectangle $R : |x - 1| \leq 0.2, |y - 1| \leq 0.2$.

Solution. Note that f is differentiable at $(1, 1)$.

$\nabla f|_{(1,1)} = (1/x, 1/y)|_{(1,1)} = (1, 1)$ and $L(x, y) = x + y - 2$.

Note that $f_{xx} = -1/x^2$, $f_{yy} = -1/y^2$ and $f_{xy} = 0$. Also, the second order partial derivatives are continuous in an open set containing R .

Example

Find the linearization of $f(x, y) = \ln x + \ln y$ at $(1, 1)$ and the error of linearisation in the rectangle $R : |x - 1| \leq 0.2, |y - 1| \leq 0.2$.

Solution. Note that f is differentiable at $(1, 1)$.

$\nabla f|_{(1,1)} = (1/x, 1/y)|_{(1,1)} = (1, 1)$ and $L(x, y) = x + y - 2$.

Note that $f_{xx} = -1/x^2$, $f_{yy} = -1/y^2$ and $f_{xy} = 0$. Also, the second order partial derivatives are continuous in an open set containing R .

And $|f_{xx}| = |-1/x^2| \leq \frac{1}{(0.8)^2} = 100/64$, since in the rectangle R , we have $0.8 \leq x \leq 1.2$ and $0.8 \leq y \leq 1.2$. Similarly, $|f_{yy}| \leq 100/64$.

Example

Find the linearization of $f(x, y) = \ln x + \ln y$ at $(1, 1)$ and the error of linearisation in the rectangle $R : |x - 1| \leq 0.2, |y - 1| \leq 0.2$.

Solution. Note that f is differentiable at $(1, 1)$.

$\nabla f|_{(1,1)} = (1/x, 1/y)|_{(1,1)} = (1, 1)$ and $L(x, y) = x + y - 2$.

Note that $f_{xx} = -1/x^2$, $f_{yy} = -1/y^2$ and $f_{xy} = 0$. Also, the second order partial derivatives are continuous in an open set containing R .

And $|f_{xx}| = |-1/x^2| \leq \frac{1}{(0.8)^2} = 100/64$, since in the rectangle R , we have $0.8 \leq x \leq 1.2$ and $0.8 \leq y \leq 1.2$. Similarly, $|f_{yy}| \leq 100/64$.

Since $M = \max \{|f_{xx}|, |f_{xy}|, |f_{yy}|\} = 100/64$, the error is given by

$$|E(x, y)| \leq \frac{1}{2}M(|x - x_0| + |y - y_0|)^2.$$

Hence,

$$|E(x, y)| \leq 1/2 \times 100/64 \times (0.2 + 0.2)^2 = 1/2 \times 100/64 \times 0.16 = 1/8.$$

Total differential for the functions of two variables

Definition

If we move from the point (x_0, y_0) to a point $(x_0 + dx, y_0 + dy)$ nearby then the resulting change in the linearization of the function

$$df = f_x(x_0, y_0)dx + f_y(x_0, y_0)dy$$

is called the **total differential** of f and dx, dy are called **differentials**.

Total differential for the functions of two variables

Definition

If we move from the point (x_0, y_0) to a point $(x_0 + dx, y_0 + dy)$ nearby then the resulting change in the linearization of the function

$$df = f_x(x_0, y_0)dx + f_y(x_0, y_0)dy$$

is called the **total differential** of f and dx, dy are called **differentials**.

Example 1. How accurately the volume of a right circular cylinder be measured if the error in the measurement of the base radius and the height is about 1 percent?

Total differential for the functions of two variables

Definition

If we move from the point (x_0, y_0) to a point $(x_0 + dx, y_0 + dy)$ nearby then the resulting change in the linearization of the function

$$df = f_x(x_0, y_0)dx + f_y(x_0, y_0)dy$$

is called the **total differential** of f and dx, dy are called **differentials**.

Example 1. How accurately the volume of a right circular cylinder be measured if the error in the measurement of the base radius and the height is about 1 percent? (**Volume** $=\pi r^2 h$; **atmost 3 percent of error in volume**)

Total differential for the functions of two variables

Definition

If we move from the point (x_0, y_0) to a point $(x_0 + dx, y_0 + dy)$ nearby then the resulting change in the linearization of the function

$$df = f_x(x_0, y_0)dx + f_y(x_0, y_0)dy$$

is called the **total differential** of f and dx, dy are called **differentials**.

Example 1. How accurately the volume of a right circular cylinder be measured if the error in the measurement of the base radius and the height is about 1 percent? (**Volume** $=\pi r^2 h$; **atmost 3 percent of error in volume**)

Example 2. A cylindrical steel storage tank is to be built that is 10ft tall and 4ft across in diameter. It is known that the steel will expand/contract with temperature changes; is the overall volume of the tank more sensitive to changes in the diameter or in the height of the tank?

Total differential for the functions of two variables

Definition

If we move from the point (x_0, y_0) to a point $(x_0 + dx, y_0 + dy)$ nearby then the resulting change in the linearization of the function

$$df = f_x(x_0, y_0)dx + f_y(x_0, y_0)dy$$

is called the **total differential** of f and dx, dy are called **differentials**.

Example 1. How accurately the volume of a right circular cylinder be measured if the error in the measurement of the base radius and the height is about 1 percent? (**Volume** $=\pi r^2 h$; **atmost 3 percent of error in volume**)

Example 2. A cylindrical steel storage tank is to be built that is 10ft tall and 4ft across in diameter. It is known that the steel will expand/contract with temperature changes; is the overall volume of the tank more sensitive to changes in the diameter or in the height of the tank? (**Volume of the tank is more sensitive to changes in radius than in height**).

Linearization in Three Variables

Linearization in Three Variables

The **linearization** $L(x, y, z)$ of $f(x, y, z)$ at a point (x_0, y_0, z_0) is

$$\begin{aligned} L(x, y, z) = & f(x_0, y_0, z_0) + f_x(x_0, y_0, z_0)(x - x_0) \\ & + f_y(x_0, y_0, z_0)(y - y_0) + f_z(x_0, y_0, z_0)(z - z_0). \end{aligned}$$

Linearization in Three Variables

Linearization in Three Variables

The **linearization** $L(x, y, z)$ of $f(x, y, z)$ at a point (x_0, y_0, z_0) is

$$\begin{aligned} L(x, y, z) = & f(x_0, y_0, z_0) + f_x(x_0, y_0, z_0)(x - x_0) \\ & + f_y(x_0, y_0, z_0)(y - y_0) + f_z(x_0, y_0, z_0)(z - z_0). \end{aligned}$$

Suppose that R is a closed rectangular solid centered at (x_0, y_0, z_0) and lying in an open region on which **the second partial derivatives of f are continuous**. Suppose also that $|f_{xx}|, |f_{xy}|, |f_{yy}|, |f_{yz}|$, and $|f_{xz}|$ are less than or equal to M throughout R (that is, $M = \max \{|f_{xx}|, |f_{xy}|, |f_{yy}|, |f_{yz}|, |f_{xz}|\}$), then the **error** in the standard linear approximation is bounded by

$$|E(x, y, z)| \leq \frac{1}{2}M (|x - x_0| + |y - y_0| + |z - z_0|)^2.$$

Linearization in Three Variables

Linearization in Three Variables

The **linearization** $L(x, y, z)$ of $f(x, y, z)$ at a point (x_0, y_0, z_0) is

$$\begin{aligned} L(x, y, z) = & f(x_0, y_0, z_0) + f_x(x_0, y_0, z_0)(x - x_0) \\ & + f_y(x_0, y_0, z_0)(y - y_0) + f_z(x_0, y_0, z_0)(z - z_0). \end{aligned}$$

Suppose that R is a closed rectangular solid centered at (x_0, y_0, z_0) and lying in an open region on which **the second partial derivatives of f are continuous**. Suppose also that $|f_{xx}|, |f_{xy}|, |f_{yy}|, |f_{yz}|, |f_{xz}|$ are less than or equal to M throughout R (that is, $M = \max \{|f_{xx}|, |f_{xy}|, |f_{yy}|, |f_{yz}|, |f_{xz}|\}$), then the **error** in the standard linear approximation is bounded by

$$|E(x, y, z)| \leq \frac{1}{2} M (|x - x_0| + |y - y_0| + |z - z_0|)^2.$$

The **total differential** is given by

$$df = f_x(x_0, y_0, z_0)dx + f_y(x_0, y_0, z_0)dy + f_z(x_0, y_0, z_0)dz.$$

Examples

- 1 Find the linearization of $f(x, y, z) = x^2 + y^2 + z^2$ at the point $(1, 1, 1)$.

Examples

- ① Find the linearization of $f(x, y, z) = x^2 + y^2 + z^2$ at the point $(1, 1, 1)$.

Solution. Here, f is differentiable and $f_x(1, 1, 1) = 2 = f_y(1, 1, 1) = f_z(1, 1, 1)$.

Examples

- ① Find the linearization of $f(x, y, z) = x^2 + y^2 + z^2$ at the point $(1, 1, 1)$.

Solution. Here, f is differentiable and $f_x(1, 1, 1) = 2 = f_y(1, 1, 1) = f_z(1, 1, 1)$. The linearization $L(x, y, z)$ of f is given by

$$\begin{aligned} L(x, y, z) &= f(1, 1, 1) + 2(x - 1) + 2(y - 1) + 2(z - 1) \\ &= 3 + 2x + 2y + 2z - 6 = 2x + 2y + 2z - 3. \end{aligned}$$

Examples

- ① Find the linearization of $f(x, y, z) = x^2 + y^2 + z^2$ at the point $(1, 1, 1)$.

Solution. Here, f is differentiable and $f_x(1, 1, 1) = 2 = f_y(1, 1, 1) = f_z(1, 1, 1)$. The linearization $L(x, y, z)$ of f is given by

$$\begin{aligned} L(x, y, z) &= f(1, 1, 1) + 2(x - 1) + 2(y - 1) + 2(z - 1) \\ &= 3 + 2x + 2y + 2z - 6 = 2x + 2y + 2z - 3. \end{aligned}$$

- ② Find an upper bound for the magnitude of the error in the approximation of $f(x, y, z) = xz - 3yz + 2$ at $P_0(1, 1, 2)$, where $R : |x - 1| \leq 0.01, |y - 1| \leq 0.01, |z - 2| \leq 0.02$.

Examples

- ① Find the linearization of $f(x, y, z) = x^2 + y^2 + z^2$ at the point $(1, 1, 1)$.

Solution. Here, f is differentiable and $f_x(1, 1, 1) = 2 = f_y(1, 1, 1) = f_z(1, 1, 1)$. The linearization $L(x, y, z)$ of f is given by

$$\begin{aligned} L(x, y, z) &= f(1, 1, 1) + 2(x - 1) + 2(y - 1) + 2(z - 1) \\ &= 3 + 2x + 2y + 2z - 6 = 2x + 2y + 2z - 3. \end{aligned}$$

- ② Find an upper bound for the magnitude of the error in the approximation of $f(x, y, z) = xz - 3yz + 2$ at $P_0(1, 1, 2)$, where $R : |x - 1| \leq 0.01, |y - 1| \leq 0.01, |z - 2| \leq 0.02$.

Solution. Here $f_{xx} = f_{yy} = f_{zz} = f_{xy} = 0, f_{yz} = -3, f_{xz} = 1$. These derivatives are continuous in an open region containing R .

Examples

- ① Find the linearization of $f(x, y, z) = x^2 + y^2 + z^2$ at the point $(1, 1, 1)$.

Solution. Here, f is differentiable and $f_x(1, 1, 1) = 2 = f_y(1, 1, 1) = f_z(1, 1, 1)$. The linearization $L(x, y, z)$ of f is given by

$$\begin{aligned} L(x, y, z) &= f(1, 1, 1) + 2(x - 1) + 2(y - 1) + 2(z - 1) \\ &= 3 + 2x + 2y + 2z - 6 = 2x + 2y + 2z - 3. \end{aligned}$$

- ② Find an upper bound for the magnitude of the error in the approximation of $f(x, y, z) = xz - 3yz + 2$ at $P_0(1, 1, 2)$, where $R : |x - 1| \leq 0.01, |y - 1| \leq 0.01, |z - 2| \leq 0.02$.

Solution. Here $f_{xx} = f_{yy} = f_{zz} = f_{xy} = 0, f_{yz} = -3, f_{xz} = 1$. These derivatives are continuous in an open region containing R . Here, $M = \max \{|f_{xx}|, |f_{xy}|, |f_{yy}|, |f_{yz}|, |f_{xz}|\} = 3$. Then,

$$|E(x, y, z)| \leq \frac{1}{2}(3)(0.01 + 0.01 + 0.02)^2 = 0.0024.$$

Examples

- ③ Find an upper bound for the magnitude of the error of the linear approximation of the function $f(x, y) = 1 + y + x \cos y$ at the point $(0,0)$ for the rectangle $|x| \leq 0.2$ and $|y| \leq 0.2$.

Examples

- ③ Find an upper bound for the magnitude of the error of the linear approximation of the function $f(x, y) = 1 + y + x \cos y$ at the point $(0, 0)$ for the rectangle $|x| \leq 0.2$ and $|y| \leq 0.2$.

Solution. Observe that all the second order partial derivatives of f are continuous in an open region containing given rectangle.

Examples

- ③ Find an upper bound for the magnitude of the error of the linear approximation of the function $f(x, y) = 1 + y + x \cos y$ at the point $(0,0)$ for the rectangle $|x| \leq 0.2$ and $|y| \leq 0.2$.

Solution. Observe that all the second order partial derivatives of f are continuous in an open region containing given rectangle.

$$|f_{xx}| = 0, |f_{yy}| = |-x \cos y| \leq |x| \leq 0.2, |f_{xy}| = |-\sin y| \leq |y| \leq 0.2.$$

Examples

- ③ Find an upper bound for the magnitude of the error of the linear approximation of the function $f(x, y) = 1 + y + x \cos y$ at the point $(0,0)$ for the rectangle $|x| \leq 0.2$ and $|y| \leq 0.2$.

Solution. Observe that all the second order partial derivatives of f are continuous in an open region containing given rectangle.

$$|f_{xx}| = 0, |f_{yy}| = |-x \cos y| \leq |x| \leq 0.2, |f_{xy}| = |-\sin y| \leq |y| \leq 0.2.$$

So we can choose our bound M to be 0.2.

Examples

- ③ Find an upper bound for the magnitude of the error of the linear approximation of the function $f(x, y) = 1 + y + x \cos y$ at the point $(0, 0)$ for the rectangle $|x| \leq 0.2$ and $|y| \leq 0.2$.

Solution. Observe that all the second order partial derivatives of f are continuous in an open region containing given rectangle.

$$|f_{xx}| = 0, |f_{yy}| = |-x \cos y| \leq |x| \leq 0.2, |f_{xy}| = |-\sin y| \leq |y| \leq 0.2.$$

So we can choose our bound M to be 0.2.

Hence, the error is given by

$$1/2 \times 0.2 \times (0.2 + 0.2)^2 = 1/2 \times 0.032 = 0.016.$$

Extreme Values

Definition

Let $f(x, y)$ be defined on a region R containing the point (a, b) .

- ① $f(a, b)$ is a **local maximum** value of f if $f(a, b) \geq f(x, y)$ for all domain points (x, y) in an open disk centered at (a, b) .
- ② $f(a, b)$ is a **local minimum** value of f if $f(a, b) \leq f(x, y)$ for all domain points (x, y) in an open disk centered at (a, b) .

Extreme Values


Definition

Let $f(x, y)$ be defined on a region R containing the point (a, b) .



- ① $f(a, b)$ is a **local maximum** value of f if $f(a, b) \geq f(x, y)$ for all domain points (x, y) in an open disk centered at (a, b) .
- ② $f(a, b)$ is a **local minimum** value of f if $f(a, b) \leq f(x, y)$ for all domain points (x, y) in an open disk centered at (a, b) .

The local maxima and local minima are together some time called **local extrema values** or **relative extrema**.

Local maximum and minimum



Figures



First Derivative Test for Local Extreme Values

Theorem (First Derivative Test for Local Extreme Values)

If $f(x, y)$ has a local maximum or minimum value at an interior point (a, b) of its domain and if the first partial derivatives exist there, then

$$f_x(a, b) = 0 \text{ and } f_y(a, b) = 0.$$

First Derivative Test for Local Extreme Values

Theorem (First Derivative Test for Local Extreme Values)

If $f(x, y)$ has a local maximum or minimum value at an interior point (a, b) of its domain and if the first partial derivatives exist there, then

$$f_x(a, b) = 0 \text{ and } f_y(a, b) = 0.$$

Proof: Let us say that the interior point (a, b) is a local extrema.

First Derivative Test for Local Extreme Values

Theorem (First Derivative Test for Local Extreme Values)

If $f(x, y)$ has a local maximum or minimum value at an interior point (a, b) of its domain and if the first partial derivatives exist there, then

$$f_x(a, b) = 0 \text{ and } f_y(a, b) = 0.$$

Proof: Let us say that the interior point (a, b) is a local extrema. Then consider the single variable function $g(x) = f(x, b)$.

First Derivative Test for Local Extreme Values

Theorem (First Derivative Test for Local Extreme Values)

*If $f(x, y)$ has a **local maximum or minimum value** at an interior point (a, b) of its domain and if the first partial derivatives exist there, then*

$$f_x(a, b) = 0 \text{ and } f_y(a, b) = 0.$$

Proof: Let us say that the interior point (a, b) is a local extrema. Then consider the single variable function $g(x) = f(x, b)$. By the definition of the extrema, this single variable function $g(x)$ has a local extrema at the point $x = a$.

First Derivative Test for Local Extreme Values

Theorem (First Derivative Test for Local Extreme Values)

If $f(x, y)$ has a local maximum or minimum value at an interior point (a, b) of its domain and if the first partial derivatives exist there, then

$$f_x(a, b) = 0 \text{ and } f_y(a, b) = 0.$$

Proof: Let us say that the interior point (a, b) is a local extrema. Then consider the single variable function $g(x) = f(x, b)$. By the definition of the extrema, this single variable function $g(x)$ has a local extrema at the point $x = a$. Hence by the single variable theory we know $g'(a) = 0 = f_x(a, b)$.

First Derivative Test for Local Extreme Values

Theorem (First Derivative Test for Local Extreme Values)

If $f(x, y)$ has a local maximum or minimum value at an interior point (a, b) of its domain and if the first partial derivatives exist there, then

$$f_x(a, b) = 0 \text{ and } f_y(a, b) = 0.$$

Proof: Let us say that the interior point (a, b) is a local extrema. Then consider the single variable function $g(x) = f(x, b)$. By the definition of the extrema, this single variable function $g(x)$ has a local extrema at the point $x = a$. Hence by the single variable theory we know $g'(a) = 0 = f_x(a, b)$. A similar arguments with $h(y) = f(a, y)$ shows that $f_y(a, b) = 0$.

Critical Points

Definition

An **interior point** of the domain of a function f is called a **critical point** of the function if

- ① either both the partial f_x, f_y vanish at the point
- ② or at least one of the partial derivatives f_x and f_y does not exist at the point.

Critical Points

Definition

An **interior point** of the domain of a function f is called a **critical point** of the function if

- ① either both the partial f_x, f_y vanish at the point
 - ② or at least one of the partial derivatives f_x and f_y does not exist at the point.
- First Derivative Test says that the only points where a function $f(x, y)$ can assume extreme values are critical points and boundary points.

Critical Points

Definition

An **interior point** of the domain of a function f is called a **critical point** of the function if

- ① either both the partial f_x, f_y vanish at the point
- ② or at least one of the partial derivatives f_x and f_y does not exist at the point.

- First Derivative Test says that the only points where a function $f(x, y)$ can assume extreme values are critical points and boundary points.
- Not every critical point of a differentiable function is a local extremum.

Critical Points

Definition

An **interior point** of the domain of a function f is called a **critical point** of the function if

- 1 either both the partial f_x , f_y vanish at the point
- 2 or at least one of the partial derivatives f_x and f_y does not exist at the point.

- First Derivative Test says that the only points where a function $f(x, y)$ can assume extreme values are critical points and boundary points.
- Not every critical point of a differentiable function is a local extremum.

Just as in the one variable situation we could have two dimensional analogue of “inflection points”. These analogues are called “saddle points” of the function.

Saddle Points

Definition

A differentiable function $f(x, y)$ has a saddle point at a critical point (a, b) if in every open disk centered at (a, b) there are domain points (x, y) where $f(x, y) > f(a, b)$ and domain points (x, y) where $f(x, y) < f(a, b)$.

Saddle Points




Definition

A differentiable function $f(x, y)$ has a saddle point at a critical point (a, b) if in every open disk centered at (a, b) there are domain points (x, y) where $f(x, y) > f(a, b)$ and domain points (x, y) where $f(x, y) < f(a, b)$. The corresponding point $(a, b, f(a, b))$ on the surface $z = f(x, y)$ is called a saddle point of the surface.

Saddle Points

Definition

A differentiable function $f(x, y)$ has a saddle point at a critical point (a, b) if in every open disk centered at (a, b) there are domain points (x, y) where $f(x, y) > f(a, b)$ and domain points (x, y) where $f(x, y) < f(a, b)$. The corresponding point $(a, b, f(a, b))$ on the surface $z = f(x, y)$ is called a saddle point of the surface.



Examples

Example 1. Find local extreme values of $f(x, y) = x^2 + y^2$.

Examples

Example 1. Find local extreme values of $f(x, y) = x^2 + y^2$.

Solution: Since the domain of f is entire plane (so there are no boundary points) and the partial derivatives exist everywhere, the local extreme values can occur only where the first derivatives vanish.

Examples

Example 1. Find local extreme values of $f(x, y) = x^2 + y^2$.

Solution: Since the domain of f is entire plane (so there are no boundary points) and the partial derivatives exist everywhere, the local extreme values can occur only where the first derivatives vanish.

The first derivatives are $f_x = 2x$,
 $f_y = 2y$.

Examples

Example 1. Find local extreme values of $f(x, y) = x^2 + y^2$.

Solution: Since the domain of f is entire plane (so there are no boundary points) and the partial derivatives exist everywhere, the local extreme values can occur only where the first derivatives vanish.

The first derivatives are $f_x = 2x$,
 $f_y = 2y$.

The only possibility is the point $(0, 0)$. The function being always non-negative, the critical point $(0, 0)$ is a local minimum.


Examples

Example 1. Find local extreme values of $f(x, y) = x^2 + y^2$.

Solution: Since the domain of f is entire plane (so there are no boundary points) and the partial derivatives exist everywhere, the local extreme values can occur only where the first derivatives vanish.

The first derivatives are $f_x = 2x$,
 $f_y = 2y$.

The only possibility is the point $(0, 0)$. The function being always non-negative, the critical point $(0, 0)$ is a local minimum.



Examples

- Find the local extreme values of $f(x, y) = y^2 - x^2$.
- Find the critical points of $f(x, y) = x^2 + 2xy$.
- Find the critical points of $f(x, y) = y \sin x$.