

MATHEMATICS-I

ANUSHAYA MOHAPATRA

Department of Mathematics
BITS PILANI K K Birla Goa Campus, Goa

September 5, 2024

Lecture 13

Power series

Radius of Convergence

Corollary 0.1.

The convergence of the series $\sum a_n(x - a)^n$ is described by one of the following three possibilities:

- *There is a positive number R such that the series diverges for x with $|x - a| > R$ but converges absolutely for x with $|x - a| < R$. The series may or may not converge at either of the endpoints $x = a - R$ and $x = a + R$.*

Radius of Convergence

Corollary 0.1.

The convergence of the series $\sum a_n(x - a)^n$ is described by one of the following three possibilities:

- There is a positive number R such that the series diverges for x with $|x - a| > R$ but converges absolutely for x with $|x - a| < R$. The series may or may not converge at either of the endpoints $x = a - R$ and $x = a + R$.*
- The series converges absolutely for every x ($R = \infty$).*

Radius of Convergence

Corollary 0.1.

The convergence of the series $\sum a_n(x - a)^n$ is described by one of the following three possibilities:

- There is a positive number R such that the series diverges for x with $|x - a| > R$ but converges absolutely for x with $|x - a| < R$. The series may or may not converge at either of the endpoints $x = a - R$ and $x = a + R$.*
- The series converges absolutely for every x ($R = \infty$).*
- The series converges at $x = a$ and diverges elsewhere ($R = 0$).*

Definition 0.2.

- 1 The number R in the above Corollary is called the **radius of convergence** of the power series $\sum a_n(x - a)^n$.

Definition 0.2.

- 1 The number R in the above Corollary is called the **radius of convergence** of the power series $\sum a_n(x - a)^n$.
- 2 The interval of radius R centered at $x = a$ is called the **interval of convergence**. The interval of convergence may be open, closed, or half-open, depending on the particular series.

Examples

$$\bullet \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$$

Examples

- $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$.
- We have seen that by ratio test, if $|x| < 1$ the series converges absolutely.

Examples

- $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$.
- We have seen that by ratio test, if $|x| < 1$ the series converges absolutely.
- If $|x| > 1$, then $\{(-1)^{n-1} \frac{x^n}{n}\}$ is divergent and hence the series diverges.

Examples

- $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$.
- We have seen that by ratio test, if $|x| < 1$ the series converges absolutely.
- If $|x| > 1$, then $\{(-1)^{n-1} \frac{x^n}{n}\}$ is divergent and hence the series diverges.
- For $x = 1$, the series is an alternating harmonic series and hence it converges.

Examples

- $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$.
- We have seen that by ratio test, if $|x| < 1$ the series converges absolutely.
- If $|x| > 1$, then $\{(-1)^{n-1} \frac{x^n}{n}\}$ is divergent and hence the series diverges.
- For $x = 1$, the series is an alternating harmonic series and hence it converges.
- For $x = -1$ the series becomes negative of harmonic series hence it diverges.

Examples

- $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$.
- We have seen that by ratio test, if $|x| < 1$ the series converges absolutely.
- If $|x| > 1$, then $\{(-1)^{n-1} \frac{x^n}{n}\}$ is divergent and hence the series diverges.
- For $x = 1$, the series is an alternating harmonic series and hence it converges.
- For $x = -1$ the series becomes negative of harmonic series hence it diverges.
- Therefore, the radius of convergence is $R = 1$ and $(-1, 1]$ is the interval of convergence.

- Find the radius and interval of convergence of

$$\sum_{n=1}^{\infty} \frac{6^n}{n} (4x - 1)^n.$$

- Find the radius and interval of convergence of

$$\sum_{n=1}^{\infty} \frac{6^n}{n} (4x - 1)^n.$$

- Apply ratio test for absolute convergence.

- Find the radius and interval of convergence of

$$\sum_{n=1}^{\infty} \frac{6^n}{n} (4x - 1)^n.$$

- Apply ratio test for absolute convergence.

$$\left| \frac{u_{n+1}}{u_n} \right| \rightarrow 6|4x - 1|.$$

- Find the radius and interval of convergence of

$$\sum_{n=1}^{\infty} \frac{6^n}{n} (4x - 1)^n.$$

- Apply ratio test for absolute convergence.

$$\left| \frac{u_{n+1}}{u_n} \right| \rightarrow 6|4x - 1|.$$

- For $|x - 1/4| < 1/24$ the power series is convergent absolutely.

- Find the radius and interval of convergence of

$$\sum_{n=1}^{\infty} \frac{6^n}{n} (4x - 1)^n.$$

- Apply ratio test for absolute convergence.

$$\left| \frac{u_{n+1}}{u_n} \right| \rightarrow 6|4x - 1|.$$

- For $|x - 1/4| < 1/24$ the power series is convergent absolutely.
For $|x - 1/4| > 1/24$ the power series is not convergent.

- Find the radius and interval of convergence of

$$\sum_{n=1}^{\infty} \frac{6^n}{n} (4x - 1)^n.$$

- Apply ratio test for absolute convergence.

$$\left| \frac{u_{n+1}}{u_n} \right| \rightarrow 6|4x - 1|.$$

- For $|x - 1/4| < 1/24$ the power series is convergent absolutely.
For $|x - 1/4| > 1/24$ the power series is not convergent.
- For $x = 5/24$, the series converges by Alternating series test.

- Find the radius and interval of convergence of

$$\sum_{n=1}^{\infty} \frac{6^n}{n} (4x - 1)^n.$$

- Apply ratio test for absolute convergence.

$$\left| \frac{u_{n+1}}{u_n} \right| \rightarrow 6|4x - 1|.$$

- For $|x - 1/4| < 1/24$ the power series is convergent absolutely.
For $|x - 1/4| > 1/24$ the power series is not convergent.
- For $x = 5/24$, the series converges by Alternating series test.
- For $x = 7/24$, the series is divergent

- Find the radius and interval of convergence of

$$\sum_{n=1}^{\infty} \frac{6^n}{n} (4x - 1)^n.$$

- Apply ratio test for absolute convergence.

$$\left| \frac{u_{n+1}}{u_n} \right| \rightarrow 6|4x - 1|.$$

- For $|x - 1/4| < 1/24$ the power series is convergent absolutely.
For $|x - 1/4| > 1/24$ the power series is not convergent.
- For $x = 5/24$, the series converges by Alternating series test.
- For $x = 7/24$, the series is divergent
- Therefore, the radius of convergence is $R = 1/24$ and $[5/24, 7/24)$ is the interval of convergence

Operations on Power Series

Theorem 0.3 (Multiplication Theorem for Power Series).

If $A(x) = \sum a_n x^n$ and $B(x) = \sum b_n x^n$ converge absolutely for all $|x| < r$ and

$$c_n = a_n b_0 + a_{n-1} b_1 + a_{n-2} b_2 + \cdots + a_0 b_n = \sum_{k=0}^n a_{n-k} b_k,$$

then $\sum c_n x^n$ converges to $A(x)B(x)$ absolutely for $|x| < r$:

$$\left(\sum_{n=0}^{\infty} a_n x^n \right) \cdot \left(\sum_{n=0}^{\infty} b_n x^n \right) = \sum_{n=0}^{\infty} c_n x^n.$$

The Term-by-Term Differentiation Theorem

Theorem 0.4.

If $\sum a_n(x - a)^n$ has radius of convergence $R > 0$, it defines a function

$$f(x) = \sum_{n=0}^{\infty} a_n(x - a)^n \quad \text{on the interval } a - R < x < a + R.$$

This function f has derivatives of all orders inside the interval, and we obtain the derivatives by differentiating the original series term by term:

$$f'(x) = \sum_{n=1}^{\infty} n a_n(x - a)^{n-1}; \quad f''(x) = \sum_{n=2}^{\infty} n(n-1) a_n(x - a)^{n-2},$$

and so on. Each of these derived series converges at every point of the interval $a - R < x < a + R$.

The Term-by-Term Integration Theorem

Theorem 0.5.

Suppose that $f(x) = \sum_{n=0}^{\infty} a_n(x-a)^n$ converges for all $|x-a| < R$. Then $\sum_{n=0}^{\infty} a_n \frac{(x-a)^{n+1}}{n+1}$ converges for all $|x-a| < R$ and

$$\int f(x) dx = \sum_{n=0}^{\infty} a_n \frac{(x-a)^{n+1}}{n+1} + C$$

for $|x-a| < R$.

Example

By knowing that $\frac{1}{1-x} = 1 + x + x^2 + \dots$ for $|x| < 1$, show that the following:

$$\textcircled{1} \quad \frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + \dots = \sum_{n=1}^{\infty} nx^{n-1} \text{ for } |x| < 1.$$

Example

By knowing that $\frac{1}{1-x} = 1 + x + x^2 + \dots$ for $|x| < 1$, show that the following:

$$\textcircled{1} \quad \frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + \dots = \sum_{n=1}^{\infty} nx^{n-1} \text{ for } |x| < 1.$$

$$\textcircled{2} \quad \frac{1}{(1-x)^3} = 1 + 3x + 6x^2 + \dots = \sum_{n=2}^{\infty} \frac{n(n-1)}{2} x^{n-2} \text{ for } |x| < 1.$$

Example

By knowing that $\frac{1}{1-x} = 1 + x + x^2 + \dots$ for $|x| < 1$, show that the following:

$$\textcircled{1} \quad \frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + \dots = \sum_{n=1}^{\infty} nx^{n-1} \text{ for } |x| < 1.$$

$$\textcircled{2} \quad \frac{1}{(1-x)^3} = 1 + 3x + 6x^2 + \dots = \sum_{n=2}^{\infty} \frac{n(n-1)}{2} x^{n-2} \text{ for } |x| < 1.$$

$$\textcircled{3} \quad \ln(1-x) = x + \frac{x^2}{2} + \frac{x^3}{3} + \dots = \sum_{n=1}^{\infty} \frac{1}{n} x^n \text{ for } |x| < 1.$$

Questions

- ❶ Find a power series representation for the following function and determine its interval of convergence.

$$f(x) = \frac{1}{1+x^3}$$

- ❷ Find the interval of convergence and radius of convergence:

- (a) $\sum_{n=1}^{\infty} (-3)^{n-1} \frac{(x-1)^n}{n}$

10.8 Taylor and Maclaurin Series

Introduction

If the power series $\sum_{n=1}^{\infty} a_n(x - a)^n$ has positive radius of convergence $R > 0$, then we know that

$$g(x) = \sum_{n=1}^{\infty} a_n(x - a)^n$$

is differentiable for infinitely many times on $(a - R, a + R)$.

But what about the other way around?

Introduction

If the power series $\sum_{n=1}^{\infty} a_n(x - a)^n$ has positive radius of convergence $R > 0$, then we know that

$$g(x) = \sum_{n=1}^{\infty} a_n(x - a)^n$$

is differentiable for infinitely many times on $(a - R, a + R)$.

But what about the other way around?

If a function $f(x)$ has derivatives of all orders on an interval, can it be expressed as a power series on at least some part of that interval?

Introduction

If the power series $\sum_{n=1}^{\infty} a_n(x - a)^n$ has positive radius of convergence $R > 0$, then we know that

$$g(x) = \sum_{n=1}^{\infty} a_n(x - a)^n$$

is differentiable for infinitely many times on $(a - R, a + R)$.

But what about the other way around?

If a function $f(x)$ has derivatives of all orders on an interval, can it be expressed as a power series on at least some part of that interval?

And if it can, what are its coefficients?

Taylor and Maclaurin Series

Definition: Let f be a function with derivatives of all orders throughout some interval containing a as an interior point.

Taylor and Maclaurin Series

Definition: Let f be a function with derivatives of all orders throughout some interval containing a as an interior point. Then the **Taylor series generated by f at $x = a$** is

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!} (x-a)^k = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!} (x-a)^2 + \cdots + \frac{f^{(n)}(a)}{n!} (x-a)^n + \cdots$$

Taylor and Maclaurin Series

Definition: the **Maclaurin series generated by** f is

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(0)}{k!} x^k = f(0) + f'(0)x + \frac{f''(0)}{2!} x^2 + \cdots + \frac{f^{(n)}(0)}{n!} x^n + \cdots ,$$

the Taylor series generated by f at $x = 0$.