

MATH F111- Mathematics I

Saranya G. Nair
Department of Mathematics

BITS Pilani

August 9, 2024



Notations

\mathbb{N} - Set of Natural numbers

\mathbb{Q} - Set of rational numbers

\mathbb{R} - Set of real numbers

\forall - For all

\exists - There exists

Intervals

Definition

A subset I of \mathbb{R} is said to be an interval if $a, b \in I$ and $a < x < b \implies x \in I$.

Let $a, b \in \mathbb{R}$ and $a < b$.

- $(a, b) := \{x \in \mathbb{R} : a < x < b\}$ (open interval)
- $[a, b] := \{x \in \mathbb{R} : a \leq x \leq b\}$ (closed interval)
- $[a, b) := \{x \in \mathbb{R} : a \leq x < b\}$ and $(a, b] := \{x \in \mathbb{R} : a < x \leq b\}$ are half-open (or half-closed) intervals.
- $(a, \infty) := \{x \in \mathbb{R} : x > a\}$ and $(-\infty, a) := \{x \in \mathbb{R} : x < a\}$ are infinite open intervals.
- $[a, \infty) := \{x \in \mathbb{R} : x \geq a\}$ and $(-\infty, a] := \{x \in \mathbb{R} : x \leq a\}$ are infinite closed intervals.

Let $a \in \mathbb{R}$ and $\epsilon > 0$. Then $(a - \epsilon, a + \epsilon)$ is called the **ϵ -neighborhood of a** .

Sequences

Definition

A **sequence of real numbers** (or a **sequence in \mathbb{R}**) is a function $x : \mathbb{N} \rightarrow \mathbb{R}$.

Sequences

Definition

A **sequence of real numbers** (or a **sequence in \mathbb{R}**) is a function $x : \mathbb{N} \rightarrow \mathbb{R}$.

- If $x : \mathbb{N} \rightarrow \mathbb{R}$ is a sequence, we will usually denote the value of $x(n)$ by the symbol x_n .

Sequences

Definition

A **sequence of real numbers** (or a **sequence in \mathbb{R}**) is a function $x : \mathbb{N} \rightarrow \mathbb{R}$.

- If $x : \mathbb{N} \rightarrow \mathbb{R}$ is a sequence, we will usually denote the value of $x(n)$ by the symbol x_n .
- The values x_n are also called the terms or the elements of the sequence and x_n (that is, the value of x at n) is called the n -th term of the sequence.

We will denote this sequence by the notations

$$(x_n), \quad \text{or} \quad (x_n : n \in \mathbb{N}).$$

In this course, we will consider only Real sequences.

Example

- $(n : n \in \mathbb{N}) = (1, 2, 3, 4, \dots)$

Example

- $(n : n \in \mathbb{N}) = (1, 2, 3, 4, \dots)$
- $(1/n : n \in \mathbb{N}) = (1, 1/2, 1/3, 1/4, \dots)$

Example

- $(n : n \in \mathbb{N}) = (1, 2, 3, 4, \dots)$
- $(1/n : n \in \mathbb{N}) = (1, 1/2, 1/3, 1/4, \dots)$
- $(n^2 : n \in \mathbb{N}) = (1, 4, 9, 16, \dots)$

Example

- $(n : n \in \mathbb{N}) = (1, 2, 3, 4, \dots)$
- $(1/n : n \in \mathbb{N}) = (1, 1/2, 1/3, 1/4, \dots)$
- $(n^2 : n \in \mathbb{N}) = (1, 4, 9, 16, \dots)$
- *If $b \in \mathbb{R}$, the sequence (b, b, b, \dots) , all of whose terms equal b , is called the constant sequence b .*

Example

- $(n : n \in \mathbb{N}) = (1, 2, 3, 4, \dots)$
- $(1/n : n \in \mathbb{N}) = (1, 1/2, 1/3, 1/4, \dots)$
- $(n^2 : n \in \mathbb{N}) = (1, 4, 9, 16, \dots)$
- *If $b \in \mathbb{R}$, the sequence (b, b, b, \dots) , all of whose terms equal b , is called the constant sequence b .*
- $(2^n : n \in \mathbb{N}) = (2, 4, 8, 16, \dots)$

Example

- $(n : n \in \mathbb{N}) = (1, 2, 3, 4, \dots)$
- $(1/n : n \in \mathbb{N}) = (1, 1/2, 1/3, 1/4, \dots)$
- $(n^2 : n \in \mathbb{N}) = (1, 4, 9, 16, \dots)$
- *If $b \in \mathbb{R}$, the sequence (b, b, b, \dots) , all of whose terms equal b , is called the constant sequence b .*
- $(2^n : n \in \mathbb{N}) = (2, 4, 8, 16, \dots)$
- $((-1)^n : n \in \mathbb{N}) = (-1, 1, -1, 1, \dots)$

Example

- $(n : n \in \mathbb{N}) = (1, 2, 3, 4, \dots)$
- $(1/n : n \in \mathbb{N}) = (1, 1/2, 1/3, 1/4, \dots)$
- $(n^2 : n \in \mathbb{N}) = (1, 4, 9, 16, \dots)$
- If $b \in \mathbb{R}$, the sequence (b, b, b, \dots) , all of whose terms equal b , is called the constant sequence b .
- $(2^n : n \in \mathbb{N}) = (2, 4, 8, 16, \dots)$
- $((-1)^n : n \in \mathbb{N}) = (-1, 1, -1, 1, \dots)$
- $x_1 := 1, x_2 := 1$ and $x_n := x_{n-1} + x_{n-2}$ for $n \geq 3$:
 $(1, 1, 2, 3, 5, 8, 13, 21, 34, 55, \dots)$ This sequence is known as the *Fibonacci sequence*.

Bounded Sequences

A sequence (a_n) of real numbers is said to be **bounded above** if there is a real number α such that $a_n \leq \alpha$ for every $(\forall) n \in \mathbb{N}$.

Bounded Sequences

A sequence (a_n) of real numbers is said to be **bounded above** if there is a real number α such that $a_n \leq \alpha$ for every $(\forall) n \in \mathbb{N}$. eg. $(a_n) = -n$.

A sequence (a_n) of real numbers is said to be **bounded below** if there is a real number β such that $\beta \leq a_n$ for every $n \in \mathbb{N}$.

Bounded Sequences

A sequence (a_n) of real numbers is said to be **bounded above** if there is a real number α such that $a_n \leq \alpha$ for every $(\forall) n \in \mathbb{N}$. eg. $(a_n) = -n$.

A sequence (a_n) of real numbers is said to be **bounded below** if there is a real number β such that $\beta \leq a_n$ for every $n \in \mathbb{N}$. eg. $(a_n) = n^2$

A sequence (a_n) of real numbers is said to be **bounded** if there are real numbers α, β such that $\beta \leq a_n \leq \alpha$ for every $n \in \mathbb{N}$.

Bounded Sequences

A sequence (a_n) of real numbers is said to be **bounded above** if there is a real number α such that $a_n \leq \alpha$ for every $(\forall) n \in \mathbb{N}$. eg. $(a_n) = -n$.

A sequence (a_n) of real numbers is said to be **bounded below** if there is a real number β such that $\beta \leq a_n$ for every $n \in \mathbb{N}$. eg. $(a_n) = n^2$

A sequence (a_n) of real numbers is said to be **bounded** if there are real numbers α, β such that $\beta \leq a_n \leq \alpha$ for every $n \in \mathbb{N}$. eg. $(a_n) = \frac{1}{n}$, $(a_n) = (-1)^n$

If a sequence is not bounded, it is said to be **unbounded**. eg. $(a_n) = (-1)^n n$

- $(a_n) = \{1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n} \dots\}$

- $(a_n) = \{1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n} \dots\}$ approaches 0 as n gets large.
- $(a_n) = \{0, \frac{1}{2}, \frac{2}{3}, \dots, 1 - \frac{1}{n} \dots\}$

- $(a_n) = \{1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n} \dots\}$ approaches 0 as n gets large.
- $(a_n) = \{0, \frac{1}{2}, \frac{2}{3}, \dots, 1 - \frac{1}{n} \dots\}$ approaches 1 as n gets large.
- $(a_n) = \{\sqrt{1}, \sqrt{2}, \sqrt{3}, \dots, \sqrt{n} \dots\}$

- $(a_n) = \{1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n} \dots\}$ approaches 0 as n gets large.
- $(a_n) = \{0, \frac{1}{2}, \frac{2}{3}, \dots, 1 - \frac{1}{n} \dots\}$ approaches 1 as n gets large.
- $(a_n) = \{\sqrt{1}, \sqrt{2}, \sqrt{3}, \dots, \sqrt{n} \dots\}$ have terms that get larger than any number as n increases.

- $(a_n) = \{1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n} \dots\}$ approaches 0 as n gets large.
- $(a_n) = \{0, \frac{1}{2}, \frac{2}{3}, \dots, 1 - \frac{1}{n} \dots\}$ approaches 1 as n gets large.
- $(a_n) = \{\sqrt{1}, \sqrt{2}, \sqrt{3}, \dots, \sqrt{n} \dots\}$ have terms that get larger than any number as n increases.
- $(a_n) = \{1, -1, 1, -1, \dots\}$

- $(a_n) = \{1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n} \dots\}$ approaches 0 as n gets large.
- $(a_n) = \{0, \frac{1}{2}, \frac{2}{3}, \dots, 1 - \frac{1}{n} \dots\}$ approaches 1 as n gets large.
- $(a_n) = \{\sqrt{1}, \sqrt{2}, \sqrt{3}, \dots, \sqrt{n} \dots\}$ have terms that get larger than any number as n increases.
- $(a_n) = \{1, -1, 1, -1, \dots\}$ bounce back and forth between 1 and -1 , never approaching to a single value.

- $(a_n) = \{1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n} \dots\}$ approaches 0 as n gets large.
- $(a_n) = \{0, \frac{1}{2}, \frac{2}{3}, \dots, 1 - \frac{1}{n} \dots\}$ approaches 1 as n gets large.
- $(a_n) = \{\sqrt{1}, \sqrt{2}, \sqrt{3}, \dots, \sqrt{n} \dots\}$ have terms that get larger than any number as n increases.
- $(a_n) = \{1, -1, 1, -1, \dots\}$ bounce back and forth between 1 and -1 , never approaching to a single value.

Remark

Question: What do we mean by a sequence converges?

It says that if we go far enough out in the sequence, the difference between a_n and the limit of the sequence becomes less than any preselected number $\epsilon > 0$.

Let us see this with $(a_n) = \frac{1}{n}$.

- Can you find an integer N such that $|a_n - 0| < \frac{1}{2}, \forall n \geq N$?

Let us see this with $(a_n) = \frac{1}{n}$.

- Can you find an integer N such that $|a_n - 0| < \frac{1}{2}, \forall n \geq N$? Yes, choose $N = 3$.

Let us see this with $(a_n) = \frac{1}{n}$.

- Can you find an integer N such that $|a_n - 0| < \frac{1}{2}, \forall n \geq N$? Yes, choose $N = 3$.
- Can you find an integer N such that $|a_n - 0| < \frac{1}{10000}, \forall n \geq N$?

Let us see this with $(a_n) = \frac{1}{n}$.

- Can you find an integer N such that $|a_n - 0| < \frac{1}{2}, \forall n \geq N$? Yes, choose $N = 3$.
- Can you find an integer N such that $|a_n - 0| < \frac{1}{10000}, \forall n \geq N$? Yes, choose $N = 10001$.

Let us see this with $(a_n) = \frac{1}{n}$.

- Can you find an integer N such that $|a_n - 0| < \frac{1}{2}, \forall n \geq N$? Yes, choose $N = 3$.
- Can you find an integer N such that $|a_n - 0| < \frac{1}{10000}, \forall n \geq N$? Yes, choose $N = 10001$.
- For any preselected positive number, say $\epsilon > 0$, can you find an integer N such that $|a_n - 0| < \epsilon, \forall n \geq N$?

Let us see this with $(a_n) = \frac{1}{n}$.

- Can you find an integer N such that $|a_n - 0| < \frac{1}{2}, \forall n \geq N$? Yes, choose $N = 3$.
- Can you find an integer N such that $|a_n - 0| < \frac{1}{10000}, \forall n \geq N$? Yes, choose $N = 10001$.
- For any preselected positive number, say $\epsilon > 0$, can you find an integer N such that $|a_n - 0| < \epsilon, \forall n \geq N$? Yes, choose $N > \frac{1}{\epsilon}$. In particular choose $N = \lfloor \frac{1}{\epsilon} \rfloor + 1$.

The Limit of a Sequence

Definition

A sequence (a_n) in \mathbb{R} is said to converge to $\ell \in \mathbb{R}$, or ℓ is said to be a limit of (a_n) ,

if **for every** $\epsilon > 0$, **there exists** an integer $N \in \mathbb{N}$ such that

$$|a_n - \ell| < \epsilon, \text{ for all } n \geq N.$$

The Limit of a Sequence

Definition

A sequence (a_n) in \mathbb{R} is said to converge to $\ell \in \mathbb{R}$, or ℓ is said to be a limit of (a_n) ,

if **for every** $\epsilon > 0$, **there exists** an integer $N \in \mathbb{N}$ such that

$$|a_n - \ell| < \epsilon, \text{ for all } n \geq N.$$

ie,

$$a_n \in (\ell - \epsilon, \ell + \epsilon), \forall n \geq N.$$

The Limit of a Sequence

Definition

A sequence (a_n) in \mathbb{R} is said to converge to $\ell \in \mathbb{R}$, or ℓ is said to be a limit of (a_n) ,

if **for every** $\epsilon > 0$, **there exists** an integer $N \in \mathbb{N}$ such that

$$|a_n - \ell| < \epsilon, \text{ for all } n \geq N.$$

ie,

$$a_n \in (\ell - \epsilon, \ell + \epsilon), \forall n \geq N.$$

Remark

The choice of N depends on the value of ϵ .

When a sequence (a_n) has limit ℓ , we will use the notation

$$\lim a_n = \ell.$$

We will sometimes use the symbolism $a_n \rightarrow \ell$, which indicates the intuitive idea that the values a_n “approach” the number ℓ as $n \rightarrow \infty$.

- If a sequence has a limit, we say that the sequence is **convergent**
- If a sequence has no limit, we say that the sequence is **divergent**.

There is also a notion of divergence if the sequence is not bounded.

Definition

The sequence (a_n) **diverges to infinity** or

$$\lim_{n \rightarrow \infty} a_n = \infty$$

Definition

The sequence (a_n) **diverges to infinity** or

$$\lim_{n \rightarrow \infty} a_n = \infty$$

if for every number M there is an integer N such that $\forall n > N, a_n > M$.

Definition

The sequence (a_n) **diverges to infinity** or

$$\lim_{n \rightarrow \infty} a_n = \infty$$

if for every number M there is an integer N such that $\forall n > N, a_n > M$.
Similarly we say (a_n) **diverges to negative infinity** or

$$\lim_{n \rightarrow \infty} a_n = -\infty$$

if for every number m , there is an integer N such that $\forall n > N$, we have $a_n < m$.

The convergence of a sequence is unaltered if a finite number of its terms are replaced by some other terms.

The convergence of a sequence is unaltered if a finite number of its terms are replaced by some other terms.

Examples:

- (i) Let $a \in \mathbb{R}$ and $a_n := a$ for all $n \in \mathbb{N}$. Then $a_n \rightarrow a$. We can let $N := 1$ for any choice of ϵ .

The convergence of a sequence is unaltered if a finite number of its terms are replaced by some other terms.

Examples:

- (i) Let $a \in \mathbb{R}$ and $a_n := a$ for all $n \in \mathbb{N}$. Then $a_n \rightarrow a$. We can let $N := 1$ for any choice of ϵ .
- (ii) $a_n := 1/n$ for all $n \in \mathbb{N}$. Then $a_n \rightarrow 0$.

$$(iii) \ a_n := 2/(n^2 + 1)$$

(iii) $a_n := 2/(n^2 + 1)$

$$\left| \frac{2}{n^2 + 1} - 0 \right| = \frac{2}{n^2 + 1} < \frac{2}{n^2} \quad \text{for all } n \in \mathbb{N}.$$

(iii) $a_n := 2/(n^2 + 1)$

$$\left| \frac{2}{n^2 + 1} - 0 \right| = \frac{2}{n^2 + 1} < \frac{2}{n^2} \quad \text{for all } n \in \mathbb{N}.$$

Choose $N \in \mathbb{N}$ such that

$$N > \sqrt{\frac{2}{\epsilon}}.$$

Then

$$|a_n - 0| < \frac{2}{n^2} < \epsilon, \quad \forall n \geq N.$$

(iii) $a_n := 2/(n^2 + 1)$

$$\left| \frac{2}{n^2 + 1} - 0 \right| = \frac{2}{n^2 + 1} < \frac{2}{n^2} \quad \text{for all } n \in \mathbb{N}.$$

Choose $N \in \mathbb{N}$ such that

$$N > \sqrt{\frac{2}{\epsilon}}.$$

Then

$$|a_n - 0| < \frac{2}{n^2} < \epsilon, \quad \forall n \geq N.$$

Then $a_n \rightarrow 0$.

(iii) $a_n := 2/(n^2 + 1)$

$$\left| \frac{2}{n^2 + 1} - 0 \right| = \frac{2}{n^2 + 1} < \frac{2}{n^2} \quad \text{for all } n \in \mathbb{N}.$$

Choose $N \in \mathbb{N}$ such that

$$N > \sqrt{\frac{2}{\epsilon}}.$$

Then

$$|a_n - 0| < \frac{2}{n^2} < \epsilon, \quad \forall n \geq N.$$

Then $a_n \rightarrow 0$.

(iv) $a_n := 5/(3n + 1)$

(iii) $a_n := 2/(n^2 + 1)$

$$\left| \frac{2}{n^2 + 1} - 0 \right| = \frac{2}{n^2 + 1} < \frac{2}{n^2} \quad \text{for all } n \in \mathbb{N}.$$

Choose $N \in \mathbb{N}$ such that

$$N > \sqrt{\frac{2}{\epsilon}}.$$

Then

$$|a_n - 0| < \frac{2}{n^2} < \epsilon, \quad \forall n \geq N.$$

Then $a_n \rightarrow 0$.

(iv) $a_n := 5/(3n + 1)$

$$\frac{5}{3n + 1} < \frac{5}{3n} \quad \text{for all } n \in \mathbb{N}.$$

(iii) $a_n := 2/(n^2 + 1)$

$$\left| \frac{2}{n^2 + 1} - 0 \right| = \frac{2}{n^2 + 1} < \frac{2}{n^2} \quad \text{for all } n \in \mathbb{N}.$$

Choose $N \in \mathbb{N}$ such that

$$N > \sqrt{\frac{2}{\epsilon}}.$$

Then

$$|a_n - 0| < \frac{2}{n^2} < \epsilon, \quad \forall n \geq N.$$

Then $a_n \rightarrow 0$.

(iv) $a_n := 5/(3n + 1)$

$$\frac{5}{3n + 1} < \frac{5}{3n} \quad \text{for all } n \in \mathbb{N}.$$

Choose $N \in \mathbb{N}$ such that $N > 5/3\epsilon$. Then $|a_n - 0| < \frac{5}{3n} < \epsilon$ for all $n \geq N$. Then $a_n \rightarrow 0$.

Examples

(v). $\lim_{n \rightarrow \infty} r^n = 0$ for $|r| < 1$.

Examples

(v). $\lim_{n \rightarrow \infty} r^n = 0$ for $|r| < 1$.

Case 1. $r = 0$

In this case the sequence is $\{0, 0, 0, \dots\}$ which converges to 0.

Examples

(v). $\lim_{n \rightarrow \infty} r^n = 0$ for $|r| < 1$.

Case 1. $r = 0$

In this case the sequence is $\{0, 0, 0, \dots\}$ which converges to 0.

Case 2. $r \neq 0$ and $|r| < 1$.

Since $|r| < 1$, $\frac{1}{|r|} > 1$. Let $\frac{1}{|r|} = 1 + a$ where $a > 0$.

Examples

(v). $\lim_{n \rightarrow \infty} r^n = 0$ for $|r| < 1$.

Case 1. $r = 0$

In this case the sequence is $\{0, 0, 0, \dots\}$ which converges to 0.

Case 2. $r \neq 0$ and $|r| < 1$.

Since $|r| < 1$, $\frac{1}{|r|} > 1$. Let $\frac{1}{|r|} = 1 + a$ where $a > 0$. Then

$$|r^n - 0| = |r|^n = \frac{1}{(1 + a)^n}.$$

Examples

(v). $\lim_{n \rightarrow \infty} r^n = 0$ for $|r| < 1$.

Case 1. $r = 0$

In this case the sequence is $\{0, 0, 0, \dots\}$ which converges to 0.

Case 2. $r \neq 0$ and $|r| < 1$.

Since $|r| < 1$, $\frac{1}{|r|} > 1$. Let $\frac{1}{|r|} = 1 + a$ where $a > 0$. Then

$$|r^n - 0| = |r|^n = \frac{1}{(1 + a)^n}.$$

We have $(1 + a)^n > na$ for all $n \in \mathbb{N}$ and hence,

$$|r^n - 0| < \frac{1}{na} \text{ for all } n \in \mathbb{N}.$$

Let $\epsilon > 0$ be given. Then

$$|r^n - 0| < \epsilon \text{ holds if } n > \frac{1}{a\epsilon}.$$

Choose any $N \in \mathbb{N}$ such that $N > \frac{1}{a\epsilon}$. Then

$$\forall n \geq N, |r^n - 0| < \epsilon.$$

Since $\epsilon > 0$ is arbitrary,

$$\lim_{n \rightarrow \infty} r^n = 0.$$