

Directional Derivatives & Gradient Vectors

Devika S

Department of Mathematics
BITS Pilani, K K Birla Goa Campus

October 21, 2024



ANNOUNCEMENT:

An additional class will be held this **Saturday (26 October 2024)** from **12:00 PM to 1:00 PM** in **LT3**.

Recall - Directional derivative

Definition

The derivative of $f(x, y)$ at $P_0(x_0, y_0)$ in the direction of the unit vector $\mathbf{u} = u_1\mathbf{i} + u_2\mathbf{j}$ is the number

$$(D_{\mathbf{u}}f)_{P_0} = \left(\frac{df}{ds} \right)_{\mathbf{u}, P_0} := \lim_{s \rightarrow 0} \frac{f(x_0 + u_1s, y_0 + u_2s) - f(x_0, y_0)}{s}$$

provided the limit exists.

Recall - Directional derivative

Definition

The derivative of $f(x, y)$ at $P_0(x_0, y_0)$ in the direction of the unit vector $\mathbf{u} = u_1\mathbf{i} + u_2\mathbf{j}$ is the number

$$(D_{\mathbf{u}}f)_{P_0} = \left(\frac{df}{ds} \right)_{\mathbf{u}, P_0} := \lim_{s \rightarrow 0} \frac{f(x_0 + u_1s, y_0 + u_2s) - f(x_0, y_0)}{s}$$

provided the limit exists.

- $f_x(x_0, y_0)$ - directional derivative of f at P_0 in the direction of \mathbf{i}
- $f_y(x_0, y_0)$ - directional derivative of f at P_0 in the direction of \mathbf{j}
- directional derivative $(D_{\mathbf{u}}f)_{P_0}$ - rate of change of f at P_0 in the direction of \mathbf{u}

Recall - Directional derivative

Definition

The derivative of $f(x, y)$ at $P_0(x_0, y_0)$ in the direction of the unit vector $\mathbf{u} = u_1\mathbf{i} + u_2\mathbf{j}$ is the number

$$(D_{\mathbf{u}}f)_{P_0} = \left(\frac{df}{ds} \right)_{\mathbf{u}, P_0} := \lim_{s \rightarrow 0} \frac{f(x_0 + u_1s, y_0 + u_2s) - f(x_0, y_0)}{s}$$

provided the limit exists.

- $f_x(x_0, y_0)$ - directional derivative of f at P_0 in the direction of \mathbf{i}
- $f_y(x_0, y_0)$ - directional derivative of f at P_0 in the direction of \mathbf{j}
- directional derivative $(D_{\mathbf{u}}f)_{P_0}$ - rate of change of f at P_0 in the direction of \mathbf{u}
- For an angle θ measured from the positive x -axis,
 $\mathbf{u} = \cos \theta \mathbf{i} + \sin \theta \mathbf{j}$.

Formula to Calculate the Directional Derivative

Let $f(x, y)$ be a differentiable function and $g(s) := f(x_0 + u_1s, y_0 + u_2s)$.

Formula to Calculate the Directional Derivative

Let $f(x, y)$ be a differentiable function and $g(s) := f(x_0 + u_1s, y_0 + u_2s)$. Then,

$$g'(0) := \lim_{s \rightarrow 0} \frac{g(s) - g(0)}{s} = \lim_{s \rightarrow 0} \frac{f(x_0 + u_1s, y_0 + u_2s) - f(x_0, y_0)}{s}$$

Formula to Calculate the Directional Derivative

Let $f(x, y)$ be a differentiable function and $g(s) := f(x_0 + u_1s, y_0 + u_2s)$. Then,

$$g'(0) := \lim_{s \rightarrow 0} \frac{g(s) - g(0)}{s} = \lim_{s \rightarrow 0} \frac{f(x_0 + u_1s, y_0 + u_2s) - f(x_0, y_0)}{s} = (D_{\mathbf{u}}f)_{P_0}.$$

Formula to Calculate the Directional Derivative

Let $f(x, y)$ be a differentiable function and $g(s) := f(x_0 + u_1s, y_0 + u_2s)$. Then,

$$g'(0) := \lim_{s \rightarrow 0} \frac{g(s) - g(0)}{s} = \lim_{s \rightarrow 0} \frac{f(x_0 + u_1s, y_0 + u_2s) - f(x_0, y_0)}{s} = (D_{\mathbf{u}}f)_{P_0}.$$

Also, $g = f(x, y)$, where $x = x_0 + u_1s = x(s)$ and $y = y_0 + u_2s = y(s)$.

Formula to Calculate the Directional Derivative

Let $f(x, y)$ be a differentiable function and $g(s) := f(x_0 + u_1s, y_0 + u_2s)$. Then,

$$g'(0) := \lim_{s \rightarrow 0} \frac{g(s) - g(0)}{s} = \lim_{s \rightarrow 0} \frac{f(x_0 + u_1s, y_0 + u_2s) - f(x_0, y_0)}{s} = (D_{\mathbf{u}}f)_{P_0}.$$

Also, $g = f(x, y)$, where $x = x_0 + u_1s = x(s)$ and $y = y_0 + u_2s = y(s)$.

By chain rule,

$$g'(s) = \frac{dg}{ds} = \frac{\partial f}{\partial x} \frac{dx}{ds} + \frac{\partial f}{\partial y} \frac{dy}{ds} = f_x(x, y)u_1 + f_y(x, y)u_2.$$

Formula to Calculate the Directional Derivative

Let $f(x, y)$ be a differentiable function and $g(s) := f(x_0 + u_1s, y_0 + u_2s)$. Then,

$$g'(0) := \lim_{s \rightarrow 0} \frac{g(s) - g(0)}{s} = \lim_{s \rightarrow 0} \frac{f(x_0 + u_1s, y_0 + u_2s) - f(x_0, y_0)}{s} = (D_{\mathbf{u}}f)_{P_0}.$$

Also, $g = f(x, y)$, where $x = x_0 + u_1s = x(s)$ and $y = y_0 + u_2s = y(s)$.

By chain rule,

$$g'(s) = \frac{dg}{ds} = \frac{\partial f}{\partial x} \frac{dx}{ds} + \frac{\partial f}{\partial y} \frac{dy}{ds} = f_x(x, y)u_1 + f_y(x, y)u_2.$$

Now, choose $s = 0$. Then $x = x_0$, $y = y_0$ and

$$g'(0) = f_x(x_0, y_0)u_1 + f_y(x_0, y_0)u_2 = (f_x|_{P_0}, f_y|_{P_0}) \cdot (u_1, u_2).$$

Formula to Calculate the Directional Derivative

Let $f(x, y)$ be a differentiable function and $g(s) := f(x_0 + u_1s, y_0 + u_2s)$. Then,

$$g'(0) := \lim_{s \rightarrow 0} \frac{g(s) - g(0)}{s} = \lim_{s \rightarrow 0} \frac{f(x_0 + u_1s, y_0 + u_2s) - f(x_0, y_0)}{s} = (D_{\mathbf{u}}f)_{P_0}.$$

Also, $g = f(x, y)$, where $x = x_0 + u_1s = x(s)$ and $y = y_0 + u_2s = y(s)$.
By chain rule,

$$g'(s) = \frac{dg}{ds} = \frac{\partial f}{\partial x} \frac{dx}{ds} + \frac{\partial f}{\partial y} \frac{dy}{ds} = f_x(x, y)u_1 + f_y(x, y)u_2.$$

Now, choose $s = 0$. Then $x = x_0$, $y = y_0$ and

$$g'(0) = f_x(x_0, y_0)u_1 + f_y(x_0, y_0)u_2 = (f_x|_{P_0}, f_y|_{P_0}) \cdot (u_1, u_2).$$

Consequently, $(D_{\mathbf{u}}f)_{P_0} = (f_x|_{P_0}, f_y|_{P_0}) \cdot (u_1, u_2)$.

This says that the derivative of a differentiable function f in the direction of \mathbf{u} at P_0 is the dot product of \mathbf{u} with a special vector $(f_x|_{P_0}, f_y|_{P_0})$.

Definition

The **gradient vector** (or **gradient**) of $f(x, y)$ is the vector

$$\nabla f = \frac{\partial f}{\partial x} \mathbf{i} + \frac{\partial f}{\partial y} \mathbf{j} = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right).$$

Definition

The **gradient vector** (or **gradient**) of $f(x, y)$ is the vector

$$\nabla f = \frac{\partial f}{\partial x} \mathbf{i} + \frac{\partial f}{\partial y} \mathbf{j} = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right).$$

The value of the gradient vector obtained by evaluating the partial derivatives at a point $P_0(x_0, y_0)$ is written $\nabla f|_{P_0}$ or $\nabla f(x_0, y_0)$.

Definition

The **gradient vector** (or **gradient**) of $f(x, y)$ is the vector

$$\nabla f = \frac{\partial f}{\partial x} \mathbf{i} + \frac{\partial f}{\partial y} \mathbf{j} = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right).$$

The value of the gradient vector obtained by evaluating the partial derivatives at a point $P_0(x_0, y_0)$ is written $\nabla f|_{P_0}$ or $\nabla f(x_0, y_0)$.

Theorem (The Directional Derivative is a Dot Product)

If $f(x, y)$ is *differentiable*, then

$$\left(\frac{df}{ds} \right)_{\mathbf{u}, P_0} = (\nabla f)_{P_0} \cdot \mathbf{u},$$

the dot product of ∇f at P_0 and \mathbf{u} . In brief, $D_{\mathbf{u}}f = \nabla f \cdot \mathbf{u}$.

The above theorem says that if a function f is differentiable at $P_0(x_0, y_0)$, then all its directional derivatives exist.

The above theorem says that if a function f is differentiable at $P_0(x_0, y_0)$, then all its directional derivatives exist. What about the converse? Existence of all directional derivatives of f does not imply f is differentiable.

The above theorem says that if a function f is differentiable at $P_0(x_0, y_0)$, then all its directional derivatives exist. What about the converse? Existence of all directional derivatives of f does not imply f is differentiable.

Consider

$$f(x, y) = \begin{cases} \frac{x^2 y}{x^4 + y^2} & \text{if } (x, y) \neq (0, 0), \\ 0 & \text{if } (x, y) = (0, 0). \end{cases}$$

The above theorem says that if a function f is differentiable at $P_0(x_0, y_0)$, then all its directional derivatives exist. What about the converse? Existence of all directional derivatives of f does not imply f is differentiable.

Consider

$$f(x, y) = \begin{cases} \frac{x^2 y}{x^4 + y^2} & \text{if } (x, y) \neq (0, 0), \\ 0 & \text{if } (x, y) = (0, 0). \end{cases}$$

This function is not continuous at $(0, 0)$ (choose $y = mx^2, m \neq 0$ and show that limit does not exist) and hence is not differentiable at $(0, 0)$.

The above theorem says that if a function f is differentiable at $P_0(x_0, y_0)$, then all its directional derivatives exist. **What about the converse?** Existence of all directional derivatives of f does not imply f is differentiable.

Consider

$$f(x, y) = \begin{cases} \frac{x^2 y}{x^4 + y^2} & \text{if } (x, y) \neq (0, 0), \\ 0 & \text{if } (x, y) = (0, 0). \end{cases}$$

This function is not continuous at $(0, 0)$ (**choose $y = mx^2, m \neq 0$ and show that limit does not exist**) and hence is not differentiable at $(0, 0)$.

Let $\mathbf{u} = (u_1, u_2)$ be a unit vector. Then,

$$\begin{aligned} (D_{\mathbf{u}}f)|_{(0,0)} &= \lim_{s \rightarrow 0} \frac{f(u_1 s, u_2 s) - f(0, 0)}{s} = \lim_{s \rightarrow 0} \frac{s^3 u_1^2 u_2}{s(s^4 u_1^4 + s^2 u_2^2)} \\ &= \lim_{s \rightarrow 0} \frac{u_1^2 u_2}{s^2 u_1^4 + u_2^2} = \begin{cases} \frac{u_1^2}{u_2} & \text{if } u_2 \neq 0, \\ 0 & \text{if } u_2 = 0. \end{cases} \end{aligned}$$

This shows that the directional derivatives in all directions at $(0, 0)$ exist.

Examples

- 1 Find the derivative of $f(x, y) = 2xy - 3y^2$ at $P_0(5, 5)$ in the direction of $\mathbf{u} = 4\mathbf{i} + 3\mathbf{j}$ using gradient.

Examples

- ① Find the derivative of $f(x, y) = 2xy - 3y^2$ at $P_0(5, 5)$ in the direction of $\mathbf{u} = 4\mathbf{i} + 3\mathbf{j}$ using gradient.

Solution:

Recall $\mathbf{v} = \frac{4}{5}\mathbf{i} + \frac{3}{5}\mathbf{j}$ is the unit vector in the direction of \mathbf{u} .

Examples

- ① Find the derivative of $f(x, y) = 2xy - 3y^2$ at $P_0(5, 5)$ in the direction of $\mathbf{u} = 4\mathbf{i} + 3\mathbf{j}$ using gradient.

Solution:

Recall $\mathbf{v} = \frac{4}{5}\mathbf{i} + \frac{3}{5}\mathbf{j}$ is the unit vector in the direction of \mathbf{u} .

To apply the above theorem, we need to check the differentiability of f at $(5, 5)$.

Examples

- ① Find the derivative of $f(x, y) = 2xy - 3y^2$ at $P_0(5, 5)$ in the direction of $\mathbf{u} = 4\mathbf{i} + 3\mathbf{j}$ using gradient.

Solution:

Recall $\mathbf{v} = \frac{4}{5}\mathbf{i} + \frac{3}{5}\mathbf{j}$ is the unit vector in the direction of \mathbf{u} .

To apply the above theorem, we need to check the differentiability of f at $(5, 5)$. We have $f_x = 2y$ and $f_y = 2x - 6y$. The partial derivatives f_x and f_y exists, and also f_x and f_y are everywhere continuous (**Why?**).

Examples

- ① Find the derivative of $f(x, y) = 2xy - 3y^2$ at $P_0(5, 5)$ in the direction of $\mathbf{u} = 4\mathbf{i} + 3\mathbf{j}$ using gradient.

Solution:

Recall $\mathbf{v} = \frac{4}{5}\mathbf{i} + \frac{3}{5}\mathbf{j}$ is the unit vector in the direction of \mathbf{u} .

To apply the above theorem, we need to check the differentiability of f at $(5, 5)$. We have $f_x = 2y$ and $f_y = 2x - 6y$. The partial derivatives f_x and f_y exists, and also f_x and f_y are everywhere continuous (**Why?**). Hence, f is differentiable at $(5, 5)$.

Examples

- ① Find the derivative of $f(x, y) = 2xy - 3y^2$ at $P_0(5, 5)$ in the direction of $\mathbf{u} = 4\mathbf{i} + 3\mathbf{j}$ using gradient.

Solution:

Recall $\mathbf{v} = \frac{4}{5}\mathbf{i} + \frac{3}{5}\mathbf{j}$ is the unit vector in the direction of \mathbf{u} .

To apply the above theorem, we need to check the differentiability of f at $(5, 5)$. We have $f_x = 2y$ and $f_y = 2x - 6y$. The partial derivatives f_x and f_y exists, and also f_x and f_y are everywhere continuous (**Why?**). Hence, f is differentiable at $(5, 5)$.

$$\begin{aligned}(D_{\mathbf{v}}f)_{P_0} &= \nabla f|_{P_0} \cdot \mathbf{v} = (f_x, f_y)|_{P_0} \cdot \mathbf{v} \\ &= (10, -20) \cdot \left(\frac{4}{5}, \frac{3}{5}\right) = 10 \left(\frac{4}{5}\right) - 20 \left(\frac{3}{5}\right) = -4.\end{aligned}$$

Examples (Using Gradient)

- ② Find the derivative of $f(x, y, z) = xy + yz + zx$ at $P_0(1, -1, 2)$ in the direction of $\mathbf{u} = 3\mathbf{i} + 6\mathbf{j} - 2\mathbf{k}$. (Ans: 3)
- ③ Find the derivative of $g(x, y) = \frac{x-y}{xy+2}$ at $P_0(1, -1)$ in the direction of $\mathbf{u} = 12\mathbf{i} + 5\mathbf{j}$. (Ans: -4)
- ④ Find the derivative of $h(x, y) = \tan^{-1}(y/x) + \sqrt{3}\sin^{-1}(xy/2)$ at $P_0(1, 1)$ in the direction of $\mathbf{u} = 3\mathbf{i} - 2\mathbf{j}$. (Ans: $3/(2\sqrt{13})$)
- ⑤ Find the derivative of $g(x, y, z) = 3e^x \cos(yz)$ at $P_0(0, 0, 0)$ in the direction of $\mathbf{u} = 2\mathbf{i} + \mathbf{j} - 2\mathbf{k}$. (Ans: 2)

Examples (Using Gradient)

- ② Find the derivative of $f(x, y, z) = xy + yz + zx$ at $P_0(1, -1, 2)$ in the direction of $\mathbf{u} = 3\mathbf{i} + 6\mathbf{j} - 2\mathbf{k}$. (Ans: 3)
- ③ Find the derivative of $g(x, y) = \frac{x-y}{xy+2}$ at $P_0(1, -1)$ in the direction of $\mathbf{u} = 12\mathbf{i} + 5\mathbf{j}$. (Ans: -4)
- ④ Find the derivative of $h(x, y) = \tan^{-1}(y/x) + \sqrt{3}\sin^{-1}(xy/2)$ at $P_0(1, 1)$ in the direction of $\mathbf{u} = 3\mathbf{i} - 2\mathbf{j}$. (Ans: $3/(2\sqrt{13})$)
- ⑤ Find the derivative of $g(x, y, z) = 3e^x \cos(yz)$ at $P_0(0, 0, 0)$ in the direction of $\mathbf{u} = 2\mathbf{i} + \mathbf{j} - 2\mathbf{k}$. (Ans: 2)

For two vectors \mathbf{v} and \mathbf{w} ,

$$\mathbf{v} \cdot \mathbf{w} = |\mathbf{v}||\mathbf{w}|\cos\theta,$$

where $0 \leq \theta \leq \pi$ is the angle between \mathbf{v} and \mathbf{w} .

Properties of $D_{\mathbf{u}}f = \nabla f \cdot \mathbf{u} = |\nabla f| \cos \theta$

- 1 The function f increases most rapidly when $\cos \theta = 1$ or when $\theta = 0$ and \mathbf{u} is the direction of ∇f . That is, at each point P in its domain, f increases most rapidly in the direction of the gradient vector ∇f at P . The derivative in this direction is $D_{\mathbf{u}}f = |\nabla f| \cos 0 = |\nabla f|$.

Properties of $D_{\mathbf{u}}f = \nabla f \cdot \mathbf{u} = |\nabla f| \cos \theta$

- 1 The function f increases most rapidly when $\cos \theta = 1$ or when $\theta = 0$ and \mathbf{u} is the direction of ∇f . That is, at each point P in its domain, f increases most rapidly in the direction of the gradient vector ∇f at P . The derivative in this direction is $D_{\mathbf{u}}f = |\nabla f| \cos 0 = |\nabla f|$.
- 2 f decreases most rapidly in the direction of $-\nabla f$. The derivative in this direction is $D_{\mathbf{u}}f = |\nabla f| \cos \pi = -|\nabla f|$.

Properties of $D_{\mathbf{u}}f = \nabla f \cdot \mathbf{u} = |\nabla f| \cos \theta$

- 1 The function f increases most rapidly when $\cos \theta = 1$ or when $\theta = 0$ and \mathbf{u} is the direction of ∇f . That is, at each point P in its domain, f increases most rapidly in the direction of the gradient vector ∇f at P . The derivative in this direction is $D_{\mathbf{u}}f = |\nabla f| \cos 0 = |\nabla f|$.
- 2 f decreases most rapidly in the direction of $-\nabla f$. The derivative in this direction is $D_{\mathbf{u}}f = |\nabla f| \cos \pi = -|\nabla f|$.
- 3 Any direction \mathbf{u} orthogonal to a gradient $\nabla f \neq 0$ is a direction of zero change in f because θ then equals to $\pi/2$ and $D_{\mathbf{u}}f = |\nabla f| \cos(\pi/2) = 0$.

Properties of $D_{\mathbf{u}}f = \nabla f \cdot \mathbf{u} = |\nabla f| \cos \theta$


- 1 The function f increases most rapidly when $\cos \theta = 1$ or when $\theta = 0$ and \mathbf{u} is the direction of ∇f . That is, at each point P in its domain, f increases most rapidly in the direction of the gradient vector ∇f at P . The derivative in this direction is $D_{\mathbf{u}}f = |\nabla f| \cos 0 = |\nabla f|$.
- 2 f decreases most rapidly in the direction of $-\nabla f$. The derivative in this direction is $D_{\mathbf{u}}f = |\nabla f| \cos \pi = -|\nabla f|$.
- 3 Any direction \mathbf{u} orthogonal to a gradient $\nabla f \neq 0$ is a direction of zero change in f because θ then equals to $\pi/2$ and $D_{\mathbf{u}}f = |\nabla f| \cos(\pi/2) = 0$.

Hence,

$$D_{\mathbf{u}}f \in [-|\nabla f|, |\nabla f|].$$

$D_{\mathbf{u}}f$ is maximum when $\theta = 0$ and minimum when $\theta = \pi$.

Figure for $f(x, y) = \frac{x^2}{2} + \frac{y^2}{2}$



Examples

- 1 Find the directions in which $f(x, y) = x^2y + e^{xy} \sin y$ increase and decrease most rapidly at $P_0(1, 0)$. Then find the derivatives of f in these directions.

Examples

- ① Find the directions in which $f(x, y) = x^2y + e^{xy} \sin y$ increase and decrease most rapidly at $P_0(1, 0)$. Then find the derivatives of f in these directions.

Solution:

The function f increases most rapidly in the direction of ∇f at $(1, 0)$.
The gradient at $(1, 0)$ is

$$\nabla f|_{(1,0)} = (2xy + ye^{xy} \sin y)\mathbf{i} + (x^2 + xe^{xy} \sin y + e^{xy} \cos y)\mathbf{j}|_{(1,0)} = 2\mathbf{j}.$$

Examples

- ① Find the directions in which $f(x, y) = x^2y + e^{xy} \sin y$ increase and decrease most rapidly at $P_0(1, 0)$. Then find the derivatives of f in these directions.

Solution:

The function f increases most rapidly in the direction of ∇f at $(1, 0)$. The gradient at $(1, 0)$ is

$$\nabla f|_{(1,0)} = (2xy + ye^{xy} \sin y)\mathbf{i} + (x^2 + xe^{xy} \sin y + e^{xy} \cos y)\mathbf{j}|_{(1,0)} = 2\mathbf{j}.$$

Its direction is

$$\mathbf{u} = \frac{2\mathbf{j}}{|2\mathbf{j}|} = \mathbf{j}.$$

f increases most rapidly in the direction $\mathbf{u} = \mathbf{j}$. The derivative of f in this direction is $D_{\mathbf{u}}(f) = |\nabla f| = 2$.

Examples

- ① Find the directions in which $f(x, y) = x^2y + e^{xy} \sin y$ increase and decrease most rapidly at $P_0(1, 0)$. Then find the derivatives of f in these directions.

Solution:

The function f increases most rapidly in the direction of ∇f at $(1, 0)$. The gradient at $(1, 0)$ is

$$\nabla f|_{(1,0)} = (2xy + ye^{xy} \sin y)\mathbf{i} + (x^2 + xe^{xy} \sin y + e^{xy} \cos y)\mathbf{j}|_{(1,0)} = 2\mathbf{j}.$$

Its direction is

$$\mathbf{u} = \frac{2\mathbf{j}}{|2\mathbf{j}|} = \mathbf{j}.$$

f increases most rapidly in the direction $\mathbf{u} = \mathbf{j}$. The derivative of f in this direction is $D_{\mathbf{u}}(f) = |\nabla f| = 2$.

f decreases most rapidly in the direction $-\mathbf{u} = -\mathbf{j}$ and the derivative of f in this direction is $D_{-\mathbf{u}}(f) = -2$.

Examples

- ② Find the directions in which $g(x, y, z) = xe^y + z^2$, increase and decrease most rapidly at $P_0(1, \ln 2, 1/2)$. Then find the derivatives of g in these directions. (Ans: 3, -3)

Examples

- ② Find the directions in which $g(x, y, z) = xe^y + z^2$, increase and decrease most rapidly at $P_0(1, \ln 2, 1/2)$. Then find the derivatives of g in these directions. (Ans: $3, -3$)
- ③ In what direction is the derivative of $f(x, y) = xy + y^2$ at $P(3, 2)$ is zero? (Ans: $\frac{7}{\sqrt{53}}\mathbf{i} - \frac{2}{\sqrt{53}}\mathbf{j}, \frac{-7}{\sqrt{53}}\mathbf{i} + \frac{2}{\sqrt{53}}\mathbf{j}$)

Examples

- 2 Find the directions in which $g(x, y, z) = xe^y + z^2$, increase and decrease most rapidly at $P_0(1, \ln 2, 1/2)$. Then find the derivatives of g in these directions. (Ans: 3, -3)
- 3 In what direction is the derivative of $f(x, y) = xy + y^2$ at $P(3, 2)$ is zero? (Ans: $\frac{7}{\sqrt{53}}\mathbf{i} - \frac{2}{\sqrt{53}}\mathbf{j}$, $\frac{-7}{\sqrt{53}}\mathbf{i} + \frac{2}{\sqrt{53}}\mathbf{j}$)
- 4 Is there a direction \mathbf{u} in which the rate of change of the temperature function $T(x, y, z) = 2xy - yz$ (temperature in degrees Celsius, distance in feet) at $P(1, -1, 1)$ is $-3^\circ\text{C}/\text{ft}$? Give reasons for your answer.

Examples

- ② Find the directions in which $g(x, y, z) = xe^y + z^2$, increase and decrease most rapidly at $P_0(1, \ln 2, 1/2)$. Then find the derivatives of g in these directions. (Ans: 3, -3)
- ③ In what direction is the derivative of $f(x, y) = xy + y^2$ at $P(3, 2)$ is zero? (Ans: $\frac{7}{\sqrt{53}}\mathbf{i} - \frac{2}{\sqrt{53}}\mathbf{j}$, $\frac{-7}{\sqrt{53}}\mathbf{i} + \frac{2}{\sqrt{53}}\mathbf{j}$)
- ④ Is there a direction \mathbf{u} in which the rate of change of the temperature function $T(x, y, z) = 2xy - yz$ (temperature in degrees Celsius, distance in feet) at $P(1, -1, 1)$ is $-3^\circ\text{C}/\text{ft}$? Give reasons for your answer. (No - Compute max. and min. of $D_{\mathbf{u}}f$.)
- ⑤ The derivative of $f(x, y)$ at $P_0(1, 2)$ in the direction of $\mathbf{i} + \mathbf{j}$ is $2\sqrt{2}$ and in the direction of $-2\mathbf{j}$ is -3 . What is the derivative of f in the direction of $-\mathbf{i} - 2\mathbf{j}$?

Examples

- ② Find the directions in which $g(x, y, z) = xe^y + z^2$, increase and decrease most rapidly at $P_0(1, \ln 2, 1/2)$. Then find the derivatives of g in these directions. (Ans: $3, -3$)
- ③ In what direction is the derivative of $f(x, y) = xy + y^2$ at $P(3, 2)$ is zero? (Ans: $\frac{7}{\sqrt{53}}\mathbf{i} - \frac{2}{\sqrt{53}}\mathbf{j}, \frac{-7}{\sqrt{53}}\mathbf{i} + \frac{2}{\sqrt{53}}\mathbf{j}$)
- ④ Is there a direction \mathbf{u} in which the rate of change of the temperature function $T(x, y, z) = 2xy - yz$ (temperature in degrees Celsius, distance in feet) at $P(1, -1, 1)$ is $-3^\circ\text{C}/\text{ft}$? Give reasons for your answer. (No - Compute max. and min. of $D_{\mathbf{u}}f$.)
- ⑤ The derivative of $f(x, y)$ at $P_0(1, 2)$ in the direction of $\mathbf{i} + \mathbf{j}$ is $2\sqrt{2}$ and in the direction of $-2\mathbf{j}$ is -3 . What is the derivative of f in the direction of $-\mathbf{i} - 2\mathbf{j}$? (Ans: $\frac{-7}{\sqrt{5}}$)

Algebra Rules for Gradients

- **Sum Rule:** $\nabla(f + g) = \nabla f + \nabla g$.
- **Difference Rule:** $\nabla(f - g) = \nabla f - \nabla g$.
- **Constant Multiple Rule:** $\nabla(kf) = k\nabla(f)$ for any constant k
- **Product Rule:** $\nabla(fg) = f\nabla(g) + g\nabla(f)$.
- **Quotient Rule:** $\nabla\left(\frac{f}{g}\right) = \frac{g\nabla f - f\nabla g}{g^2}$

Functions of Three Variables

For functions of three variables we can define directional derivatives in a similar manner.

Functions of Three Variables

For functions of three variables we can define directional derivatives in a similar manner.

The derivative of $f(x, y, z)$ at $P_0(x_0, y_0, z_0)$ in the direction of the unit vector $\mathbf{u} = u_1\mathbf{i} + u_2\mathbf{j} + u_3\mathbf{k}$ is the number

$$(D_{\mathbf{u}}f)_{P_0} = \lim_{s \rightarrow 0} \frac{f(x_0 + u_1s, y_0 + u_2s, z_0 + u_3s) - f(x_0, y_0, z_0)}{s}$$

provided the limit exists.

Functions of Three Variables

For functions of three variables we can define directional derivatives in a similar manner.

The derivative of $f(x, y, z)$ at $P_0(x_0, y_0, z_0)$ in the direction of the unit vector $\mathbf{u} = u_1\mathbf{i} + u_2\mathbf{j} + u_3\mathbf{k}$ is the number

$$(D_{\mathbf{u}}f)_{P_0} = \lim_{s \rightarrow 0} \frac{f(x_0 + u_1s, y_0 + u_2s, z_0 + u_3s) - f(x_0, y_0, z_0)}{s}$$

provided the limit exists. If f is differentiable, then

$$(D_{\mathbf{u}}f)_{P_0} = \nabla f \cdot \mathbf{u}, \text{ where } \nabla f = (f_x, f_y, f_z).$$

Functions of Three Variables

For functions of three variables we can define directional derivatives in a similar manner.

The derivative of $f(x, y, z)$ at $P_0(x_0, y_0, z_0)$ in the direction of the unit vector $\mathbf{u} = u_1\mathbf{i} + u_2\mathbf{j} + u_3\mathbf{k}$ is the number

$$(D_{\mathbf{u}}f)_{P_0} = \lim_{s \rightarrow 0} \frac{f(x_0 + u_1s, y_0 + u_2s, z_0 + u_3s) - f(x_0, y_0, z_0)}{s}$$

provided the limit exists. If f is differentiable, then

$$(D_{\mathbf{u}}f)_{P_0} = \nabla f \cdot \mathbf{u}, \text{ where } \nabla f = (f_x, f_y, f_z).$$

At any given point, f increases most rapidly in the direction of ∇f and decreases most rapidly in the direction of $-\nabla f$. In any direction orthogonal to ∇f , the derivative is zero.

Gradients and Tangents to Level Curves

If a differentiable function $f(x, y)$ has a constant value c along a smooth curve $\mathbf{r} = g(t)\mathbf{i} + h(t)\mathbf{j}$, then $f(g(t), h(t)) = c$.

Differentiating both sides of this equation with respect to t leads to

$$\frac{d}{dt}f(g(t), h(t)) = \frac{d}{dt}(c)$$

Using chain rule, $\frac{\partial f}{\partial x} \frac{dg}{dt} + \frac{\partial f}{\partial y} \frac{dh}{dt} = 0 \implies \nabla f \cdot \frac{d\mathbf{r}}{dt} = 0.$

Gradients and Tangents to Level Curves

If a differentiable function $f(x, y)$ has a constant value c along a smooth curve $\mathbf{r} = g(t)\mathbf{i} + h(t)\mathbf{j}$, then $f(g(t), h(t)) = c$.

Differentiating both sides of this equation with respect to t leads to

$$\frac{d}{dt}f(g(t), h(t)) = \frac{d}{dt}(c)$$

$$\text{Using chain rule, } \frac{\partial f}{\partial x} \frac{dg}{dt} + \frac{\partial f}{\partial y} \frac{dh}{dt} = 0 \implies \nabla f \cdot \frac{d\mathbf{r}}{dt} = 0.$$

This says that ∇f is orthogonal to the tangent vector $d\mathbf{r}/dt$, so it is normal to the curve.

At every point (x_0, y_0) in the domain of a differentiable function $f(x, y)$, the gradient of f is normal to the level curve through (x_0, y_0) .

Gradients to Level Curves




FIGURE 14.30 The gradient of a differentiable function of two variables at a point is always normal to the function's level curve through that point.

Tangent Lines

Tangent lines - lines that are tangent to the level curves

\implies tangent lines are the lines normal to the gradients. Why?

Tangent Lines

Tangent lines - lines that are tangent to the level curves

\implies tangent lines are the lines normal to the gradients. Why?

The line passing through a point $P_0(x_0, y_0)$ normal to the vector $\mathbf{N} = A\mathbf{i} + B\mathbf{j}$ has the equation

$$A(x - x_0) + B(y - y_0) = 0.$$

Tangent Lines

Tangent lines - lines that are tangent to the level curves

\implies tangent lines are the lines normal to the gradients. **Why?**

The line passing through a point $P_0(x_0, y_0)$ normal to the vector $\mathbf{N} = A\mathbf{i} + B\mathbf{j}$ has the equation

$$A(x - x_0) + B(y - y_0) = 0.$$

If $\mathbf{N} = \nabla f$, then the equation of the tangent line to level curve $f(x, y) = c$ at $P_0(x_0, y_0)$ is given by

$$f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0) = 0.$$

Tangent Lines

Tangent lines - lines that are tangent to the level curves

\implies tangent lines are the lines normal to the gradients. **Why?**

The line passing through a point $P_0(x_0, y_0)$ normal to the vector $\mathbf{N} = A\mathbf{i} + B\mathbf{j}$ has the equation

$$A(x - x_0) + B(y - y_0) = 0.$$

If $\mathbf{N} = \nabla f$, then the equation of the tangent line to level curve $f(x, y) = c$ at $P_0(x_0, y_0)$ is given by

$$f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0) = 0.$$

- Write an equation for the tangent line $x^2 - xy + y^2 = 7$, $P_0(-1, 2)$.

Tangent Lines

Tangent lines - lines that are tangent to the level curves

\implies tangent lines are the lines normal to the gradients. **Why?**

The line passing through a point $P_0(x_0, y_0)$ normal to the vector $\mathbf{N} = A\mathbf{i} + B\mathbf{j}$ has the equation

$$A(x - x_0) + B(y - y_0) = 0.$$

If $\mathbf{N} = \nabla f$, then the equation of the tangent line to level curve $f(x, y) = c$ at $P_0(x_0, y_0)$ is given by

$$f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0) = 0.$$

- Write an equation for the tangent line $x^2 - xy + y^2 = 7$, $P_0(-1, 2)$. (**Ans:** $y = x - 4$)

Tangent Lines

Tangent lines - lines that are tangent to the level curves

\implies tangent lines are the lines normal to the gradients. **Why?**

The line passing through a point $P_0(x_0, y_0)$ normal to the vector $\mathbf{N} = A\mathbf{i} + B\mathbf{j}$ has the equation

$$A(x - x_0) + B(y - y_0) = 0.$$

If $\mathbf{N} = \nabla f$, then the equation of the tangent line to level curve $f(x, y) = c$ at $P_0(x_0, y_0)$ is given by

$$f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0) = 0.$$

- Write an equation for the tangent line $x^2 - xy + y^2 = 7$, $P_0(-1, 2)$. (**Ans:** $y = x - 4$)
- Write an equation for the tangent line $xy = -4$, $P_0(2, -2)$. (**Ans:** $-4x + 5y - 14 = 0$.)