

Lagrange Multipliers

Devika S

Department of Mathematics
BITS Pilani, K K Birla Goa Campus

October 28, 2024



Lagrange Multipliers

Consider the following problems:

- Find the maximum value of $s(x, y, z) = xy + yz + xz$ where $x + y + z = 6$.

Lagrange Multipliers

Consider the following problems:

- Find the maximum value of $s(x, y, z) = xy + yz + xz$ where $x + y + z = 6$.
- Find the points on the hyperbolic cylinder $x^2 - z^2 - 1 = 0$ that are closest to the origin.

Lagrange Multipliers

Consider the following problems:

- Find the maximum value of $s(x, y, z) = xy + yz + xz$ where $x + y + z = 6$.
- Find the points on the hyperbolic cylinder $x^2 - z^2 - 1 = 0$ that are closest to the origin.
- Find the extreme values of the function $f(x, y, z) = x^2yz + 1$ on the intersection of $z = 1$ and $x^2 + y^2 + z^2 = 10$.

Lagrange Multipliers

Consider the following problems:

- Find the maximum value of $s(x, y, z) = xy + yz + xz$ where $x + y + z = 6$.
- Find the points on the hyperbolic cylinder $x^2 - z^2 - 1 = 0$ that are closest to the origin.
- Find the extreme values of the function $f(x, y, z) = x^2yz + 1$ on the intersection of $z = 1$ and $x^2 + y^2 + z^2 = 10$.

To deal with the situation when we have to find maximum or minimum of a function $f(x, y, z)$ subject to a constraint $g(x, y, z) = 0$ (or more than one constraints, say, $g_1(x, y, z) = 0$ and $g_2(x, y, z) = 0$), we use **Lagrange multiplier method**.

Lagrange Multipliers

Consider the following problems:

- Find the maximum value of $s(x, y, z) = xy + yz + xz$ where $x + y + z = 6$.
- Find the points on the hyperbolic cylinder $x^2 - z^2 - 1 = 0$ that are closest to the origin.
- Find the extreme values of the function $f(x, y, z) = x^2yz + 1$ on the intersection of $z = 1$ and $x^2 + y^2 + z^2 = 10$.

To deal with the situation when we have to find maximum or minimum of a function $f(x, y, z)$ subject to a constraint $g(x, y, z) = 0$ (or more than one constraints, say, $g_1(x, y, z) = 0$ and $g_2(x, y, z) = 0$), we use **Lagrange multiplier method**.

Theorem (The Orthogonal Gradient Theorem)

Suppose that $f(x, y, z)$ is differentiable in a region whose interior contains a smooth curve $C: \mathbf{r}(t) = g(t)\mathbf{i} + h(t)\mathbf{j} + k(t)\mathbf{k}$.

If P_0 is a point on C where f has a local maximum or minimum relative to its values on C , then ∇f is orthogonal to C at P_0 .

Observation

Theorem (The Orthogonal Gradient Theorem)

Suppose that $f(x, y, z)$ is differentiable in a region whose interior contains a smooth curve $C: \mathbf{r}(t) = g(t)\mathbf{i} + h(t)\mathbf{j} + k(t)\mathbf{k}$.

If P_0 is a point on C where f has a local maximum or minimum relative to its values on C , then ∇f is orthogonal to C at P_0 .

Proof: We want to show that $\nabla f \cdot \mathbf{v} = 0$ where $\mathbf{v} = \frac{d\mathbf{r}}{dt}$.

The f on C is given by $f(g(t), h(t), k(t))$. Hence,

$$\frac{df}{dt} = \frac{\partial f}{\partial x} \frac{dg}{dt} + \frac{\partial f}{\partial y} \frac{dh}{dt} + \frac{\partial f}{\partial z} \frac{dk}{dt} = \nabla f \cdot \mathbf{v}.$$

Theorem (The Orthogonal Gradient Theorem)

Suppose that $f(x, y, z)$ is differentiable in a region whose interior contains a smooth curve $C: \mathbf{r}(t) = g(t)\mathbf{i} + h(t)\mathbf{j} + k(t)\mathbf{k}$.

If P_0 is a point on C where f has a local maximum or minimum relative to its values on C , then ∇f is orthogonal to C at P_0 .

Proof: We want to show that $\nabla f \cdot \mathbf{v} = 0$ where $\mathbf{v} = \frac{d\mathbf{r}}{dt}$.

The f on C is given by $f(g(t), h(t), k(t))$. Hence,

$$\frac{df}{dt} = \frac{\partial f}{\partial x} \frac{dg}{dt} + \frac{\partial f}{\partial y} \frac{dh}{dt} + \frac{\partial f}{\partial z} \frac{dk}{dt} = \nabla f \cdot \mathbf{v}.$$

At any point P_0 , where f has a local maxima or minima relative to its values on the curve, $\frac{df}{dt} = 0$. This completes the proof.

Corollary (Side remark: Two variable case)

At the points on a smooth curve $\mathbf{r}(t) = g(t)\mathbf{i} + h(t)\mathbf{j}$ where a differentiable function $f(x, y)$ takes on its local maxima and minima relative to its values on the curve, $\nabla f \cdot \mathbf{v} = 0$, where $\mathbf{v} = d\mathbf{r}/dt$.

Corollary (Side remark: Two variable case)

At the points on a smooth curve $\mathbf{r}(t) = g(t)\mathbf{i} + h(t)\mathbf{j}$ where a differentiable function $f(x, y)$ takes on its local maxima and minima relative to its values on the curve, $\nabla f \cdot \mathbf{v} = 0$, where $\mathbf{v} = d\mathbf{r}/dt$.

Theorem (The Method of Lagrange Multipliers)

Suppose that $f(x, y, z)$ and $g(x, y, z)$ are differentiable and $\nabla g \neq 0$ when $g(x, y, z) = 0$. To find the local maximum and minimum values of f subject to the constraint $g(x, y, z) = 0$ (if these exist), find the values of x, y, z , and λ that simultaneously satisfy the equations

$$\nabla f = \lambda \nabla g \quad \text{and} \quad g(x, y, z) = 0.$$

For functions of two independent variables, the condition is similar, but without the variable z .

Here, the scalar λ is called a **Lagrange multiplier**.

- The geometry of the orthogonal gradient theorem tells us that at the point of the surface $g = 0$ where f assumes maximum or minimum value, the gradient of f is orthogonal to the velocity vector of any curve on the surface $g = 0$ at that point.

- The geometry of the orthogonal gradient theorem tells us that at the point of the surface $g = 0$ where f assumes maximum or minimum value, the gradient of f is orthogonal to the velocity vector of any curve on the surface $g = 0$ at that point.
- And the velocity vector of any curve at the point is orthogonal to the gradient of the defining equation of the surface g .

- The geometry of the orthogonal gradient theorem tells us that at the point of the surface $g = 0$ where f assumes maximum or minimum value, the gradient of f is orthogonal to the velocity vector of any curve on the surface $g = 0$ at that point.
- And the velocity vector of any curve at the point is orthogonal to the gradient of the defining equation of the surface g .
- Putting these two together we see that at the point of the surface where the function f take on extreme values, the gradient of the function f must be a scalar multiple of the gradient of the defining equation of the surface g .

Examples

Example 1. Find the dimensions of the closed right circular cylindrical can of smallest surface area whose volume is $16\pi \text{ cm}^3$.

Examples

Example 1. Find the dimensions of the closed right circular cylindrical can of smallest surface area whose volume is $16\pi \text{ cm}^3$.

Solution: Let us assume the radius of the can is r and the height is h .

Examples

Example 1. Find the dimensions of the closed right circular cylindrical can of smallest surface area whose volume is $16\pi \text{ cm}^3$.

Solution: Let us assume the radius of the can is r and the height is h .

We want to minimize the surface area which is $f(r, h) = 2\pi rh + 2\pi r^2$ given that the volume is $\pi r^2 h = 16\pi$.

Examples

Example 1. Find the dimensions of the closed right circular cylindrical can of smallest surface area whose volume is $16\pi \text{ cm}^3$.

Solution: Let us assume the radius of the can is r and the height is h .

We want to minimize the surface area which is $f(r, h) = 2\pi rh + 2\pi r^2$ given that the volume is $\pi r^2 h = 16\pi$.

The problem is essentially to minimize the function $f(r, h)$ on the surface $g(r, h) = r^2 h - 16 = 0$.

- We set up the Lagrange multiplier method first.

$$\nabla f(r, h) = \lambda \nabla g(r, h), \quad g(r, h) = 0.$$

Examples

- Since

$$\nabla f(r, h) = (2\pi h + 4\pi r)\mathbf{i} + 2\pi r\mathbf{j}, \quad \nabla g(r, h) = 2rh\mathbf{i} + r^2\mathbf{j}.$$

Examples

- Since

$$\nabla f(r, h) = (2\pi h + 4\pi r)\mathbf{i} + 2\pi r\mathbf{j}, \quad \nabla g(r, h) = 2rh\mathbf{i} + r^2\mathbf{j}.$$

So $\nabla f(r, h) = \lambda \nabla g(r, h)$ becomes

$$(2\pi h + 4\pi r)\mathbf{i} + 2\pi r\mathbf{j} = 2rh\lambda\mathbf{i} + r^2\lambda\mathbf{j}$$

Examples

- Since

$$\nabla f(r, h) = (2\pi h + 4\pi r)\mathbf{i} + 2\pi r\mathbf{j}, \quad \nabla g(r, h) = 2rh\mathbf{i} + r^2\mathbf{j}.$$

So $\nabla f(r, h) = \lambda \nabla g(r, h)$ becomes

$$(2\pi h + 4\pi r)\mathbf{i} + 2\pi r\mathbf{j} = 2rh\lambda\mathbf{i} + r^2\lambda\mathbf{j}$$

- Equating, $2\pi h + 4\pi r = 2\pi rh\lambda$ and $2\pi r = r^2\lambda$.

Examples

- Since

$$\nabla f(r, h) = (2\pi h + 4\pi r)\mathbf{i} + 2\pi r\mathbf{j}, \quad \nabla g(r, h) = 2rh\mathbf{i} + r^2\mathbf{j}.$$

So $\nabla f(r, h) = \lambda \nabla g(r, h)$ becomes

$$(2\pi h + 4\pi r)\mathbf{i} + 2\pi r\mathbf{j} = 2rh\lambda\mathbf{i} + r^2\lambda\mathbf{j}$$

- Equating, $2\pi h + 4\pi r = 2\pi rh\lambda$ and $2\pi r = r^2\lambda$.
- Since $r \neq 0$, $\lambda = 2\pi/r$ from the second equation. Substituting this in the first equation yields $h = 2r$.

Examples

- Since

$$\nabla f(r, h) = (2\pi h + 4\pi r)\mathbf{i} + 2\pi r\mathbf{j}, \quad \nabla g(r, h) = 2rh\mathbf{i} + r^2\mathbf{j}.$$

So $\nabla f(r, h) = \lambda \nabla g(r, h)$ becomes

$$(2\pi h + 4\pi r)\mathbf{i} + 2\pi r\mathbf{j} = 2rh\lambda\mathbf{i} + r^2\lambda\mathbf{j}$$

- Equating, $2\pi h + 4\pi r = 2\pi rh\lambda$ and $2\pi r = r^2\lambda$.
- Since $r \neq 0$, $\lambda = 2\pi/r$ from the second equation. Substituting this in the first equation yields $h = 2r$.
- Now, substitute this value of h in the other equation

$$r^2h - 16 = 0$$

Examples

- Since

$$\nabla f(r, h) = (2\pi h + 4\pi r)\mathbf{i} + 2\pi r\mathbf{j}, \quad \nabla g(r, h) = 2rh\mathbf{i} + r^2\mathbf{j}.$$

So $\nabla f(r, h) = \lambda \nabla g(r, h)$ becomes

$$(2\pi h + 4\pi r)\mathbf{i} + 2\pi r\mathbf{j} = 2rh\lambda\mathbf{i} + r^2\lambda\mathbf{j}$$

- Equating, $2\pi h + 4\pi r = 2\pi rh\lambda$ and $2\pi r = r^2\lambda$.
- Since $r \neq 0$, $\lambda = 2\pi/r$ from the second equation. Substituting this in the first equation yields $h = 2r$.
- Now, substitute this value of h in the other equation

$$r^2h - 16 = 0$$

- Solving, we get $r = 2, h = 4$.

Examples

Example. Find the maximum value of the function $f(x, y) = x^2 + y^2$ subject to the condition $x^2 - 2x + y^2 - 4y = 0$.

Examples

Example. Find the maximum value of the function $f(x, y) = x^2 + y^2$ subject to the condition $x^2 - 2x + y^2 - 4y = 0$.

Solution. We set up the Lagrange multiplier method.

Examples

Example. Find the maximum value of the function $f(x, y) = x^2 + y^2$ subject to the condition $x^2 - 2x + y^2 - 4y = 0$.

Solution. We set up the Lagrange multiplier method.

- Equating ∇f to $\lambda \nabla g$ gives:

$$2x\mathbf{i} + 2y\mathbf{j} = (2x - 2)\lambda\mathbf{i} + (2y - 4)\lambda\mathbf{j}.$$

Examples

Example. Find the maximum value of the function $f(x, y) = x^2 + y^2$ subject to the condition $x^2 - 2x + y^2 - 4y = 0$.

Solution. We set up the Lagrange multiplier method.

- Equating ∇f to $\lambda \nabla g$ gives:

$$2x\mathbf{i} + 2y\mathbf{j} = (2x - 2)\lambda\mathbf{i} + (2y - 4)\lambda\mathbf{j}.$$

- Solving, $x = \frac{\lambda}{\lambda - 1}$ and $y = \frac{2\lambda}{\lambda - 1}$, $\lambda \neq 1$ ($\lambda = 1$ implies $2x = 2x - 2$, impossible!)

Examples

Example. Find the maximum value of the function $f(x, y) = x^2 + y^2$ subject to the condition $x^2 - 2x + y^2 - 4y = 0$.

Solution. We set up the Lagrange multiplier method.

- Equating ∇f to $\lambda \nabla g$ gives:

$$2x\mathbf{i} + 2y\mathbf{j} = (2x - 2)\lambda\mathbf{i} + (2y - 4)\lambda\mathbf{j}.$$

- Solving, $x = \frac{\lambda}{\lambda - 1}$ and $y = \frac{2\lambda}{\lambda - 1}$, $\lambda \neq 1$ ($\lambda = 1$ implies $2x = 2x - 2$, impossible!)
- Now $g(x, y, z) = x^2 - 2x + y^2 - 4y = 0$, substituting the above values into this equation gives: $5\lambda^2 - 10\lambda = 0$.

Examples

Example. Find the maximum value of the function $f(x, y) = x^2 + y^2$ subject to the condition $x^2 - 2x + y^2 - 4y = 0$.

Solution. We set up the Lagrange multiplier method.

- Equating ∇f to $\lambda \nabla g$ gives:

$$2x\mathbf{i} + 2y\mathbf{j} = (2x - 2)\lambda\mathbf{i} + (2y - 4)\lambda\mathbf{j}.$$

- Solving, $x = \frac{\lambda}{\lambda - 1}$ and $y = \frac{2\lambda}{\lambda - 1}$, $\lambda \neq 1$ ($\lambda = 1$ implies $2x = 2x - 2$, impossible!)
- Now $g(x, y, z) = x^2 - 2x + y^2 - 4y = 0$, substituting the above values into this equation gives: $5\lambda^2 - 10\lambda = 0$.
- $\lambda = 0, 2$ or the points $(0, 0)$ and $(2, 4)$, where the values of the functions are 0 and 20 respectively. So 0 is minimum and 20 is maximum.

Examples

Example 3. Find the maximum and minimum values of the function $f(x, y, z) = x - 2y + 5z$ on the sphere $x^2 + y^2 + z^2 = 30$.

Examples

Example 3. Find the maximum and minimum values of the function $f(x, y, z) = x - 2y + 5z$ on the sphere $x^2 + y^2 + z^2 = 30$.

Solution. We set up the Lagrange multiplier method.

Examples

Example 3. Find the maximum and minimum values of the function $f(x, y, z) = x - 2y + 5z$ on the sphere $x^2 + y^2 + z^2 = 30$.

Solution. We set up the Lagrange multiplier method.

- $\nabla f = \lambda \nabla(x^2 + y^2 + z^2 - 30)$ and $x^2 + y^2 + z^2 - 30 = 0$.

Examples

Example 3. Find the maximum and minimum values of the function $f(x, y, z) = x - 2y + 5z$ on the sphere $x^2 + y^2 + z^2 = 30$.

Solution. We set up the Lagrange multiplier method.

- $\nabla f = \lambda \nabla(x^2 + y^2 + z^2 - 30)$ and $x^2 + y^2 + z^2 - 30 = 0$.
- $\mathbf{i} - 2\mathbf{j} + 5\mathbf{k} = 2x\lambda\mathbf{i} + 2y\lambda\mathbf{j} + 2z\lambda\mathbf{k}$.

Examples

Example 3. Find the maximum and minimum values of the function $f(x, y, z) = x - 2y + 5z$ on the sphere $x^2 + y^2 + z^2 = 30$.

Solution. We set up the Lagrange multiplier method.

- $\nabla f = \lambda \nabla(x^2 + y^2 + z^2 - 30)$ and $x^2 + y^2 + z^2 - 30 = 0$.
- $\mathbf{i} - 2\mathbf{j} + 5\mathbf{k} = 2x\lambda\mathbf{i} + 2y\lambda\mathbf{j} + 2z\lambda\mathbf{k}$.
- $2x\lambda = 1, 2y\lambda = -2, 2z\lambda = 5$.
- This implies $x = \frac{1}{2\lambda}, y = \frac{-1}{\lambda}, z = \frac{5}{2\lambda}, \lambda \neq 0$.

Examples

Example 3. Find the maximum and minimum values of the function $f(x, y, z) = x - 2y + 5z$ on the sphere $x^2 + y^2 + z^2 = 30$.

Solution. We set up the Lagrange multiplier method.

- $\nabla f = \lambda \nabla(x^2 + y^2 + z^2 - 30)$ and $x^2 + y^2 + z^2 - 30 = 0$.
- $\mathbf{i} - 2\mathbf{j} + 5\mathbf{k} = 2x\lambda\mathbf{i} + 2y\lambda\mathbf{j} + 2z\lambda\mathbf{k}$.
- $2x\lambda = 1, 2y\lambda = -2, 2z\lambda = 5$.
- This implies $x = \frac{1}{2\lambda}, y = \frac{-1}{\lambda}, z = \frac{5}{2\lambda}, \lambda \neq 0$.
- Substituting these into $g(x, y, z) = 0$ gives: $\lambda = \frac{1}{2}, -\frac{1}{2}$.

Examples

Example 3. Find the maximum and minimum values of the function $f(x, y, z) = x - 2y + 5z$ on the sphere $x^2 + y^2 + z^2 = 30$.

Solution. We set up the Lagrange multiplier method.

- $\nabla f = \lambda \nabla(x^2 + y^2 + z^2 - 30)$ and $x^2 + y^2 + z^2 - 30 = 0$.
- $\mathbf{i} - 2\mathbf{j} + 5\mathbf{k} = 2x\lambda\mathbf{i} + 2y\lambda\mathbf{j} + 2z\lambda\mathbf{k}$.
- $2x\lambda = 1, 2y\lambda = -2, 2z\lambda = 5$.
- This implies $x = \frac{1}{2\lambda}, y = \frac{-1}{\lambda}, z = \frac{5}{2\lambda}, \lambda \neq 0$.
- Substituting these into $g(x, y, z) = 0$ gives: $\lambda = \frac{1}{2}, -\frac{1}{2}$.
- The corresponding points are $(1, -2, 5), (-1, 2, -5)$ respectively and the values are 30 and -30 respectively which are the maximum and the minimum value in that order.

Lagrange Multipliers with Two Constraints

To find the extreme values of a differentiable function $f(x, y, z)$ whose variables are subject to two constraints. If the constraints are

$$g_1(x, y, z) = 0 \text{ and } g_2(x, y, z) = 0$$

and g_1 and g_2 are differentiable, with ∇g_1 not parallel to ∇g_2 , we find the constrained local maxima and minima of f by introducing two Lagrange multipliers λ and μ .

Lagrange Multipliers with Two Constraints

To find the extreme values of a differentiable function $f(x, y, z)$ whose variables are subject to two constraints. If the constraints are

$$g_1(x, y, z) = 0 \text{ and } g_2(x, y, z) = 0$$

and g_1 and g_2 are differentiable, with ∇g_1 not parallel to ∇g_2 , we find the constrained local maxima and minima of f by introducing two Lagrange multipliers λ and μ . That is, locate the points $P(x, y, z)$ where f takes on its constrained extreme values by finding the values of x, y, z, λ , and μ that simultaneously satisfy the equations

$$\nabla f = \lambda \nabla g_1 + \mu \nabla g_2, \quad g_1(x, y, z) = 0, \quad g_2(x, y, z) = 0.$$

The geometry of the setup is similar to the one constraint system.

- At the points where f takes local extremum, the gradient of f is perpendicular to the velocity vector of the curve C of intersection of the two surfaces $g_1 = 0$, $g_2 = 0$.

The geometry of the setup is similar to the one constraint system.

- At the points where f takes local extremum, the gradient of f is perpendicular to the velocity vector of the curve C of intersection of the two surfaces $g_1 = 0$, $g_2 = 0$.
- But ∇g_1 and ∇g_2 are also perpendicular to the velocity vector of C at those points.

The geometry of the setup is similar to the one constraint system.

- At the points where f takes local extremum, the gradient of f is perpendicular to the velocity vector of the curve C of intersection of the two surfaces $g_1 = 0$, $g_2 = 0$.
- But ∇g_1 and ∇g_2 are also perpendicular to the velocity vector of C at those points.
- Therefore, the gradient ∇f lies in the plane determined by ∇g_1 and ∇g_2 i.e., ∇f is a linear combination of the two gradient vectors ∇g_1 , ∇g_2 .

The geometry of the setup is similar to the one constraint system.

- At the points where f takes local extremum, the gradient of f is perpendicular to the velocity vector of the curve C of intersection of the two surfaces $g_1 = 0, g_2 = 0$.
- But ∇g_1 and ∇g_2 are also perpendicular to the velocity vector of C at those points.
- Therefore, the gradient ∇f lies in the plane determined by ∇g_1 and ∇g_2 i.e., ∇f is a linear combination of the two gradient vectors $\nabla g_1, \nabla g_2$.
- Hence we have λ_1, λ_2 such that $\nabla f = \lambda_1 \nabla g_1 + \lambda_2 \nabla g_2$

Examples

Example 1. Find the extreme values of the function $f(x, y, z) = x^2yz + 1$ on the intersection of the plane $z = 1$ and the sphere $x^2 + y^2 + z^2 = 10$.

Examples

Example 1. Find the extreme values of the function $f(x, y, z) = x^2yz + 1$ on the intersection of the plane $z = 1$ and the sphere $x^2 + y^2 + z^2 = 10$.

Solution. Let us set up the Lagrange multiplier system.

- $\nabla f = \lambda_1 \nabla(z - 1) + \lambda_2 \nabla(x^2 + y^2 + z^2 - 10)$

Examples

Example 1. Find the extreme values of the function $f(x, y, z) = x^2yz + 1$ on the intersection of the plane $z = 1$ and the sphere $x^2 + y^2 + z^2 = 10$.

Solution. Let us set up the Lagrange multiplier system.

- $\nabla f = \lambda_1 \nabla(z - 1) + \lambda_2 \nabla(x^2 + y^2 + z^2 - 10)$
- $2xyz = 2x\lambda_2, x^2z = 2y\lambda_2, x^2y = \lambda_1 + 2z\lambda_2.$

Examples

Example 1. Find the extreme values of the function $f(x, y, z) = x^2yz + 1$ on the intersection of the plane $z = 1$ and the sphere $x^2 + y^2 + z^2 = 10$.

Solution. Let us set up the Lagrange multiplier system.

- $\nabla f = \lambda_1 \nabla(z - 1) + \lambda_2 \nabla(x^2 + y^2 + z^2 - 10)$
- $2xyz = 2x\lambda_2, x^2z = 2y\lambda_2, x^2y = \lambda_1 + 2z\lambda_2.$
- From the first equation, either $x = 0$ or $yz = \lambda_2.$

Examples

Example 1. Find the extreme values of the function $f(x, y, z) = x^2yz + 1$ on the intersection of the plane $z = 1$ and the sphere $x^2 + y^2 + z^2 = 10$.

Solution. Let us set up the Lagrange multiplier system.

- $\nabla f = \lambda_1 \nabla(z - 1) + \lambda_2 \nabla(x^2 + y^2 + z^2 - 10)$
- $2xyz = 2x\lambda_2, x^2z = 2y\lambda_2, x^2y = \lambda_1 + 2z\lambda_2.$
- From the first equation, either $x = 0$ or $yz = \lambda_2.$
- If $x \neq 0$, then the second equation with $z = 1$ leads to $x^2 = 2y^2.$
Substituting this in the equation of the sphere, we get:

Examples

Example 1. Find the extreme values of the function $f(x, y, z) = x^2yz + 1$ on the intersection of the plane $z = 1$ and the sphere $x^2 + y^2 + z^2 = 10$.

Solution. Let us set up the Lagrange multiplier system.

- $\nabla f = \lambda_1 \nabla(z - 1) + \lambda_2 \nabla(x^2 + y^2 + z^2 - 10)$
- $2xyz = 2x\lambda_2, x^2z = 2y\lambda_2, x^2y = \lambda_1 + 2z\lambda_2.$
- From the first equation, either $x = 0$ or $yz = \lambda_2.$
- If $x \neq 0$, then the second equation with $z = 1$ leads to $x^2 = 2y^2.$

Substituting this in the equation of the sphere, we get:

$x = \sqrt{6}, -\sqrt{6}$ and $y = \sqrt{3}, -\sqrt{3}.$ The critical points would be $(\pm\sqrt{6}, \pm\sqrt{3}, 1).$ The function value would be $1 \pm 6\sqrt{3}.$

Examples

Example 1. Find the extreme values of the function $f(x, y, z) = x^2yz + 1$ on the intersection of the plane $z = 1$ and the sphere $x^2 + y^2 + z^2 = 10$.

Solution. Let us set up the Lagrange multiplier system.

- $\nabla f = \lambda_1 \nabla(z - 1) + \lambda_2 \nabla(x^2 + y^2 + z^2 - 10)$
- $2xyz = 2x\lambda_2, x^2z = 2y\lambda_2, x^2y = \lambda_1 + 2z\lambda_2$.
- From the first equation, either $x = 0$ or $yz = \lambda_2$.
- If $x \neq 0$, then the second equation with $z = 1$ leads to $x^2 = 2y^2$. Substituting this in the equation of the sphere, we get:
 $x = \sqrt{6}, -\sqrt{6}$ and $y = \sqrt{3}, -\sqrt{3}$. The critical points would be $(\pm\sqrt{6}, \pm\sqrt{3}, 1)$. The function value would be $1 \pm 6\sqrt{3}$.
- When $x = 0$, $y = \pm 3$ (use the equation of the sphere) or the points are $(0, \pm 3, 1)$ and the function values are 1.

Examples

Example 1. Find the extreme values of the function $f(x, y, z) = x^2yz + 1$ on the intersection of the plane $z = 1$ and the sphere $x^2 + y^2 + z^2 = 10$.

Solution. Let us set up the Lagrange multiplier system.

- $\nabla f = \lambda_1 \nabla(z - 1) + \lambda_2 \nabla(x^2 + y^2 + z^2 - 10)$
- $2xyz = 2x\lambda_2, x^2z = 2y\lambda_2, x^2y = \lambda_1 + 2z\lambda_2$.
- From the first equation, either $x = 0$ or $yz = \lambda_2$.
- If $x \neq 0$, then the second equation with $z = 1$ leads to $x^2 = 2y^2$. Substituting this in the equation of the sphere, we get:
 $x = \sqrt{6}, -\sqrt{6}$ and $y = \sqrt{3}, -\sqrt{3}$. The critical points would be $(\pm\sqrt{6}, \pm\sqrt{3}, 1)$. The function value would be $1 \pm 6\sqrt{3}$.
- When $x = 0$, $y = \pm 3$ (use the equation of the sphere) or the points are $(0, \pm 3, 1)$ and the function values are 1.
- The extreme values are $1 + 6\sqrt{3}, 1 - 6\sqrt{3}$ at the points $(\pm\sqrt{6}, \sqrt{3}, 1), (\pm\sqrt{6}, -\sqrt{3}, 1)$ respectively.

Examples

Example 2. Find the maximum volume of a box with a given total surface area.

Examples

Example 2. Find the maximum volume of a box with a given total surface area.

Solution. The volume $f(x, y, z) = xyz$ to be maximized given that the surface area $g(x, y, z) = 2(xy + yz + zx) - c = 0$.

Examples

Example 2. Find the maximum volume of a box with a given total surface area.

Solution. The volume $f(x, y, z) = xyz$ to be maximized given that the surface area $g(x, y, z) = 2(xy + yz + zx) - c = 0$.

First we find the gradients $\nabla f = yz\mathbf{i} + xz\mathbf{j} + xy\mathbf{k}$ and $\nabla g = 2((y + z)\mathbf{i} + (x + z)\mathbf{j} + (x + y)\mathbf{k})$ then we have to set up the Lagrange multiplier system.

Examples

Example 2. Find the maximum volume of a box with a given total surface area.

Solution. The volume $f(x, y, z) = xyz$ to be maximized given that the surface area $g(x, y, z) = 2(xy + yz + zx) - c = 0$.

First we find the gradients $\nabla f = yz\mathbf{i} + xz\mathbf{j} + xy\mathbf{k}$ and $\nabla g = 2((y + z)\mathbf{i} + (x + z)\mathbf{j} + (x + y)\mathbf{k})$ then we have to set up the Lagrange multiplier system.

$$\nabla f = \lambda \nabla g$$

gives the equations

Example cont.

$$yz = 2\lambda(y + z)$$

$$xz = 2\lambda(x + z)$$

$$xy = 2\lambda(x + y)$$

Example cont.

$$yz = 2\lambda(y + z)$$

$$xz = 2\lambda(x + z)$$

$$xy = 2\lambda(x + y)$$

Or we get

$$\frac{y}{x} = \frac{y + z}{x + z}$$

$$\frac{z}{y} = \frac{x + z}{x + y}$$

Example cont.

$$yz = 2\lambda(y + z)$$

$$xz = 2\lambda(x + z)$$

$$xy = 2\lambda(x + y)$$

Or we get

$$\frac{y}{x} = \frac{y + z}{x + z}$$

$$\frac{z}{y} = \frac{x + z}{x + y}$$

Solving for x, y, z , we get $x = y = z$. Now, utilize the equation $g = 0$ to obtain $x = y = z = \sqrt{\frac{c}{6}}$. So the maximum volume is $(\frac{c}{6})^{\frac{3}{2}}$.

Few notes from Lagrange multiplier method

- Lagrange multiplier finds extreme values of a function $f(x, y)$ (or $f(x, y, z)$) given one or more than one constraints, $g(x, y) = 0$ (or $g(x, y, z) = 0$).

Few notes from Lagrange multiplier method

- Lagrange multiplier finds extreme values of a function $f(x, y)$ (or $f(x, y, z)$) given one or more than one constraints, $g(x, y) = 0$ (or $g(x, y, z) = 0$).
- One can visualize it as finding the maximum or minimum values of the function $f(x, y)$ (or $f(x, y, z)$) on the curve $g(x, y) = 0$ (or surface $g(x, y, z) = 0$).

Few notes from Lagrange multiplier method

- Lagrange multiplier finds extreme values of a function $f(x, y)$ (or $f(x, y, z)$) given one or more than one constraints, $g(x, y) = 0$ (or $g(x, y, z) = 0$).
- One can visualize it as finding the maximum or minimum values of the function $f(x, y)$ (or $f(x, y, z)$) on the curve $g(x, y) = 0$ (or surface $g(x, y, z) = 0$).
- The method finds all the relative maximum or minimum for the function in the curve or surface defined by the constraint (s).
- $f(x, y)$ (or $f(x, y, z)$) is called the objective function, $g(x, y)$ (or $g(x, y, z)$) is called the constraint function.

Few notes from Lagrange multiplier method

- Lagrange multiplier finds extreme values of a function $f(x, y)$ (or $f(x, y, z)$) given one or more than one constraints, $g(x, y) = 0$ (or $g(x, y, z) = 0$).
- One can visualize it as finding the maximum or minimum values of the function $f(x, y)$ (or $f(x, y, z)$) on the curve $g(x, y) = 0$ (or surface $g(x, y, z) = 0$).
- The method finds all the relative maximum or minimum for the function in the curve or surface defined by the constraint (s).
- $f(x, y)$ (or $f(x, y, z)$) is called the objective function, $g(x, y)$ (or $g(x, y, z)$) is called the constraint function.
- The Lagrange multiplier method is only a necessary condition need not be sufficient.

Examples

- 1 A rectangular box without a lid is to be made from 27 m^2 of cardboard. Find the maximum volume of such a box.

Examples

- ① A rectangular box without a lid is to be made from 27 m^2 of cardboard. Find the maximum volume of such a box. ($\min xyz$ s.t $xy + 2yz + 2xz - 27 = 0$, Ans: 13.5 m^3)

Examples

- ① A rectangular box without a lid is to be made from 27 m^2 of cardboard. Find the maximum volume of such a box. ($\min xyz$ s.t $xy + 2yz + 2xz - 27 = 0$, Ans: 13.5 m^3)
- ② Find three real numbers whose sum is 9 and the sum of whose squares is as small as possible.

Examples

- ① A rectangular box without a lid is to be made from 27 m^2 of cardboard. Find the maximum volume of such a box. ($\min xyz$ s.t $xy + 2yz + 2xz - 27 = 0$, Ans: 13.5 m^3)
- ② Find three real numbers whose sum is 9 and the sum of whose squares is as small as possible. ($3, 3, 3$)

Examples

- ① A rectangular box without a lid is to be made from 27 m^2 of cardboard. Find the maximum volume of such a box. ($\min xyz$ s.t $xy + 2yz + 2xz - 27 = 0$, Ans: 13.5 m^3)
- ② Find three real numbers whose sum is 9 and the sum of whose squares is as small as possible. ($3, 3, 3$)
- ③ Find points on the curve $xy^2 = 54$ nearest to origin.

Examples

- ① A rectangular box without a lid is to be made from 27 m^2 of cardboard. Find the maximum volume of such a box. ($\min xyz$ s.t $xy + 2yz + 2xz - 27 = 0$, Ans: 13.5 m^3)
- ② Find three real numbers whose sum is 9 and the sum of whose squares is as small as possible. $(3, 3, 3)$
- ③ Find points on the curve $xy^2 = 54$ nearest to origin. $((3, \pm 3\sqrt{2}))$

Examples

- ① A rectangular box without a lid is to be made from 27 m^2 of cardboard. Find the maximum volume of such a box. ($\min xyz$ s.t $xy + 2yz + 2xz - 27 = 0$, Ans: 13.5 m^3)
- ② Find three real numbers whose sum is 9 and the sum of whose squares is as small as possible. $(3, 3, 3)$
- ③ Find points on the curve $xy^2 = 54$ nearest to origin. $((3, \pm 3\sqrt{2}))$
- ④ The temperature at a point (x, y) on a metal plate is $T(x, y) = 4x^2 - 4xy + y^2$. An ant on the plate walks around the circle of radius 5 centered at the origin. What are the highest and lowest temperatures encountered by the ant?

Examples

- ① A rectangular box without a lid is to be made from 27 m^2 of cardboard. Find the maximum volume of such a box. ($\min xyz$ s.t $xy + 2yz + 2xz - 27 = 0$, Ans: 13.5 m^3)
- ② Find three real numbers whose sum is 9 and the sum of whose squares is as small as possible. ($3, 3, 3$)
- ③ Find points on the curve $xy^2 = 54$ nearest to origin. ($(3, \pm 3\sqrt{2})$)
- ④ The temperature at a point (x, y) on a metal plate is $T(x, y) = 4x^2 - 4xy + y^2$. An ant on the plate walks around the circle of radius 5 centered at the origin. What are the highest and lowest temperatures encountered by the ant? ($0^\circ, 125^\circ$)
- ⑤ Find the extreme values of the function $f(x, y, z) = xy + z^2$ on the circle in which the plane $y - x = 0$ intersects the sphere $x^2 + y^2 + z^2 = 4$.

Examples

- ① A rectangular box without a lid is to be made from 27 m^2 of cardboard. Find the maximum volume of such a box. ($\min xyz$ s.t $xy + 2yz + 2xz - 27 = 0$, Ans: 13.5 m^3)
- ② Find three real numbers whose sum is 9 and the sum of whose squares is as small as possible. ($3, 3, 3$)
- ③ Find points on the curve $xy^2 = 54$ nearest to origin. ($(3, \pm 3\sqrt{2})$)
- ④ The temperature at a point (x, y) on a metal plate is $T(x, y) = 4x^2 - 4xy + y^2$. An ant on the plate walks around the circle of radius 5 centered at the origin. What are the highest and lowest temperatures encountered by the ant? ($0^\circ, 125^\circ$)
- ⑤ Find the extreme values of the function $f(x, y, z) = xy + z^2$ on the circle in which the plane $y - x = 0$ intersects the sphere $x^2 + y^2 + z^2 = 4$. ($f(0, 0, \pm 2) = 4, f(\pm\sqrt{2}, \pm\sqrt{2}, 0) = 2$)