Lecture 19

SIMPLE HARMONIC MOTION FORCED HARMONIC MOTION

Radhika Vathsan, Physics@BITS-Goa, 2024

Lecture 19

SIMPLE HARMONIC MOTION FORCED HARMONIC MOTION

- Forced Oscillations
- Equation of motion and solutions

Radhika Vathsan, Physics@BITS-Goa, 2024



• An external driving force F(t) acts on an oscillator

- ullet An external driving force F(t) acts on an oscillator
- This continuously supplies energy to the system.

- An external driving force F(t) acts on an oscillator
- This continuously supplies energy to the system.
- Assumed to be harmonic: $F_0 \cos \omega t$: can solve the motion analytically.

- An external driving force F(t) acts on an oscillator
- This continuously supplies energy to the system.
- Assumed to be harmonic: $F_0 \cos \omega t$: can solve the motion analytically.
- Physical examples:

- An external driving force F(t) acts on an oscillator
- This continuously supplies energy to the system.
- Assumed to be harmonic: $F_0 \cos \omega t$: can solve the motion analytically.
- Physical examples:

- An external driving force F(t) acts on an oscillator
- This continuously supplies energy to the system.
- Assumed to be harmonic: $F_0 \cos \omega t$: can solve the motion analytically.
- Physical examples:
 - ▶ Light incident on an electron in an atom ⇒ atom emits light

- An external driving force F(t) acts on an oscillator
- This continuously supplies energy to the system.
- Assumed to be harmonic: $F_0 \cos \omega t$: can solve the motion analytically.
- Physical examples:
 - ▶ Light incident on an electron in an atom ⇒ atom emits light
 - lacktriangledown periodic gravity of the sun/moon \implies tides in a lake or ocean

- An external driving force F(t) acts on an oscillator
- This continuously supplies energy to the system.
- Assumed to be harmonic: $F_0 \cos \omega t$: can solve the motion analytically.
- Physical examples:
 - lacktriangle Light incident on an electron in an atom \Longrightarrow atom emits light
 - lacktriangledown periodic gravity of the sun/moon \implies tides in a lake or ocean
 - AC voltage supplied to an electrical oscillator (LCR)

- An external driving force F(t) acts on an oscillator
- This continuously supplies energy to the system.
- Assumed to be harmonic: $F_0 \cos \omega t$: can solve the motion analytically.
- Physical examples:
 - ▶ Light incident on an electron in an atom ⇒ atom emits light
 - ▶ periodic gravity of the sun/moon ⇒ tides in a lake or ocean
 - AC voltage supplied to an electrical oscillator (LCR)
 - Absorption lines in the spectra of stars

- An external driving force F(t) acts on an oscillator
- This continuously supplies energy to the system.
- Assumed to be harmonic: $F_0 \cos \omega t$: can solve the motion analytically.
- Physical examples:
 - ▶ Light incident on an electron in an atom ⇒ atom emits light
 - ▶ periodic gravity of the sun/moon ⇒ tides in a lake or ocean
 - AC voltage supplied to an electrical oscillator (LCR)
 - Absorption lines in the spectra of stars
 - Nuclear Magnetic Resonance (NMR)

Undamped oscillator: equation and solution

$$m\ddot{x} + kx =$$

Mathematical analysis for solution

Undamped oscillator: equation and solution

$$m\ddot{x} + kx = F_0 \cos \Omega t$$
: Driving Force $F(t)$.

Mathematical analysis for solution

Undamped oscillator: equation and solution

$$m\ddot{x} + kx = F_0 \cos \Omega t$$
: Driving Force $F(t)$.

Mathematical analysis for solution

Complexify: add $i(m\ddot{y} + ky = F_0 \sin \Omega t)$

Undamped oscillator: equation and solution

$$m\ddot{x} + kx = F_0 \cos \Omega t$$
: Driving Force $F(t)$.

Mathematical analysis for solution

Complexify: add $i(m\ddot{y} + ky = F_0 \sin \Omega t)$ to get $m\ddot{z} + kz = F_0 e^{i\Omega t}$;

Undamped oscillator: equation and solution

$$m\ddot{x} + kx = F_0 \cos \Omega t$$
: Driving Force $F(t)$.

Mathematical analysis for solution

Complexify: add $i(m\ddot{y} + ky = F_0 \sin \Omega t)$ to get $m\ddot{z} + kz = F_0 e^{i\Omega t}$; Trial exponential solution: $z = Re^{i(\omega t + \theta)}$, R, θ real.

$$\implies (-mR\omega^2 + kR)e^{i(\omega t + \theta)} = F_0e^{i\Omega t}$$

Undamped oscillator: equation and solution

$$m\ddot{x} + kx = F_0 \cos \Omega t$$
: Driving Force $F(t)$.

Mathematical analysis for solution

Complexify: add $i(m\ddot{y}+ky=F_0\sin\Omega t)$ to get $m\ddot{z}+kz=F_0e^{i\Omega t};$ Trial exponential solution: $z=Re^{i(\omega t+\theta)},\quad R,\theta$ real.

$$\implies (-mR\omega^2 + kR)e^{i(\omega t + \theta)} = F_0 e^{i\Omega t}$$
$$-mR\omega^2 + kR = F_0 e^{i((\Omega - \omega)t + \theta)}$$

Undamped oscillator: equation and solution

$$m\ddot{x} + kx = F_0 \cos \Omega t$$
: Driving Force $F(t)$.

Mathematical analysis for solution

Complexify: add $i(m\ddot{y} + ky = F_0 \sin \Omega t)$ to get $m\ddot{z} + kz = F_0 e^{i\Omega t}$; Trial exponential solution: $z = Re^{i(\omega t + \theta)}$, R, θ real.

$$\implies (-mR\omega^2 + kR)e^{i(\omega t + \theta)} = F_0 e^{i\Omega t}$$

$$-mR\omega^2 + kR = F_0 e^{i((\Omega - \omega)t + \theta)}$$

$$(-m\omega^2 + k)R = F_0 \cos((\Omega - \omega)t + \theta) + iF_0 \sin((\Omega - \omega)t + \theta)$$

Undamped oscillator: equation and solution

$$m\ddot{x} + kx = F_0 \cos \Omega t$$
: Driving Force $F(t)$.

Mathematical analysis for solution

Complexify: add $i(m\ddot{y} + ky = F_0 \sin \Omega t)$ to get $m\ddot{z} + kz = F_0 e^{i\Omega t}$; Trial exponential solution: $z = Re^{i(\omega t + \theta)}$, R, θ real.

$$\implies (-mR\omega^2 + kR)e^{i(\omega t + \theta)} = F_0 e^{i\Omega t}$$

$$-mR\omega^2 + kR = F_0 e^{i((\Omega - \omega)t + \theta)}$$

$$(-m\omega^2 + k)R = F_0 \cos((\Omega - \omega)t + \theta) + iF_0 \sin((\Omega - \omega)t + \theta)$$

(a) Imaginary part: $\sin((\Omega - \omega)t + \theta)) = 0$.

Since this must be true for all t, $\Omega = \omega$, $\theta = 0$

Real part:
$$(-m\Omega^2 + k)R = F_0$$
.

Real part:
$$(-m\Omega^2 + k)R = F_0$$
.

$$R = \frac{F_0}{k - m\Omega^2}$$

Real part:
$$(-m\Omega^2 + k)R = F_0$$
.

$$R = \frac{F_0}{k - m\Omega^2} = \frac{F_0}{m(\omega_0^2 - \Omega^2)}$$

Real part:
$$(-m\Omega^2 + k)R = F_0$$
.

$$R = \frac{F_0}{k - m\Omega^2} = \frac{F_0}{m(\omega_0^2 - \Omega^2)}$$

$$x(t) = R(\Omega)\cos\Omega t.$$

Undamped oscillator: equation and solution

Real part:
$$(-m\Omega^2 + k)R = F_0$$
.

$$R = \frac{F_0}{k - m\Omega^2} = \frac{F_0}{m(\omega_0^2 - \Omega^2)}$$

$$x(t) = R(\Omega)\cos\Omega t.$$

• How to satisfy initial conditions?

Undamped oscillator: equation and solution

$$\label{eq:Real part} \begin{split} & \text{Real part:} (-m\Omega^2+k)R = F_0. \\ & R = \frac{F_0}{k-m\Omega^2} = \frac{F_0}{m(\omega_0^2-\Omega^2)} \end{split}$$

$$x(t) = R(\Omega)\cos\Omega t.$$

How to satisfy initial conditions?
 2nd) order diffl eqn must have two independent solutions.

Real part:
$$(-m\Omega^2+k)R=F_0.$$

$$R=\frac{F_0}{k-m\Omega^2}=\frac{F_0}{m(\omega_0^2-\Omega^2)}$$

$$x(t) = R(\Omega)\cos\Omega t.$$

- How to satisfy initial conditions?
 2nd) order diffl eqn must have two independent solutions.
- Solution to homogeneous eqn (undriven):

$$x_2(t) = B\cos(\omega_0 t + \phi)$$
: "General solution"

Undamped oscillator: equation and solution

Real part:
$$(-m\Omega^2+k)R=F_0.$$

$$R=\frac{F_0}{k-m\Omega^2}=\frac{F_0}{m(\omega_0^2-\Omega^2)}$$

$$x(t) = R(\Omega)\cos\Omega t.$$

- How to satisfy initial conditions?
 2nd) order diffl eqn must have two independent solutions.
- Solution to homogeneous eqn (undriven):

$$x_2(t) = B\cos(\omega_0 t + \phi)$$
: "General solution"

Two constants B and ϕ : satisfy initial conditions.

Undamped oscillator: equation and solution

Real part:
$$(-m\Omega^2+k)R=F_0.$$

$$R=\frac{F_0}{k-m\Omega^2}=\frac{F_0}{m(\omega_0^2-\Omega^2)}$$

$$x_1(t) = R(\Omega) \cos \Omega t$$
 : "Particular Solution"

- How to satisfy initial conditions?
 2nd) order diffl eqn must have two independent solutions.
- Solution to homogeneous eqn (undriven):

$$x_2(t) = B\cos(\omega_0 t + \phi)$$
: "General solution"

Two constants B and ϕ : satisfy initial conditions.

Undamped oscillator: equation and solution

Real part:
$$(-m\Omega^2+k)R=F_0.$$

$$R=\frac{F_0}{k-m\Omega^2}=\frac{F_0}{m(\omega_0^2-\Omega^2)}$$

$$x_1(t) = R(\Omega) \cos \Omega t$$
 : "Particular Solution"

- How to satisfy initial conditions?
 2nd) order diffl eqn must have two independent solutions.
- Solution to homogeneous eqn (undriven):

$$x_2(t) = B\cos(\omega_0 t + \phi)$$
: "General solution"


Two constants B and ϕ : satisfy initial conditions.

Full Solution:
$$x(t) = B\cos(\omega_0 t + \phi) + R\cos\Omega t$$
.

Undamped oscillator: equation and solution

Analysis of solution:

•
$$R(\Omega) = \frac{F_0/m}{\omega_0^2 - \Omega^2}, \quad \omega_0^2 = \frac{k}{m}.$$




Undamped oscillator: equation and solution

Analysis of solution:

•
$$R(\Omega) = \frac{F_0/m}{\omega_0^2 - \Omega^2}, \quad \omega_0^2 = \frac{k}{m}.$$

 \bullet Amplitude blows up at $\Omega=\omega_0$




Undamped oscillator: equation and solution

Analysis of solution:

•
$$R(\Omega) = \frac{F_0/m}{\omega_0^2 - \Omega^2}, \quad \omega_0^2 = \frac{k}{m}.$$

- Amplitude blows up at $\Omega = \omega_0$
 - Resonance.




Undamped oscillator: equation and solution

Analysis of solution:

•
$$R(\Omega) = \frac{F_0/m}{\omega_0^2 - \Omega^2}, \quad \omega_0^2 = \frac{k}{m}.$$

- Amplitude blows up at $\Omega = \omega_0$
 - Resonance.
- R changes sign thru resonance: phase change of π .



Simple Harmonic Motion Equation of motion and solutions

Undamped system is unphysical

Amplitude has a problem at $\Omega=\omega_0$

Solution: consider damping.