

Double Integrals

Devika S

Department of Mathematics
BITS Pilani, K K Birla Goa Campus

October 30, 2024




Double Integrals

Let us discuss double integrals of function $f(x, y)$ defined on a rectangular region $R : a \leq x \leq b, c \leq y \leq d$.

Double Integrals

Let us discuss double integrals of function $f(x, y)$ defined on a rectangular region $R : a \leq x \leq b, c \leq y \leq d$.

Let us divide R into rectangles as shown in the figure below.



Double Integrals

The rectangles form a partition P of R . We number these rectangles 1 through n . Define **norm** of the partition $\|P\|$ by is the largest width or height of any rectangle in the partition P .

Double Integrals

The rectangles form a partition P of R . We number these rectangles 1 through n . Define **norm** of the partition $\|P\|$ by is the largest width or height of any rectangle in the partition P .

Let $\Delta A_k = \Delta x_k \Delta y_k$, here Δx_k and Δy_k are width and height of the k -th rectangle and let (x_k, y_k) be any point in the k -th rectangle.

Double Integrals

The rectangles form a partition P of R . We number these rectangles 1 through n . Define **norm** of the partition $\|P\|$ by is the largest width or height of any rectangle in the partition P .

Let $\Delta A_k = \Delta x_k \Delta y_k$, here Δx_k and Δy_k are width and height of the k -th rectangle and let (x_k, y_k) be any point in the k -th rectangle.

If the limit

$$\lim_{\|P\| \rightarrow 0} \sum_{k=1}^n f(x_k, y_k) \Delta A_k = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(x_k, y_k) \Delta A_k$$

exists, then f is said to be integrable over R and this limit is called the **double integral of f over R** , written as

$$\iint_R f(x, y) dA \quad \text{or} \quad \iint_R f(x, y) dx dy.$$


Double Integral as Volume

Double Integral as Volume


If $f(x, y)$ is a **positive continuous function** on a rectangular region R , then the **volume of the 3-dimensional solid region over the xy -plane bounded below by R and above by the surface $z = f(x, y)$** is given by the double integral

$$\text{Volume} = \iint_R f(x, y) dA.$$


Double Integral as Volume



Double Integral as Volume



(a) $n = 16$



(b) $n = 64$



(c) $n = 256$

Fubini's Theorem

Fubini's Theorem (First Version)

If $f(x, y)$ is continuous throughout the rectangular region $R: a \leq x \leq b, c \leq y \leq d$, then

$$\iint_R f(x, y) dA = \int_a^b \int_c^d f(x, y) dy dx = \int_c^d \int_a^b f(x, y) dx dy.$$

Fubini's Theorem

Fubini's Theorem (First Version)

If $f(x, y)$ is continuous throughout the rectangular region $R: a \leq x \leq b, c \leq y \leq d$, then

$$\iint_R f(x, y) dA = \int_a^b \int_c^d f(x, y) dy dx = \int_c^d \int_a^b f(x, y) dx dy.$$

- Fubini's theorem says that the **double integral** of any continuous function over a rectangle can be calculated as an **iterated or repeated integral** in either order of integration.

Examples

Example 1. $\iint_R (4 - y^2) dA$, where $R : 0 \leq x \leq 3, 0 \leq y \leq 2$.

Examples

Example 1. $\iint_R (4 - y^2) dA$, where $R : 0 \leq x \leq 3, 0 \leq y \leq 2$.

Solution: Use Fubini's Theorem.

Examples

Example 1. $\iint_R (4 - y^2) dA$, where $R : 0 \leq x \leq 3, 0 \leq y \leq 2$.

Solution: Use Fubini's Theorem.

$$\begin{aligned}\iint_R (4 - y^2) dA &= \int_0^3 \int_0^2 (4 - y^2) dy dx \\ &= \int_0^3 \left(4y - \frac{1}{3}y^3 \right) \Big|_0^2 dx \\ &= \int_0^3 \frac{16}{3} dx = \frac{16}{3} (x) \Big|_0^3 = 16.\end{aligned}$$

Examples

Example 1. $\iint_R (4 - y^2) dA$, where $R : 0 \leq x \leq 3, 0 \leq y \leq 2$.

Solution: Use Fubini's Theorem.

$$\begin{aligned}\iint_R (4 - y^2) dA &= \int_0^3 \int_0^2 (4 - y^2) dy dx \\ &= \int_0^3 \left(4y - \frac{1}{3}y^3 \right) \Big|_0^2 dx \\ &= \int_0^3 \frac{16}{3} dx = \frac{16}{3} (x) \Big|_0^3 = 16.\end{aligned}$$

If we reverse the order of integration, then

$$\int_0^2 (4 - y^2)(x) \Big|_0^3 dy = 3 \int_0^2 (4 - y^2) dy = 3 \times 16/3 = 16.$$

Examples

Example 2. $\iint_R xy \cos y dA$, where $R : -1 \leq x \leq 1, 0 \leq y \leq \pi$.

Examples

Example 2. $\iint_R xy \cos y dA$, where $R : -1 \leq x \leq 1, 0 \leq y \leq \pi$.

Solution:

$$\begin{aligned}\iint_R xy \cos y dA &= \int_{-1}^1 \int_0^{\pi} xy \cos y dy dx \\ &= \int_{-1}^1 (xy \sin y + x \cos y) \Big|_0^{\pi} dx = \int_{-1}^1 -2x dx = 0.\end{aligned}$$

Examples

Example 2. $\iint_R xy \cos y dA$, where $R : -1 \leq x \leq 1, 0 \leq y \leq \pi$.

Solution:

$$\begin{aligned}\iint_R xy \cos y dA &= \int_{-1}^1 \int_0^{\pi} xy \cos y dy dx \\ &= \int_{-1}^1 (xy \sin y + x \cos y) \Big|_0^{\pi} dx = \int_{-1}^1 -2x dx = 0.\end{aligned}$$

Example 3. $\iint_R \frac{y}{x^2 y^2 + 1} dA$, where $R : 0 \leq x \leq 1, 0 \leq y \leq 1$.

Examples

Example 2. $\iint_R xy \cos y dA$, where $R : -1 \leq x \leq 1, 0 \leq y \leq \pi$.

Solution:

$$\begin{aligned}\iint_R xy \cos y dA &= \int_{-1}^1 \int_0^\pi xy \cos y dy dx \\ &= \int_{-1}^1 (xy \sin y + x \cos y) \Big|_0^\pi dx = \int_{-1}^1 -2x dx = 0.\end{aligned}$$

Example 3. $\iint_R \frac{y}{x^2 y^2 + 1} dA$, where $R : 0 \leq x \leq 1, 0 \leq y \leq 1$.

Solution:

$$\begin{aligned}\iint_R \frac{y}{x^2 y^2 + 1} dA &= \int_0^1 \int_0^1 \frac{y}{(xy)^2 + 1} dx dy \\ &= \int_0^1 (\tan^{-1}(xy)) \Big|_0^1 dy = y \tan^{-1} y - \frac{1}{2} \ln(1 + y^2) \Big|_0^1 = \frac{\pi}{4} - \frac{1}{2} \ln 2.\end{aligned}$$

Examples

Example 4. Find the volume of the region bounded above by the plane $z = 2 - x - y$ and below by the square R : $0 \leq x \leq 1$, $0 \leq y \leq 1$.

Examples

Example 4. Find the volume of the region bounded above by the plane $z = 2 - x - y$ and below by the square R : $0 \leq x \leq 1$, $0 \leq y \leq 1$.

Solution: Volume V is given by

$$\begin{aligned} V &= \iint_R (2 - x - y) dA \\ &= \int_0^1 \int_0^1 (2 - x - y) dy dx = \int_0^1 \left(2y - xy - \frac{y^2}{2} \right) \Big|_0^1 dx \\ &= \int_0^1 \left(\frac{3}{2} - x \right) dx = \left(\frac{3}{2}x - \frac{x^2}{2} \right) \Big|_0^1 = 1. \end{aligned}$$

Double Integrals over Bounded, Nonrectangular Regions




FIGURE 15.8 A rectangular grid partitioning a bounded, nonrectangular region into rectangular cells.

Double Integrals over Bounded, Nonrectangular Regions

To define the double integral of a function $f(x, y)$ over a bounded, nonrectangular region R , we again begin by covering R with a grid of small rectangular cells whose union contains all points of R .

Double Integrals over Bounded, Nonrectangular Regions

To define the double integral of a function $f(x, y)$ over a bounded, nonrectangular region R , we again begin by covering R with a grid of small rectangular cells whose union contains all points of R .

This time, however, we cannot exactly fill R with a finite number of rectangles lying inside R , since its boundary is curved, and some of the small rectangles in the grid lie partly outside R .

Double Integrals over Bounded, Nonrectangular Regions

To define the double integral of a function $f(x, y)$ over a bounded, nonrectangular region R , we again begin by covering R with a grid of small rectangular cells whose union contains all points of R .

This time, however, we cannot exactly fill R with a finite number of rectangles lying inside R , since its boundary is curved, and some of the small rectangles in the grid lie partly outside R .

A partition of R is formed by taking the rectangles that lie completely inside it, not using any that are either partly or completely outside. For commonly arising regions, more and more of R is included as the norm of a partition approaches zero.

Double Integrals over Bounded, Nonrectangular Regions

To define the double integral of a function $f(x, y)$ over a bounded, nonrectangular region R , we again begin by covering R with a grid of small rectangular cells whose union contains all points of R .

This time, however, we cannot exactly fill R with a finite number of rectangles lying inside R , since its boundary is curved, and some of the small rectangles in the grid lie partly outside R .

A partition of R is formed by taking the rectangles that lie completely inside it, not using any that are either partly or completely outside. For commonly arising regions, more and more of R is included as the norm of a partition approaches zero.

Double Integrals over Bounded, Nonrectangular Regions

Once we have a partition of R , we number the rectangles in some order from 1 to n and let ΔA_k be the area of the k th rectangle. We then choose a point (x_k, y_k) in the k th rectangle and form the Riemann sum:

$$S_n = \sum_{k=1}^n f(x_k, y_k) \Delta A_k.$$

As the norm of the partition forming S_n goes to zero, $\|P\| \rightarrow 0$, the width and height of each enclosed rectangle goes to zero and their number goes to infinity. If $f(x, y)$ is a continuous function, then these Riemann sums converge to a limiting value, not dependent on any of the choices we made. This limit is called the **double integral** of $f(x, y)$ over R :

$$\iint_R f(x, y) dA := \lim_{\|P\| \rightarrow 0} \sum_{k=1}^n f(x_k, y_k) \Delta A_k.$$

Fubini's Theorem General Version

Theorem

Let $f(x, y)$ be a continuous function on a region R .

- 1 If the region R is given by $a \leq x \leq b, g_1(x) \leq y \leq g_2(x)$, with g_1, g_2 continuous on the interval $[a, b]$, then

Fubini's Theorem General Version

Theorem

Let $f(x, y)$ be a continuous function on a region R .

- 1 If the region R is given by $a \leq x \leq b$, $g_1(x) \leq y \leq g_2(x)$, with g_1, g_2 continuous on the interval $[a, b]$, then

$$\iint_R f(x, y) dA = \int_a^b \int_{g_1(x)}^{g_2(x)} f(x, y) dy dx.$$

Fubini's Theorem General Version

Theorem

Let $f(x, y)$ be a continuous function on a region R .

- ① If the region R is given by $a \leq x \leq b$, $g_1(x) \leq y \leq g_2(x)$, with g_1, g_2 continuous on the interval $[a, b]$, then

$$\iint_R f(x, y) dA = \int_a^b \int_{g_1(x)}^{g_2(x)} f(x, y) dy dx.$$

- ② If R is given by $c \leq y \leq d$, $h_1(y) \leq x \leq h_2(y)$, with h_1, h_2 continuous on $[c, d]$, then

Fubini's Theorem General Version

Theorem

Let $f(x, y)$ be a continuous function on a region R .

- ① If the region R is given by $a \leq x \leq b$, $g_1(x) \leq y \leq g_2(x)$, with g_1, g_2 continuous on the interval $[a, b]$, then

$$\iint_R f(x, y) dA = \int_a^b \int_{g_1(x)}^{g_2(x)} f(x, y) dy dx.$$

- ② If R is given by $c \leq y \leq d$, $h_1(y) \leq x \leq h_2(y)$, with h_1, h_2 continuous on $[c, d]$, then

$$\iint_R f(x, y) dA = \int_c^d \int_{h_1(y)}^{h_2(y)} f(x, y) dx dy$$

Fubini's Theorem General Version

Theorem

Let $f(x, y)$ be a continuous function on a region R .

- ① If the region R is given by $a \leq x \leq b$, $g_1(x) \leq y \leq g_2(x)$, with g_1, g_2 continuous on the interval $[a, b]$, then

$$\iint_R f(x, y) dA = \int_a^b \int_{g_1(x)}^{g_2(x)} f(x, y) dy dx.$$

- ② If R is given by $c \leq y \leq d$, $h_1(y) \leq x \leq h_2(y)$, with h_1, h_2 continuous on $[c, d]$, then

$$\iint_R f(x, y) dA = \int_c^d \int_{h_1(y)}^{h_2(y)} f(x, y) dx dy$$

In a double integral, the outer limits must be constant, but the inner limits can depend on the outer variable.

Examples


Example 1. Sketch the region of integration:

① $0 \leq x \leq 3, 0 \leq y \leq 2x$

Examples

Example 1. Sketch the region of integration:


① $0 \leq x \leq 3, 0 \leq y \leq 2x$



Examples

Example 1. Sketch the region of integration:

① $0 \leq x \leq 3, 0 \leq y \leq 2x$




② $-1 \leq x \leq 2, x - 1 \leq y \leq x^2$


Examples

Example 1. Sketch the region of integration:

① $0 \leq x \leq 3, 0 \leq y \leq 2x$



② $-1 \leq x \leq 2, x - 1 \leq y \leq x^2$




Examples

③ $-2 \leq y \leq 2, y^2 \leq x \leq 4$


Examples

③ $-2 \leq y \leq 2, y^2 \leq x \leq 4$



Examples


③ $-2 \leq y \leq 2, y^2 \leq x \leq 4$




④ $0 \leq y \leq 1, y \leq x \leq 2y$

Examples

③ $-2 \leq y \leq 2, y^2 \leq x \leq 4$



④ $0 \leq y \leq 1, y \leq x \leq 2y$




Examples

5 $0 \leq x \leq 1, e^x \leq y \leq e$


Examples

⑤ $0 \leq x \leq 1, e^x \leq y \leq e$



Examples

⑤ $0 \leq x \leq 1, e^x \leq y \leq e$




Example 2. Sketch the region of integration and evaluate the integral

$$\int_0^1 \int_0^{y^2} 3y^3 e^{xy} dx dy.$$

Examples

⑤ $0 \leq x \leq 1, e^x \leq y \leq e$



Example 2. Sketch the region of integration and evaluate the integral


$$\int_0^1 \int_0^{y^2} 3y^3 e^{xy} dx dy.$$

Solution:

$$\int_0^1 \int_0^{y^2} 3y^3 e^{xy} dx dy = \int_0^1 \left[3y^3 \frac{e^{xy}}{y} \right]_0^{y^2} dy$$

Examples

⑤ $0 \leq x \leq 1, e^x \leq y \leq e$



Example 2. Sketch the region of integration and evaluate the integral

$$\int_0^1 \int_0^{y^2} 3y^3 e^{xy} dx dy.$$

Solution:

$$\begin{aligned} \int_0^1 \int_0^{y^2} 3y^3 e^{xy} dx dy &= \int_0^1 \left[3y^3 \frac{e^{xy}}{y} \right]_0^{y^2} dy \\ &= \int_0^1 (3y^2 e^{y^3} - 3y^2) dy = e^{y^3} - y^3 \Big|_0^1 = e - 2. \end{aligned}$$

How to find the limits of integration

- Calculate $\iint_R f(x, y) dA$ where the region R is bounded by the x -axis and the lines $x = 1$, $y = x$.

How to find the limits of integration

- Calculate $\iint_R f(x, y) dA$ where the region R is bounded by the x -axis and the lines $x = 1$, $y = x$.

How to find the limits of integration

- Calculate $\iint_R f(x, y) dA$ where the region R is bounded by the x -axis and the lines $x = 1$, $y = x$.

Using vertical cross-sections: For a given region R , if we want to integrate first with respect to y then :

How to find the limits of integration

- Calculate $\iint_R f(x, y) dA$ where the region R is bounded by the x -axis and the lines $x = 1$, $y = x$.

Using vertical cross-sections: For a given region R , if we want to integrate first with respect to y then :

- 1 Sketch the region and label the curves bounding the region.

How to find the limits of integration

- Calculate $\iint_R f(x, y) dA$ where the region R is bounded by the x -axis and the lines $x = 1$, $y = x$.

Using vertical cross-sections: For a given region R , if we want to integrate first with respect to y then :

- ① Sketch the region and label the curves bounding the region.
- ② Find the y limits:

How to find the limits of integration

- Calculate $\iint_R f(x, y) dA$ where the region R is bounded by the x -axis and the lines $x = 1$, $y = x$.

Using vertical cross-sections: For a given region R , if we want to integrate first with respect to y then :

- 1 **Sketch** the region and label the curves bounding the region.
- 2 **Find the y limits**: Look at a **vertical line** through $(x, 0)$ and cutting through the region R

How to find the limits of integration

- Calculate $\iint_R f(x, y) dA$ where the region R is bounded by the x -axis and the lines $x = 1$, $y = x$.

Using vertical cross-sections: For a given region R , if we want to integrate first with respect to y then :

- 1 **Sketch** the region and label the curves bounding the region.
- 2 **Find the y limits**: Look at a **vertical line** through $(x, 0)$ and cutting through the region R then y values as a function of x where the line enters and leaves the region

How to find the limits of integration

- Calculate $\iint_R f(x, y) dA$ where the region R is bounded by the x -axis and the lines $x = 1$, $y = x$.

Using vertical cross-sections: For a given region R , if we want to integrate first with respect to y then :

- 1 **Sketch** the region and label the curves bounding the region.
- 2 **Find the y limits**: Look at a **vertical line** through $(x, 0)$ and cutting through the region R then y values as a function of x where the line enters and leaves the region are the limits for y .

How to find the limits of integration

- Calculate $\iint_R f(x, y) dA$ where the region R is bounded by the x -axis and the lines $x = 1$, $y = x$.

Using vertical cross-sections: For a given region R , if we want to integrate first with respect to y then :

- 1 **Sketch** the region and label the curves bounding the region.
- 2 **Find the y limits**: Look at a **vertical line** through $(x, 0)$ and cutting through the region R then y values as a function of x where the line enters and leaves the region are the limits for y .
- 3 **Find the x limits**: The x limits of integration are the extreme x values of the region.

How to find the limits of integration

- Calculate $\iint_R f(x, y) dA$ where the region R is bounded by the x -axis and the lines $x = 1$, $y = x$.

Using vertical cross-sections: For a given region R , if we want to integrate first with respect to y then :

- 1 **Sketch** the region and label the curves bounding the region.
- 2 **Find the y limits**: Look at a **vertical line** through $(x, 0)$ and cutting through the region R then y values as a function of x where the line enters and leaves the region are the limits for y .
- 3 **Find the x limits**: The x limits of integration are the extreme x values of the region.

Using horizontal cross-sections: If we want to **integrate with respect to x first**, we simply choose **horizontal lines** to find the x limits as functions of y .

Properties of Double Integrals

If $f(x, y)$ and $g(x, y)$ are continuous on the bounded region R , then the following properties hold:

① **Constant Multiple:**
$$\iint_R cf(x, y)dA = c \iint_R f(x, y)dA \quad (c \in \mathbb{R})$$

② **Sum and Difference:**

$$\iint_R (f(x, y) \pm g(x, y))dA = \iint_R f(x, y)dA \pm \iint_R g(x, y)dA$$

③ **Monotonicity:**
$$\iint_R f(x, y)dA \geq \iint_R g(x, y)dA, \text{ if } f(x, y) \geq g(x, y)$$

④ **Additivity:**
$$\iint_R f(x, y)dA = \iint_{R_1} f(x, y)dA + \iint_{R_2} f(x, y)dA,$$

if R is the union of two nonoverlapping regions R_1 and R_2 .

Examples

Example 1. Write an iterated integral for $\iint_R dA$ over the described region R using (a) vertical cross-sections, (b) horizontal cross-sections.

Examples

Example 1. Write an iterated integral for $\iint_R dA$ over the described region R using (a) vertical cross-sections, (b) horizontal cross-sections.

- Bounded by $y = \sqrt{x}$, $y = 0$, and $x = 9$

Examples

Example 1. Write an iterated integral for $\iint_R dA$ over the described region R using (a) vertical cross-sections, (b) horizontal cross-sections.

- Bounded by $y = \sqrt{x}$, $y = 0$, and $x = 9$

$$(a) \int_0^9 \int_0^{\sqrt{x}} dy dx,$$

Examples

Example 1. Write an iterated integral for $\iint_R dA$ over the described region R using (a) vertical cross-sections, (b) horizontal cross-sections.

- Bounded by $y = \sqrt{x}$, $y = 0$, and $x = 9$

$$(a) \int_0^9 \int_0^{\sqrt{x}} dy dx, \quad (b) \int_0^3 \int_{y^2}^9 dx dy.$$

Examples

Example 1. Write an iterated integral for $\iint_R dA$ over the described region R using (a) vertical cross-sections, (b) horizontal cross-sections.

- Bounded by $y = \sqrt{x}$, $y = 0$, and $x = 9$

$$(a) \int_0^9 \int_0^{\sqrt{x}} dy dx, \quad (b) \int_0^3 \int_{y^2}^9 dx dy.$$

- Bounded by $y = \tan x$, $x = 0$, and $y = 1$.

Examples

Example 1. Write an iterated integral for $\iint_R dA$ over the described region R using (a) vertical cross-sections, (b) horizontal cross-sections.

- Bounded by $y = \sqrt{x}$, $y = 0$, and $x = 9$

$$(a) \int_0^9 \int_0^{\sqrt{x}} dy dx, \quad (b) \int_0^3 \int_{y^2}^9 dx dy.$$

- Bounded by $y = \tan x$, $x = 0$, and $y = 1$.

$$(a) \int_0^{\pi/4} \int_{\tan x}^1 dy dx,$$

Examples

Example 1. Write an iterated integral for $\iint_R dA$ over the described region R using (a) vertical cross-sections, (b) horizontal cross-sections.

- Bounded by $y = \sqrt{x}$, $y = 0$, and $x = 9$

$$(a) \int_0^9 \int_0^{\sqrt{x}} dy dx, \quad (b) \int_0^3 \int_{y^2}^9 dx dy.$$

- Bounded by $y = \tan x$, $x = 0$, and $y = 1$.

$$(a) \int_0^{\pi/4} \int_{\tan x}^1 dy dx, \quad (b) \int_0^1 \int_0^{\tan^{-1} y} dx dy.$$

Examples

Example 2. Calculate $\iint_R \frac{\sin x}{x} dA$ where the region R is bounded by the x -axis and the lines $x = 1$, $y = x$.

Examples

Example 2. Calculate $\iint_R \frac{\sin x}{x} dA$ where the region R is bounded by the x -axis and the lines $x = 1$, $y = x$.

Solution. Let us first integrate with respect to y ,

Examples

Example 2. Calculate $\iint_R \frac{\sin x}{x} dA$ where the region R is bounded by the x -axis and the lines $x = 1$, $y = x$.

Solution. Let us first integrate with respect to y , we get

$$\begin{aligned} \int_0^1 \left(\int_0^x \frac{\sin x}{x} dy \right) dx &= \int_0^1 \left(\frac{\sin x}{x} y \right) \Big|_0^x dx \\ &= \int_0^1 \frac{\sin x}{x} x dx = \int_0^1 \sin x dx = -\cos x \Big|_0^1 = -\cos 1 + 1. \end{aligned}$$

Examples

Example 2. Calculate $\iint_R \frac{\sin x}{x} dA$ where the region R is bounded by the x -axis and the lines $x = 1$, $y = x$.

Solution. Let us first integrate with respect to y , we get

$$\begin{aligned} \int_0^1 \left(\int_0^x \frac{\sin x}{x} dy \right) dx &= \int_0^1 \left(\frac{\sin x}{x} y \right) \Big|_0^x dx \\ &= \int_0^1 \frac{\sin x}{x} x dx = \int_0^1 \sin x dx = -\cos x \Big|_0^1 = -\cos 1 + 1. \end{aligned}$$

If we try to integrate with x first then we have to integrate

$$\int_0^1 \int_y^1 \frac{\sin x}{x} dx,$$

Examples

Example 2. Calculate $\iint_R \frac{\sin x}{x} dA$ where the region R is bounded by the x -axis and the lines $x = 1$, $y = x$.

Solution. Let us first integrate with respect to y , we get

$$\begin{aligned} \int_0^1 \left(\int_0^x \frac{\sin x}{x} dy \right) dx &= \int_0^1 \left(\frac{\sin x}{x} y \right) \Big|_0^x dx \\ &= \int_0^1 \frac{\sin x}{x} x dx = \int_0^1 \sin x dx = -\cos x \Big|_0^1 = -\cos 1 + 1. \end{aligned}$$

If we try to integrate with x first then we have to integrate

$$\int_0^1 \int_y^1 \frac{\sin x}{x} dx,$$

we run into a problem because $\int \frac{\sin x}{x} dx$ cannot be expressed in terms of elementary functions (there is no simple antiderivative)!

Examples

Example 3. Find the integral of $f(u, v) = v - \sqrt{u}$ over the triangular region of the first quadrant of uv plane bounded above by the line $u + v = 1$.

Examples

Example 3. Find the integral of $f(u, v) = v - \sqrt{u}$ over the triangular region of the first quadrant of uv plane bounded above by the line $u + v = 1$.

Solution: Let us first find the limits. Let us integrate with respect to u first,

Examples

Example 3. Find the integral of $f(u, v) = v - \sqrt{u}$ over the triangular region of the first quadrant of uv plane bounded above by the line $u + v = 1$.

Solution: Let us first find the limits. Let us integrate with respect to u first, so we have to find the limits of u as functions of v ,

Examples

Example 3. Find the integral of $f(u, v) = v - \sqrt{u}$ over the triangular region of the first quadrant of uv plane bounded above by the line $u + v = 1$.

Solution: Let us first find the limits. Let us integrate with respect to u first, so we have to find the limits of u as functions of v , we get $u = 0, u = 1 - v$.

Examples

Example 3. Find the integral of $f(u, v) = v - \sqrt{u}$ over the triangular region of the first quadrant of uv plane bounded above by the line $u + v = 1$.

Solution: Let us first find the limits. Let us integrate with respect to u first, so we have to find the limits of u as functions of v , we get $u = 0, u = 1 - v$.

And v ranges from 0 to 1.

Examples

Example 3. Find the integral of $f(u, v) = v - \sqrt{u}$ over the triangular region of the first quadrant of uv plane bounded above by the line $u + v = 1$.

Solution: Let us first find the limits. Let us integrate with respect to u first, so we have to find the limits of u as functions of v , we get $u = 0, u = 1 - v$.

And v ranges from 0 to 1. So the integral turns out to be:

Examples

Example 3. Find the integral of $f(u, v) = v - \sqrt{u}$ over the triangular region of the first quadrant of uv plane bounded above by the line $u + v = 1$.

Solution: Let us first find the limits. Let us integrate with respect to u first, so we have to find the limits of u as functions of v , we get $u = 0, u = 1 - v$.

And v ranges from 0 to 1. So the integral turns out to be:

$$\begin{aligned}\int_0^1 \int_0^{1-v} (v - \sqrt{u}) du dv &= \int_0^1 \left[vu - \frac{2}{3} u^{\frac{3}{2}} \right]_0^{1-v} dv \\ &= \int_0^1 \left(v - v^2 - \frac{2}{3} (1-v)^{\frac{3}{2}} \right) dv \\ &= \frac{1}{2} v^2 - \frac{1}{3} v^3 + \frac{4}{15} (1-v)^{\frac{5}{2}} \Big|_0^1 = -\frac{1}{10}.\end{aligned}$$

Examples

Example 4. Sketch the region and evaluate $\int_{-2}^0 \int_v^{-v} 2dpdv$.

Examples

Example 4. Sketch the region and evaluate $\int_{-2}^0 \int_v^{-v} 2dpdv$.

Solution: The integral is

$$\begin{aligned}\int_{-2}^0 \int_v^{-v} 2dpdv &= \int_{-2}^0 \left[2p \right]_v^{-v} dv = \int_{-2}^0 (-4v)dv \\ &= -4 \left[\frac{1}{2}v^2 \right]_{-2}^0 = 8.\end{aligned}$$

Examples

Example 4. Sketch the region and evaluate $\int_{-2}^0 \int_v^{-v} 2dpdv$.

Solution: The integral is

$$\begin{aligned}\int_{-2}^0 \int_v^{-v} 2dpdv &= \int_{-2}^0 \left[2p \right]_v^{-v} dv = \int_{-2}^0 (-4v) dv \\ &= -4 \left[\frac{1}{2} v^2 \right]_{-2}^0 = 8.\end{aligned}$$

Example 5. Find the volume of the space enclosed by the paraboloid $z = x^2 + y^2$ and below by the triangle enclosed by the lines $y = x, x = 0, x + y = 2$ in the xy plane.

Examples

Example 4. Sketch the region and evaluate $\int_{-2}^0 \int_v^{-v} 2dpdv$.

Solution: The integral is

$$\begin{aligned}\int_{-2}^0 \int_v^{-v} 2dpdv &= \int_{-2}^0 \left[2p \right]_v^{-v} dv = \int_{-2}^0 (-4v)dv \\ &= -4 \left[\frac{1}{2}v^2 \right]_{-2}^0 = 8.\end{aligned}$$

Example 5. Find the volume of the space enclosed by the paraboloid $z = x^2 + y^2$ and below by the triangle enclosed by the lines $y = x, x = 0, x + y = 2$ in the xy plane.

Solution. The volume is by definition given by the integral

$$\iint_R (x^2 + y^2) dA \text{ where the region } R \text{ is stated as above.}$$

Examples

Let us say that we want to first integrate with respect to y first then let us find the limits of y as functions of x . The limits are $y = x$ and $y = 2 - x$, and x varies from 0 to 1.

$$\begin{aligned}\int_0^1 \int_x^{2-x} (x^2 + y^2) dy dx &= \int_0^1 \left[\frac{1}{3} y^3 + x^2 y \right]_x^{2-x} dx \\ &= \int_0^1 \left(2x^2 - \frac{7}{3} x^3 + \frac{1}{3} (2-x)^3 \right) dx \\ &= \left[\frac{2}{3} x^3 - \frac{7}{12} x^4 - \frac{1}{12} (2-x)^4 \right]_0^1 = \frac{4}{3}.\end{aligned}$$

Examples

Let us say that we want to first integrate with respect to y first then let us find the limits of y as functions of x . The limits are $y = x$ and $y = 2 - x$, and x varies from 0 to 1.

$$\begin{aligned}\int_0^1 \int_x^{2-x} (x^2 + y^2) dy dx &= \int_0^1 \left[\frac{1}{3} y^3 + x^2 y \right]_x^{2-x} dx \\ &= \int_0^1 \left(2x^2 - \frac{7}{3} x^3 + \frac{1}{3} (2-x)^3 \right) dx \\ &= \left[\frac{2}{3} x^3 - \frac{7}{12} x^4 - \frac{1}{12} (2-x)^4 \right]_0^1 = \frac{4}{3}.\end{aligned}$$

Note: If you want to integrate w.r.t x first, consider two regions $R = R_1 \cup R_2$, first region where $R_1 : 0 \leq x \leq y, 0 \leq y \leq 1$ and the second region where $R_2 : 0 \leq x \leq 2 - y, 0 \leq y \leq 1$.