

MATH F111- Mathematics I

Saranya G. Nair
Department of Mathematics

BITS Pilani

September 2, 2024



Definition

A power series about $x = 0$ is a series of the form

$$\sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 x + c_2 x^2 + \cdots + c_n x^n + \cdots$$

A power series about $x = a$ is a series of the form

$$\sum_{n=0}^{\infty} c_n (x - a)^n = c_0 + c_1 (x - a) + c_2 (x - a)^2 + \cdots + c_n (x - a)^n + \cdots$$

in which the center a and the coefficients $c_0, c_1, c_2, \cdots, c_n, \cdots$ are constants.

When we fix a value for x , say $x = 1$, the power series

$$\sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 + c_2 + \cdots + c_n + \cdots$$

is an infinite series whose convergence or divergence can be investigated.

When we fix a value for x , say $x = 1$, the power series

$$\sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 + c_2 + \cdots + c_n + \cdots$$

is an infinite series whose convergence or divergence can be investigated.

For $x = 2$, the power series $\sum_{n=0}^{\infty} c_n x^n = c_0 + 2c_1 + 4c_2 + \cdots + 2^n c_n + \cdots$.

If for an x , the series $\sum_{n=0}^{\infty} c_n x^n$ converges we can use the limit of partial sequences to define a function f at x .

When we fix a value for x , say $x = 1$, the power series

$$\sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 + c_2 + \cdots + c_n + \cdots$$

is an infinite series whose convergence or divergence can be investigated.

For $x = 2$, the power series $\sum_{n=0}^{\infty} c_n x^n = c_0 + 2c_1 + 4c_2 + \cdots + 2^n c_n + \cdots$.

If for an x , the series $\sum_{n=0}^{\infty} c_n x^n$ converges we can use the limit of partial sequences to define a function f at x .

- We will see that a power series defines a function $f(x)$ on a certain interval where it converges.

When we fix a value for x , say $x = 1$, the power series

$$\sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 + c_2 + \cdots + c_n + \cdots$$

is an infinite series whose convergence or divergence can be investigated.

For $x = 2$, the power series $\sum_{n=0}^{\infty} c_n x^n = c_0 + 2c_1 + 4c_2 + \cdots + 2^n c_n + \cdots$.

If for an x , the series $\sum_{n=0}^{\infty} c_n x^n$ converges we can use the limit of partial sequences to define a function f at x .

- We will see that a power series defines a function $f(x)$ on a certain interval where it converges.
- Finding this interval of convergence is important. Moreover, this function will be shown to be continuous and differentiable inside the interval.

Let us consider some familiar power series.

$$\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + x^3 + x^4 + \cdots + x^n + \cdots .$$

Let us consider some familiar power series.

$$\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + x^3 + x^4 + \cdots + x^n + \cdots .$$

This is the geometric series with first term 1 and common ratio x . It converges to $\frac{1}{(1-x)}$ for $|x| < 1$. We express this fact by writing

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + x^4 + \cdots + x^n + \cdots , -1 < x < 1.$$

Let us consider some familiar power series.

$$\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + x^3 + x^4 + \cdots + x^n + \cdots .$$

This is the geometric series with first term 1 and common ratio x . It converges to $\frac{1}{(1-x)}$ for $|x| < 1$. We express this fact by writing

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + x^4 + \cdots + x^n + \cdots , -1 < x < 1.$$


We think of the partial sums of the series on the right as polynomials $P_n(x)$ that approximate the function on the left.

$$P_1(x) = 1$$

$$P_2(x) = 1 + x$$

$$P_3(x) = 1 + x + x^2$$

$$P_n(x) = 1 + x + x^2 + \cdots + x^n$$



For values of x near zero, we need take only a few terms of the series to get a good approximation. The approximations do not apply when $|x| > 1$.

Consider the power series

$$1 - \frac{1}{2}(x - 2) + \frac{1}{4}(x - 2)^2 + \cdots + \left(-\frac{1}{2}\right)^n(x - 2)^n + \cdots$$

This is a geometric series with first term 1 and ratio $r = -\frac{(x-2)}{2}$. The series converges for

$$\left| \frac{x - 2}{2} \right| < 1$$

which simplifies to $0 < x < 4$.

Consider the power series

$$1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 + \cdots + \left(-\frac{1}{2}\right)^n(x-2)^n + \cdots$$

This is a geometric series with first term 1 and ratio $r = -\frac{(x-2)}{2}$. The series converges for

$$\left| \frac{x-2}{2} \right| < 1$$

which simplifies to $0 < x < 4$. The sum is $\frac{1}{1-r} = \frac{2}{x}$. Thus

$$\frac{2}{x} = 1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 + \cdots + \left(-\frac{1}{2}\right)^n(x-2)^n + \cdots, 0 < x < 4.$$

The following example illustrates how we test a power series for convergence by using the Ratio Test to see where it converges and diverges.

The following example illustrates how we test a power series for convergence by using the Ratio Test to see where it converges and diverges.

For what values of x the power series $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n}$ converge?

The following example illustrates how we test a power series for convergence by using the Ratio Test to see where it converges and diverges.

For what values of x the power series $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n}$ converge?

- $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| =$

The following example illustrates how we test a power series for convergence by using the Ratio Test to see where it converges and diverges.

For what values of x the power series $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n}$ converge?

- $$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{n+1}}{n+1} \cdot \frac{n}{x^n} \right| = \lim_{n \rightarrow \infty} \frac{n}{n+1} |x| = |x|.$$

The following example illustrates how we test a power series for convergence by using the Ratio Test to see where it converges and diverges.

For what values of x the power series $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n}$ converge?

- $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{n+1}}{n+1} \cdot \frac{n}{x^n} \right| = \lim_{n \rightarrow \infty} \frac{n}{n+1} |x| = |x|.$
- By the Ratio Test, the series converges absolutely for $|x| < 1$ and diverges for $|x| > 1$.

The following example illustrates how we test a power series for convergence by using the Ratio Test to see where it converges and diverges.

For what values of x the power series $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n}$ converge?

- $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{n+1}}{n+1} \cdot \frac{n}{x^n} \right| = \lim_{n \rightarrow \infty} \frac{n}{n+1} |x| = |x|.$
- By the Ratio Test, the series converges absolutely for $|x| < 1$ and diverges for $|x| > 1$.
- At $x = 1$,

The following example illustrates how we test a power series for convergence by using the Ratio Test to see where it converges and diverges.

For what values of x the power series $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n}$ converge?

- $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{n+1}}{n+1} \cdot \frac{n}{x^n} \right| = \lim_{n \rightarrow \infty} \frac{n}{n+1} |x| = |x|.$
- By the Ratio Test, the series converges absolutely for $|x| < 1$ and diverges for $|x| > 1$.
- At $x = 1$, we get the alternating harmonic series $1 - \frac{1}{2} + \frac{1}{3} - \dots$ which converges.

The following example illustrates how we test a power series for convergence by using the Ratio Test to see where it converges and diverges.

For what values of x the power series $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n}$ converge?

- $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{n+1}}{n+1} \cdot \frac{n}{x^n} \right| = \lim_{n \rightarrow \infty} \frac{n}{n+1} |x| = |x|.$
- By the Ratio Test, the series converges absolutely for $|x| < 1$ and diverges for $|x| > 1$.
- At $x = 1$, we get the alternating harmonic series $1 - \frac{1}{2} + \frac{1}{3} - \dots$ which converges.
- At $x = -1$, we get the harmonic series $1 + \frac{1}{2} + \frac{1}{3} + \dots$ which diverges.

The following example illustrates how we test a power series for convergence by using the Ratio Test to see where it converges and diverges.

For what values of x the power series $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n}$ converge?

- $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{n+1}}{n+1} \cdot \frac{n}{x^n} \right| = \lim_{n \rightarrow \infty} \frac{n}{n+1} |x| = |x|.$
- By the Ratio Test, the series converges absolutely for $|x| < 1$ and diverges for $|x| > 1$.
- At $x = 1$, we get the alternating harmonic series $1 - \frac{1}{2} + \frac{1}{3} - \dots$ which converges.
- At $x = -1$, we get the harmonic series $1 + \frac{1}{2} + \frac{1}{3} + \dots$ which diverges. Thus $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n}$ converges for $-1 < x \leq 1$ and diverges elsewhere.

For what values of x the power series $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{2n-1}}{2n-1}$ converges?

For what values of x the power series $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{2n-1}}{2n-1}$ converges?

- By the Ratio Test, the series converges for $x^2 < 1$ and diverges for $x^2 > 1$.

For what values of x the power series $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{2n-1}}{2n-1}$ converges?

- By the Ratio Test, the series converges for $x^2 < 1$ and diverges for $x^2 > 1$.
- At $x = \pm 1$, the alternating series converges.

For what values of x the power series $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{2n-1}}{2n-1}$ converges?

- By the Ratio Test, the series converges for $x^2 < 1$ and diverges for $x^2 > 1$.
- At $x = \pm 1$, the alternating series converges.
- Thus the series converges for $-1 \leq x \leq 1$ and diverges elsewhere.

For what values of x the power series $\sum_{n=1}^{\infty} \frac{x^n}{n!}$ converges?

For what values of x the power series $\sum_{n=1}^{\infty} \frac{x^n}{n!}$ converges?

- By ratio test, $\left| \frac{a_{n+1}}{a_n} \right| = \left| \frac{x^{n+1}}{(n+1)!} \cdot \frac{n!}{x^n} \right| = \frac{|x|}{n+1} \rightarrow 0$ for every x .

For what values of x the power series $\sum_{n=1}^{\infty} \frac{x^n}{n!}$ converges?

- By ratio test, $\left| \frac{a_{n+1}}{a_n} \right| = \left| \frac{x^{n+1}}{(n+1)!} \cdot \frac{n!}{x^n} \right| = \frac{|x|}{n+1} \rightarrow 0$ for every x .
- The series converges absolutely and hence converges for all $x \in \mathbb{R}$.

For what values of x the power series $\sum_{n=1}^{\infty} n!x^n$ converges?

For what values of x the power series $\sum_{n=1}^{\infty} n!x^n$ converges?

By ratio test, $\left| \frac{a_{n+1}}{a_n} \right| = \left| \frac{(n+1)!x^{n+1}}{n!x^n} \right| = (n+1)|x| \rightarrow \infty$ except for $x = 0$.

For what values of x the power series $\sum_{n=1}^{\infty} n!x^n$ converges?

By ratio test, $\left| \frac{a_{n+1}}{a_n} \right| = \left| \frac{(n+1)!x^{n+1}}{n!x^n} \right| = (n+1)|x| \rightarrow \infty$ except for $x = 0$. The series diverges for all values of x except at $x = 0$.

The Convergence Theorem for Power Series

Theorem

If the power series

$$\sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + \cdots + a_n x^n + \cdots$$

converges at $x = c$,

The Convergence Theorem for Power Series

Theorem

If the power series

$$\sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + \cdots + a_n x^n + \cdots$$

converges at $x = c$, then it converges absolutely for all x with $|x| < |c|$.

The Convergence Theorem for Power Series

Theorem

If the power series

$$\sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + \cdots + a_n x^n + \cdots$$

converges at $x = c$, then it converges absolutely for all x with $|x| < |c|$. If the series diverges at $x = d$, then it diverges for all x with $|x| > |d|$.

- $\sum_{n=0}^{\infty} (-1)^{n-1} \frac{x^n}{n}$: Interval of convergence is $x \in (-1, 1]$.

- $\sum_{n=0}^{\infty} (-1)^{n-1} \frac{x^n}{n}$: Interval of convergence is $x \in (-1, 1]$.
- $\sum_{n=0}^{\infty} \frac{x^n}{n!}$: Interval of convergence is $x \in (-\infty, \infty)$.

- $\sum_{n=0}^{\infty} (-1)^{n-1} \frac{x^n}{n}$: Interval of convergence is $x \in (-1, 1]$.
- $\sum_{n=0}^{\infty} \frac{x^n}{n!}$: Interval of convergence is $x \in (-\infty, \infty)$.

Definition

The maximum value of R such that the series $\sum_{n=0}^{\infty} a_n x^n$ converges for $x \in (-R, R)$ is called Radius of Convergence.

- $\sum_{n=0}^{\infty} (-1)^{n-1} \frac{x^n}{n}$: Interval of convergence is $x \in (-1, 1]$.
- $\sum_{n=0}^{\infty} \frac{x^n}{n!}$: Interval of convergence is $x \in (-\infty, \infty)$.

Definition


The maximum value of R such that the series $\sum_{n=0}^{\infty} a_n x^n$ converges for $x \in (-R, R)$ is called Radius of Convergence.

- At $x = \pm R$, the series may or may not converge.
- When $x \in (-\infty, -R) \cup (R, \infty)$ the series diverges.

Radius of convergence

For series of the form $\sum_{n=0}^{\infty} a_n(x - a)^n$, we can replace $x - a$ by y and apply the results to the series $\sum_{n=0}^{\infty} a_n y^n$. The maximum range R for which the series $\sum_{n=0}^{\infty} a_n(x - a)^n$ converges for $a - R < x < a + R$ is called the radius of convergence of the power series, and the interval of radius R centered at $x = a$ is called the interval of convergence.

Possibilities of convergence



conv.png

- The interval of convergence may be open, closed, or half-open, depending on the particular series. Outside the interval of convergence, the power series will diverge.

- The interval of convergence may be open, closed, or half-open, depending on the particular series. Outside the interval of convergence, the power series will diverge.
- At points x with $|x - a| < R$, the series converges absolutely.

- The interval of convergence may be open, closed, or half-open, depending on the particular series. Outside the interval of convergence, the power series will diverge.
- At points x with $|x - a| < R$, the series converges absolutely.
- If the series converges for all values of x , we say its radius of convergence is infinite.

- The interval of convergence may be open, closed, or half-open, depending on the particular series. Outside the interval of convergence, the power series will diverge.
- At points x with $|x - a| < R$, the series converges absolutely.
- If the series converges for all values of x , we say its radius of convergence is infinite.
- If it converges only at $x = a$, we say its radius of convergence is zero.

- The interval of convergence may be open, closed, or half-open, depending on the particular series. Outside the interval of convergence, the power series will diverge.
- At points x with $|x - a| < R$, the series converges absolutely.
- If the series converges for all values of x , we say its radius of convergence is infinite.
- If it converges only at $x = a$, we say its radius of convergence is zero.

How to calculate R ?

- Use ratio test or root test to find R such that in $|x - a| < R$, the power series $\sum_{n=0}^{\infty} a_n(x - a)^n$, converges.

How to calculate R ?

- Use ratio test or root test to find R such that in $|x - a| < R$, the power series $\sum_{n=0}^{\infty} a_n(x - a)^n$, converges.
- Check for the convergence at $|x - a| = R$ to conclude if R also a part of interval of convergence.

How to calculate R ?

- Use ratio test or root test to find R such that in $|x - a| < R$, the power series $\sum_{n=0}^{\infty} a_n(x - a)^n$, converges.
- Check for the convergence at $|x - a| = R$ to conclude if R also a part of interval of convergence.

(This is already did for $\sum_{n=0}^{\infty} (-1)^{n-1} \frac{x^n}{n}$ and $\sum_{n=0}^{\infty} \frac{x^n}{n!}$)

On the **intersection of their intervals of convergence**, two power series can be added and subtracted term by term just like series of constants.

On the **intersection of their intervals of convergence**, two power series can be added and subtracted term by term just like series of constants. They can be multiplied just as we multiply polynomials as follows:

Theorem

If $A(x) = \sum_{n=0}^{\infty} a_n x^n$ and $B(x) = \sum_{n=0}^{\infty} b_n x^n$ converge absolutely for $|x| < R$,

and $c_n = a_0 b_n + a_1 b_{n-1} + a_2 b_{n-2} + \cdots + a_n b_0 = \sum_{k=0}^n a_k b_{n-k}$, then

$\sum_{n=0}^{\infty} c_n x^n$ converges absolutely to $A(x)B(x)$ for $|x| < R$.

$$\left(\sum_{n=0}^{\infty} a_n x^n \right) \left(\sum_{n=0}^{\infty} b_n x^n \right) = \sum_{n=0}^{\infty} c_n x^n$$

Theorem

If $\sum_{n=0}^{\infty} a_n x^n$ converges absolutely for $|x| < R$ and f is a continuous function, then $\sum_{n=0}^{\infty} a_n f(x)^n$ converges absolutely on the set of points x where $|f(x)| < R$.

Theorem

If $\sum_{n=0}^{\infty} a_n x^n$ converges absolutely for $|x| < R$ and f is a continuous

function, then $\sum_{n=0}^{\infty} a_n f(x)^n$ converges absolutely on the set of points x where $|f(x)| < R$.

Remark

Note that $\sum_{n=0}^{\infty} a_n f(x)^n$ may not be a power series, it will depend upon the choice of f . But using comparison test we can conclude about absolute convergence of $\sum_{n=0}^{\infty} a_n f(x)^n$, provided $|f(x)| < R$.

1. Find the interval of convergence for $\sum_{n=0}^{\infty} (4x^2)^n$. To which function this series will converge in this interval?

1. Find the interval of convergence for $\sum_{n=0}^{\infty} (4x^2)^n$. To which function this series will converge in this interval?

Since $\sum_{n=0}^{\infty} x^n$ converges absolutely to the function $\frac{1}{1-x}$ for $|x| < 1$,

1. Find the interval of convergence for $\sum_{n=0}^{\infty} (4x^2)^n$. To which function this series will converge in this interval?

Since $\sum_{n=0}^{\infty} x^n$ converges absolutely to the function $\frac{1}{1-x}$ for $|x| < 1$, it follows from above theorem (with $f(x) = 4x^2$) that

1. Find the interval of convergence for $\sum_{n=0}^{\infty} (4x^2)^n$. To which function this series will converge in this interval?

Since $\sum_{n=0}^{\infty} x^n$ converges absolutely to the function $\frac{1}{1-x}$ for $|x| < 1$, it follows from above theorem (with $f(x) = 4x^2$) that

$\sum_{n=0}^{\infty} (4x^2)^n$ converges absolutely to $\frac{1}{1-4x^2}$ when x satisfies $|4x^2| < 1$ or equivalently when $|x| < \frac{1}{2}$.

Theorem

If $\sum_{n=0}^{\infty} a_n(x-a)^n$ has radius of convergence $R > 0$, it defines a function

$f(x) = \sum_{n=0}^{\infty} a_n(x-a)^n$ on the interval $a - R < x < a + R$. This function f has derivatives of all orders inside the interval, and the derivatives are obtained by differentiating the original series term by term

$$f'(x) = \sum_{n=1}^{\infty} n a_n (x-a)^{n-1}$$

$$f''(x) = \sum_{n=2}^{\infty} n(n-1) a_n (x-a)^{n-2}$$

and so on. Each of these derived series converges at every point of the interval $a - R < x < a + R$.

Find series for $f'(x)$ and $f''(x)$ if

$$f(x) = \frac{1}{1-x} = 1 + x + x^2 + x^3 + \cdots = \sum_{n=1}^{\infty} x^n.$$

We differentiate the power series on the right term by term:

$$f'(x) = \frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + \cdots = \sum_{n=1}^{\infty} nx^{n-1}, -1 < x < 1$$

$$f''(x) = \frac{2}{(1-x)^3} = 2 + 6x + 12x^2 \cdots = \sum_{n=1}^{\infty} n(n-1)x^{n-2}, -1 < x < 1$$

Remark

Term-by-term differentiation might not work for other kinds of series.

Remark

Term-by-term differentiation might not work for other kinds of series.

For example, the trigonometric series $\sum_{n=1}^{\infty} \frac{\sin(n!x)}{n^2}$ converges for all x . But if we differentiate term by term we get the series $\sum_{n=1}^{\infty} \frac{n! \cos(n!x)}{n^2}$ which diverges for all x . Note that this is not a power series since it is not a sum of positive integer powers of x .

Theorem

Suppose that $\sum_{n=0}^{\infty} a_n(x-a)^n$ converges for $a-R < x < a+R$. Then

$$\sum_{n=0}^{\infty} a_n \frac{(x-a)^{n+1}}{n+1} \text{ converges for } a-R < x < a+R$$

and

$$\int f(x)dx = \sum_{n=0}^{\infty} a_n \frac{(x-a)^{n+1}}{n+1} + C \text{ for } a-R < x < a+R.$$

Identify the function

$$f(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{2n+1} = x - \frac{x^3}{3} + \frac{x^5}{5} - \cdots, -1 \leq x \leq 1.$$

Identify the function

$$f(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{2n+1} = x - \frac{x^3}{3} + \frac{x^5}{5} - \cdots, -1 \leq x \leq 1.$$

Differentiating the original series term by term, we get

$$f'(x) = 1 - x^2 + x^4 - \cdots, -1 < x < 1.$$

Identify the function

$$f(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{2n+1} = x - \frac{x^3}{3} + \frac{x^5}{5} - \cdots, -1 \leq x \leq 1.$$

Differentiating the original series term by term, we get

$$f'(x) = 1 - x^2 + x^4 - \cdots, -1 < x < 1.$$

This is a geometric series with first term 1 and ratio $-x^2$,

Identify the function

$$f(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{2n+1} = x - \frac{x^3}{3} + \frac{x^5}{5} - \cdots, -1 \leq x \leq 1.$$

Differentiating the original series term by term, we get

$$f'(x) = 1 - x^2 + x^4 - \cdots, -1 < x < 1.$$

This is a geometric series with first term 1 and ratio $-x^2$, so $f'(x) = \frac{1}{1+x^2}$.

We can now integrate $f'(x) = \frac{1}{1+x^2}$ to get

$$\int f'(x) dx = \int \frac{dx}{1+x^2} = \tan^{-1} x + C.$$

The series for $f(x)$ is 0 when $x = 0$, so $C = 0$. Thus

$$f(x) = x - \frac{x^3}{3} + \frac{x^5}{5} - \cdots = \tan^{-1} x, -1 < x < 1.$$

Note that the original series $x - \frac{x^3}{3} + \frac{x^5}{5} - \dots$, converges at both endpoints of the original interval of convergence, but our theorem can only guarantee the convergence of the differentiated series inside the interval.

The series

$$\frac{1}{1+t} = 1 - t + t^2 - t^3 + \dots$$

converges on the open interval $-1 < t < 1$.

The series

$$\frac{1}{1+t} = 1 - t + t^2 - t^3 + \dots$$

converges on the open interval $-1 < t < 1$.

Therefore,

$$\ln(1+x) = \int_0^x \frac{1}{1+t} dt = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots, -1 < x < 1.$$

We have seen that within its interval of convergence I , the sum of a power series is a continuous function with derivatives of all orders. Now we ask the reverse question.

We have seen that within its interval of convergence I , the sum of a power series is a continuous function with derivatives of all orders. Now we ask the reverse question.

Remark

- *If a function $f(x)$ has derivatives of all orders on an interval, can it be expressed as a power series on at least part of that interval?*

We have seen that within its interval of convergence I , the sum of a power series is a continuous function with derivatives of all orders. Now we ask the reverse question.

Remark

- *If a function $f(x)$ has derivatives of all orders on an interval, can it be expressed as a power series on at least part of that interval?*
- *And if it can, what are its coefficients?*

We have seen that within its interval of convergence I , the sum of a power series is a continuous function with derivatives of all orders. Now we ask the reverse question.

Remark

- *If a function $f(x)$ has derivatives of all orders on an interval, can it be expressed as a power series on at least part of that interval?*
- *And if it can, what are its coefficients?*

If we can find power series representation of a function, they provide useful polynomial approximations of the original functions. Because approximation by polynomials is extremely useful to both mathematicians and scientists, we are interested to see when a function can have power series representation.

