

# Double and Triple Integrals

Devika S

Department of Mathematics  
BITS Pilani, K K Birla Goa Campus

November 6, 2024



# Double Integrals - Polar Form

## Theorem

Suppose that  $f(r, \theta)$  is continuous on the region  $R = \{(r, \theta) : g_1(\theta) \leq r \leq g_2(\theta), \alpha \leq \theta \leq \beta\}$ . Then

# Double Integrals - Polar Form

## Theorem

Suppose that  $f(r, \theta)$  is continuous on the region  $R = \{(r, \theta) : g_1(\theta) \leq r \leq g_2(\theta), \alpha \leq \theta \leq \beta\}$ . Then

$$\iint_R f(r, \theta) dA = \int_{\theta=\alpha}^{\theta=\beta} \int_{r=g_1(\theta)}^{r=g_2(\theta)} f(r, \theta) r \, dr \, d\theta.$$

# Double Integrals - Polar Form

## Theorem

Suppose that  $f(r, \theta)$  is continuous on the region  $R = \{(r, \theta) : g_1(\theta) \leq r \leq g_2(\theta), \alpha \leq \theta \leq \beta\}$ . Then

$$\iint_R f(r, \theta) dA = \int_{\theta=\alpha}^{\theta=\beta} \int_{r=g_1(\theta)}^{r=g_2(\theta)} f(r, \theta) r \, dr \, d\theta.$$

## Theorem

Suppose that  $f(r, \theta)$  is continuous on the region  $R = \{(r, \theta) : a \leq r \leq b, h_1(r) \leq \theta \leq h_2(r)\}$ . Then

# Double Integrals - Polar Form

## Theorem

Suppose that  $f(r, \theta)$  is continuous on the region  $R = \{(r, \theta) : g_1(\theta) \leq r \leq g_2(\theta), \alpha \leq \theta \leq \beta\}$ . Then

$$\iint_R f(r, \theta) dA = \int_{\theta=\alpha}^{\theta=\beta} \int_{r=g_1(\theta)}^{r=g_2(\theta)} f(r, \theta) r \, dr \, d\theta.$$

## Theorem

Suppose that  $f(r, \theta)$  is continuous on the region  $R = \{(r, \theta) : a \leq r \leq b, h_1(r) \leq \theta \leq h_2(r)\}$ . Then

$$\iint_R f(r, \theta) dA = \int_{r=a}^{r=b} \int_{\theta=h_1(r)}^{\theta=h_2(r)} f(r, \theta) r \, d\theta \, dr.$$

# Examples - Double Integrals

Example 1. Find the integral:

$$\int_0^2 \int_0^{\sqrt{1-(x-1)^2}} \frac{x+y}{x^2+y^2} dy dx.$$

# Examples - Double Integrals

Example 1. Find the integral:

$$\int_0^2 \int_0^{\sqrt{1-(x-1)^2}} \frac{x+y}{x^2+y^2} dy dx.$$

**Solution:** The integral turns out to be:

$$\int_0^{\frac{\pi}{2}} \int_0^{2\cos\theta} (\cos\theta + \sin\theta) dr d\theta = \frac{\pi}{2} + 1.$$

# Examples - Double Integrals

Example 1. Find the integral:

$$\int_0^2 \int_0^{\sqrt{1-(x-1)^2}} \frac{x+y}{x^2+y^2} dy dx.$$

**Solution:** The integral turns out to be:

$$\int_0^{\frac{\pi}{2}} \int_0^{2\cos\theta} (\cos\theta + \sin\theta) dr d\theta = \frac{\pi}{2} + 1.$$

Example 2. Find the integral:

$$\int_0^{\ln 2} \int_0^{\sqrt{(\ln 2)^2 - y^2}} e^{\sqrt{x^2 + y^2}} dx dy.$$



# Examples - Double Integrals

Example 1. Find the integral:

$$\int_0^2 \int_0^{\sqrt{1-(x-1)^2}} \frac{x+y}{x^2+y^2} dy dx.$$

**Solution:** The integral turns out to be:

$$\int_0^{\frac{\pi}{2}} \int_0^{2\cos\theta} (\cos\theta + \sin\theta) dr d\theta = \frac{\pi}{2} + 1.$$

Example 2. Find the integral:

$$\int_0^{\ln 2} \int_0^{\sqrt{(\ln 2)^2 - y^2}} e^{\sqrt{x^2 + y^2}} dx dy.$$

**Solution:**

$$\int_0^{\frac{\pi}{2}} \int_0^{\ln 2} r e^r dr d\theta = (2\ln 2 - 1) \frac{\pi}{2}.$$

## Examples

**Example 3.** Find the integral by converting to a double integral

$$\int_0^2 (\tan^{-1} \pi x - \tan^{-1} x) dx.$$

## Examples

**Example 3.** Find the integral by converting to a double integral

$$\int_0^2 (\tan^{-1} \pi x - \tan^{-1} x) dx.$$

**Solution:** First let us write the integrand as an integral  $\int_x^{\pi x} \frac{1}{1+y^2} dy$ .

## Examples

**Example 3.** Find the integral by converting to a double integral

$$\int_0^2 (\tan^{-1} \pi x - \tan^{-1} x) dx.$$

**Solution:** First let us write the integrand as an integral  $\int_x^{\pi x} \frac{1}{1+y^2} dy$ .

Then we have to integrate

$$\int_0^2 \int_x^{\pi x} \frac{1}{1+y^2} dy dx.$$

## Examples

**Example 3.** Find the integral by converting to a double integral

$$\int_0^2 (\tan^{-1} \pi x - \tan^{-1} x) dx.$$

**Solution:** First let us write the integrand as an integral  $\int_x^{\pi x} \frac{1}{1+y^2} dy$ .

Then we have to integrate

$$\int_0^2 \int_x^{\pi x} \frac{1}{1+y^2} dy dx.$$

We sketch the region. Note that we have to write the region as disjoint union of the region  $y \leq 2$  and the region  $y \geq 2$ .

## Examples

**Example 3.** Find the integral by converting to a double integral

$$\int_0^2 (\tan^{-1} \pi x - \tan^{-1} x) dx.$$

**Solution:** First let us write the integrand as an integral  $\int_x^{\pi x} \frac{1}{1+y^2} dy$ .

Then we have to integrate

$$\int_0^2 \int_x^{\pi x} \frac{1}{1+y^2} dy dx.$$

We sketch the region. Note that we have to write the region as disjoint union of the region  $y \leq 2$  and the region  $y \geq 2$ . The final integral is

$$\int_0^2 \int_{\frac{y}{\pi}}^y \frac{1}{1+y^2} dx dy + \int_2^{2\pi} \int_{\frac{y}{\pi}}^2 \frac{1}{1+y^2} dx dy,$$

which can be computed easily.

# Examples

① Integrate the following by changing to the polar coordinates:

- $\int_0^1 \int_x^1 \frac{y}{x^2 + y^2} dy dx.$
- $\int_0^1 \int_{-y/3}^{y/3} \frac{y}{\sqrt{x^2 + y^2}} dx dy.$

② Find the area of the region that lies inside the cardioid  $r = 1 + \cos \theta$  and outside the circle  $r = 1$ .  $((8 + \pi)/4)$

③ Find the volume of the solid cut from the first octant by the surface  $z = 4 - x^2 - y$ .  $(128/15)$

④ Find the volume of the sphere of radius  $a$  by double integration.  
 $(\frac{4}{3}\pi a^3 - \text{compute the volume of the portion in the first octant and multiply the answer by } 8)$

⑤ Find the average height of the hemispherical surface  $z = \sqrt{a^2 - x^2 - y^2}$  above the disk  $x^2 + y^2 \leq a^2$  in the  $xy$ -plane.  
 $(2a/3)$

# Triple Integrals

A **triple integral** or a **space integral** is integral of a function of three variables over a domain given in **three dimensional region/space**.



# Triple Integrals

A **triple integral** or a **space integral** is integral of a function of three variables over a domain given in **three dimensional region/space**.

If  $F(x, y, z)$  is a function defined on a closed bounded region  $D$  in space, such as the region occupied by a solid ball or a lump of clay, then the integral of  $F$  over  $D$  may be defined in the following way.


# Triple Integrals

A **triple integral** or a **space integral** is integral of a function of three variables over a domain given in **three dimensional region/space**.

If  $F(x, y, z)$  is a function defined on a closed bounded region  $D$  in space, such as the region occupied by a solid ball or a lump of clay, then the integral of  $F$  over  $D$  may be defined in the following way.

We **partition a rectangular box like region containing  $D$  into rectangular cells** by planes parallel to the coordinate axes.

# Triple Integrals



# Triple Integrals

We number the cells that lie completely inside  $D$  from 1 to  $n$  in some order, the  $k$ -th cell having dimensions  $\Delta x_k \times \Delta y_k \times \Delta z_k$  and volume  $\Delta V_k = \Delta x_k \Delta y_k \Delta z_k$ . We choose a point  $(x_k, y_k, z_k)$ . Define the norm of the partition  $P$  as  $\|P\| = \max\{\Delta x_k, \Delta y_k, \Delta z_k\}$ .

# Triple Integrals

We number the cells that lie completely inside  $D$  from 1 to  $n$  in some order, the  $k$ -th cell having dimensions  $\Delta x_k \times \Delta y_k \times \Delta z_k$  and volume  $\Delta V_k = \Delta x_k \Delta y_k \Delta z_k$ . We choose a point  $(x_k, y_k, z_k)$ . Define the norm of the partition  $P$  as  $\|P\| = \max\{\Delta x_k, \Delta y_k, \Delta z_k\}$ .

## Triple Integrals

Triple integral of  $F$  over  $D$  is defined by

$$\iiint_D F(x, y, z) dV = \lim_{\|P\| \rightarrow 0} \sum_{k=1}^n F(x_k, y_k, z_k) \Delta V_k,$$

provided the limit exists.

# Triple Integrals

## Volume

The volume of a closed, bounded region  $D$  in space is

$$V = \iiint_D dV.$$

# Triple Integrals

## Volume

The volume of a closed, bounded region  $D$  in space is

$$V = \iiint_D dV.$$

## Average

The average value of a function  $F$  over a region  $D$  in space is

$$\frac{1}{\text{Volume of } D} \iiint_D F dV.$$

# Triple Integral over Rectangular Region

## Fubini's Theorem

Suppose that  $f(x, y, z)$  is continuous on the region  $D = \{(x, y, z) : a \leq x \leq b, c \leq y \leq d, e \leq z \leq f\}$ . Then, triple integral can be written as triple iterated integral:

$$\iiint_D f(x, y, z) dV = \int_e^f \int_c^d \int_a^b f(x, y, z) dx dy dz.$$



# Triple Integral over Rectangular Region

## Fubini's Theorem

Suppose that  $f(x, y, z)$  is continuous on the region  $D = \{(x, y, z) : a \leq x \leq b, c \leq y \leq d, e \leq z \leq f\}$ . Then, triple integral can be written as triple iterated integral:

$$\iiint_D f(x, y, z) dV = \int_e^f \int_c^d \int_a^b f(x, y, z) dx dy dz.$$

Note that a triple integral can be evaluated with six different possible orders of integration by choosing the limits appropriately.

# Triple Integral over Rectangular Region

## Fubini's Theorem

Suppose that  $f(x, y, z)$  is continuous on the region  $D = \{(x, y, z) : a \leq x \leq b, c \leq y \leq d, e \leq z \leq f\}$ . Then, triple integral can be written as triple iterated integral:

$$\iiint_D f(x, y, z) dV = \int_e^f \int_c^d \int_a^b f(x, y, z) dx dy dz.$$

Note that a triple integral can be evaluated with six different possible orders of integration by choosing the limits appropriately.

**Example 1.** Evaluate  $\int_0^1 \int_2^4 \int_{-1}^5 (x + yz^2) dx dy dz$

# Triple Integral over Rectangular Region

## Fubini's Theorem

Suppose that  $f(x, y, z)$  is continuous on the region  $D = \{(x, y, z) : a \leq x \leq b, c \leq y \leq d, e \leq z \leq f\}$ . Then, triple integral can be written as triple iterated integral:

$$\iiint_D f(x, y, z) dV = \int_e^f \int_c^d \int_a^b f(x, y, z) dx dy dz.$$

Note that a triple integral can be evaluated with six different possible orders of integration by choosing the limits appropriately.

**Example 1.** Evaluate  $\int_0^1 \int_2^4 \int_{-1}^5 (x + yz^2) dx dy dz$  (36)

# Triple Integral over Rectangular Region

## Fubini's Theorem

Suppose that  $f(x, y, z)$  is continuous on the region  $D = \{(x, y, z) : a \leq x \leq b, c \leq y \leq d, e \leq z \leq f\}$ . Then, triple integral can be written as triple iterated integral:

$$\iiint_D f(x, y, z) dV = \int_e^f \int_c^d \int_a^b f(x, y, z) dx dy dz.$$

Note that a triple integral can be evaluated with six different possible orders of integration by choosing the limits appropriately.

**Example 1.** Evaluate  $\int_0^1 \int_2^4 \int_{-1}^5 (x + yz^2) dx dy dz$  (36)

**Example 2.** Evaluate  $\iiint_D x^2 yz dV$ ,  $D = [-2, 1] \times [0, 3] \times [1, 5]$

# Triple Integral over Rectangular Region

## Fubini's Theorem

Suppose that  $f(x, y, z)$  is continuous on the region  $D = \{(x, y, z) : a \leq x \leq b, c \leq y \leq d, e \leq z \leq f\}$ . Then, triple integral can be written as triple iterated integral:

$$\iiint_D f(x, y, z) dV = \int_e^f \int_c^d \int_a^b f(x, y, z) dx dy dz.$$

Note that a triple integral can be evaluated with six different possible orders of integration by choosing the limits appropriately.

**Example 1.** Evaluate  $\int_0^1 \int_2^4 \int_{-1}^5 (x + yz^2) dx dy dz$  (36)

**Example 2.** Evaluate  $\iiint_D x^2 yz dV$ ,  $D = [-2, 1] \times [0, 3] \times [1, 5]$  (162)

# Finding Limits of Integration in the Order $dzdydx$

First let us discuss how to find the limits of a triple integral:

# Finding Limits of Integration in the Order $dzdydx$

First let us discuss how to find the limits of a triple integral:

- First sketch the region  $D$  in three dimensional space along with the its shadow (vertical projection)  $R$  in the  $xy$  plane.

# Finding Limits of Integration in the Order $dzdydx$

First let us discuss how to find the limits of a triple integral:

- First sketch the region  $D$  in three dimensional space along with the its shadow (vertical projection)  $R$  in the  $xy$  plane.
- Find the  $z$  limits by imagining a line passing through  $(x, y, 0)$  and is parallel to the  $z$  axis entering and leaving the region, the  $z$  coordinates are the  $z$  limits as functions of  $x, y$ .




# Finding Limits of Integration in the Order $dzdydx$


First let us discuss how to find the limits of a triple integral:

- First sketch the region  $D$  in three dimensional space along with the its shadow (vertical projection)  $R$  in the  $xy$  plane.
- Find the  $z$  limits by imagining a line passing through  $(x, y, 0)$  and is parallel to the  $z$  axis entering and leaving the region, the  $z$  coordinates are the  $z$  limits as functions of  $x, y$ .
- Find the  $x$  and  $y$  limits for the projection in  $xy$ -plane as discussed in the double integrals in accordance of your order of integral.


# Triple Integrals - Step 1



# Triple Integrals - Step 2



## Triple Integrals - Step 3



# Fubini's Theorem

## Fubini's Theorem

Let  $F$  be a continuous function over the domain  $D$  in the space defined by  $D = \{(x, y, z) : a \leq x \leq b, g_1(x) \leq y \leq g_2(x), f_1(x, y) \leq z \leq f_2(x, y)\}$ , then

# Fubini's Theorem

## Fubini's Theorem

Let  $F$  be a continuous function over the domain  $D$  in the space defined by  $D = \{(x, y, z) : a \leq x \leq b, g_1(x) \leq y \leq g_2(x), f_1(x, y) \leq z \leq f_2(x, y)\}$ , then

$$\iiint_D F(x, y, z) dV = \int_{x=a}^{x=b} \int_{y=g_1(x)}^{y=g_2(x)} \int_{z=f_1(x,y)}^{z=f_2(x,y)} F(x, y, z) dz \, dy \, dx.$$

# Examples

**Example 1.** Evaluate the triple integral of the function  $f(x, y, z) = 5x - 3y$  over the solid tetrahedron bounded by the planes  $x = 0$ ,  $y = 0$ ,  $z = 0$ , and  $x + y + z = 1$ .

# Examples

**Example 1.** Evaluate the triple integral of the function  $f(x, y, z) = 5x - 3y$  over the solid tetrahedron bounded by the planes  $x = 0$ ,  $y = 0$ ,  $z = 0$ , and  $x + y + z = 1$ .

**Solution:** Sketch the region  $D$  and its projection  $R$  onto  $xy$ -plane.



# Examples

**Example 1.** Evaluate the triple integral of the function  $f(x, y, z) = 5x - 3y$  over the solid tetrahedron bounded by the planes  $x = 0$ ,  $y = 0$ ,  $z = 0$ , and  $x + y + z = 1$ .

**Solution:** Sketch the region  $D$  and its projection  $R$  onto  $xy$ -plane. Then,

$$\iiint_D F(x, y, z) dV = \int_{x=0}^1 \int_0^{1-x} \int_0^{1-x-y} (5x - 3y) dz \, dy \, dx$$

# Examples

**Example 1.** Evaluate the triple integral of the function  $f(x, y, z) = 5x - 3y$  over the solid tetrahedron bounded by the planes  $x = 0$ ,  $y = 0$ ,  $z = 0$ , and  $x + y + z = 1$ .

**Solution:** Sketch the region  $D$  and its projection  $R$  onto  $xy$ -plane. Then,

$$\iiint_D F(x, y, z) dV = \int_{x=0}^1 \int_0^{1-x} \int_0^{1-x-y} (5x - 3y) dz \, dy \, dx = \frac{1}{12}.$$

# Examples

**Example 1.** Evaluate the triple integral of the function  $f(x, y, z) = 5x - 3y$  over the solid tetrahedron bounded by the planes  $x = 0$ ,  $y = 0$ ,  $z = 0$ , and  $x + y + z = 1$ .

**Solution:** Sketch the region  $D$  and its projection  $R$  onto  $xy$ -plane. Then,

$$\iiint_D F(x, y, z) dV = \int_{x=0}^1 \int_0^{1-x} \int_0^{1-x-y} (5x - 3y) dz \, dy \, dx = \frac{1}{12}.$$

**Example 2:** Evaluate the integral in Example 1 using  $dydx dz$ .

# Examples

**Example 3.** Find the volume of the sphere  
 $D = \{(x, y, z) : x^2 + y^2 + z^2 = 9\}$  using triple integral.

# Examples

**Example 3.** Find the volume of the sphere  $D = \{(x, y, z) : x^2 + y^2 + z^2 = 9\}$  using triple integral.

**Solution:** The volume  $V$  can be expressed as

$$V = \iiint_D dV = \int_{-3}^3 \int_{-\sqrt{9-x^2}}^{\sqrt{9-x^2}} \int_{-\sqrt{9-x^2-y^2}}^{\sqrt{9-x^2-y^2}} dz dy dx.$$

# Examples

**Example 3.** Find the volume of the sphere  $D = \{(x, y, z) : x^2 + y^2 + z^2 = 9\}$  using triple integral.

**Solution:** The volume  $V$  can be expressed as

$$V = \iiint_D dV = \int_{-3}^3 \int_{-\sqrt{9-x^2}}^{\sqrt{9-x^2}} \int_{-\sqrt{9-x^2-y^2}}^{\sqrt{9-x^2-y^2}} dz dy dx.$$

After performing the integration with respect to  $z$ , transform the resulting integral into polar coordinates. The volume is  $36\pi$  cubic units.

# Examples

**Example 4:** Let  $D$  be the region bounded by the paraboloids  $z = 8 - x^2 - y^2$  and  $z = x^2 + y^2$ . Write six different triple iterated integrals for the volume of  $D$ . Evaluate one of the integrals.

# Examples

**Example 4:** Let  $D$  be the region bounded by the paraboloids  $z = 8 - x^2 - y^2$  and  $z = x^2 + y^2$ . Write six different triple iterated integrals for the volume of  $D$ . Evaluate one of the integrals.

**Solution:** Let us first sketch the region and set up the integral.



# Examples

**Example 4:** Let  $D$  be the region bounded by the paraboloids  $z = 8 - x^2 - y^2$  and  $z = x^2 + y^2$ . Write six different triple iterated integrals for the volume of  $D$ . Evaluate one of the integrals.

**Solution:** Let us first sketch the region and set up the integral.

Case 1:  $dzdydx$

# Examples

**Example 4:** Let  $D$  be the region bounded by the paraboloids  $z = 8 - x^2 - y^2$  and  $z = x^2 + y^2$ . Write six different triple iterated integrals for the volume of  $D$ . Evaluate one of the integrals.

**Solution:** Let us first sketch the region and set up the integral.

**Case 1:**  $dzdydx$  In this case,

$$V = \int_{-2}^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} \int_{x^2+y^2}^{8-x^2-y^2} dzdydx = 16\pi.$$

# Examples

**Example 4:** Let  $D$  be the region bounded by the paraboloids  $z = 8 - x^2 - y^2$  and  $z = x^2 + y^2$ . Write six different triple iterated integrals for the volume of  $D$ . Evaluate one of the integrals.

**Solution:** Let us first sketch the region and set up the integral.

**Case 1:**  $dzdydx$  In this case,

$$V = \int_{-2}^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} \int_{x^2+y^2}^{8-x^2-y^2} dzdydx = 16\pi.$$

**Case 2:**  $dzdx dy$

# Examples

**Example 4:** Let  $D$  be the region bounded by the paraboloids  $z = 8 - x^2 - y^2$  and  $z = x^2 + y^2$ . Write six different triple iterated integrals for the volume of  $D$ . Evaluate one of the integrals.

**Solution:** Let us first sketch the region and set up the integral.

**Case 1:**  $dzdydx$  In this case,

$$V = \int_{-2}^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} \int_{x^2+y^2}^{8-x^2-y^2} dzdydx = 16\pi.$$

**Case 2:**  $dzdxdy$

$$V = \int_{-2}^2 \int_{-\sqrt{4-y^2}}^{\sqrt{4-y^2}} \int_{x^2+y^2}^{8-x^2-y^2} dzdxdy.$$

# Examples

Case 3:  $dydzdx$

# Examples

## Case 3: $dydzdx$

$$V = \int_{-2}^2 \int_{x^2}^4 \int_{-\sqrt{z-x^2}}^{\sqrt{z-x^2}} dy dz dx + \int_{-2}^2 \int_4^{8-x^2} \int_{-\sqrt{8-z-x^2}}^{\sqrt{8-z-x^2}} dy dz dx.$$

# Examples

Case 3:  $dydzdx$

$$V = \int_{-2}^2 \int_{x^2}^4 \int_{-\sqrt{z-x^2}}^{\sqrt{z-x^2}} dy dz dx + \int_{-2}^2 \int_4^{8-x^2} \int_{-\sqrt{8-z-x^2}}^{\sqrt{8-z-x^2}} dy dz dx.$$

Case 4:  $dydx dz$

# Examples

## Case 3: $dydzdx$

$$V = \int_{-2}^2 \int_{x^2}^4 \int_{-\sqrt{z-x^2}}^{\sqrt{z-x^2}} dy dz dx + \int_{-2}^2 \int_4^{8-x^2} \int_{-\sqrt{8-z-x^2}}^{\sqrt{8-z-x^2}} dy dz dx.$$

## Case 4: $dydx dz$

$$V = \int_0^4 \int_{-\sqrt{z}}^{\sqrt{z}} \int_{-\sqrt{z-x^2}}^{\sqrt{z-x^2}} dy dx dz + \int_4^8 \int_{-\sqrt{8-z}}^{\sqrt{8-z}} \int_{-\sqrt{8-z-x^2}}^{\sqrt{8-z-x^2}} dy dx dz.$$



# Examples

Case :5  $dx dz dy$

# Examples

Case :5  $dx dz dy$

$$V = \int_{-2}^2 \int_{y^2}^4 \int_{-\sqrt{z-y^2}}^{\sqrt{z-y^2}} dx dz dy + \int_{-2}^2 \int_4^{8-y^2} \int_{-\sqrt{8-z-y^2}}^{\sqrt{8-z-y^2}} dx dz dy.$$

# Examples

Case 5:  $dx dz dy$

$$V = \int_{-2}^2 \int_{y^2}^4 \int_{-\sqrt{z-y^2}}^{\sqrt{z-y^2}} dx dz dy + \int_{-2}^2 \int_4^{8-y^2} \int_{-\sqrt{8-z-y^2}}^{\sqrt{8-z-y^2}} dx dz dy.$$

Case 6:  $dx dy dz$

# Examples

Case 5:  $dx dz dy$

$$V = \int_{-2}^2 \int_{y^2}^4 \int_{-\sqrt{z-y^2}}^{\sqrt{z-y^2}} dx dz dy + \int_{-2}^2 \int_4^{8-y^2} \int_{-\sqrt{8-z-y^2}}^{\sqrt{8-z-y^2}} dx dz dy.$$

Case 6:  $dx dy dz$

$$V = \int_0^4 \int_{-\sqrt{z}}^{\sqrt{z}} \int_{-\sqrt{z-y^2}}^{\sqrt{z-y^2}} dx dy dz + \int_4^8 \int_{-\sqrt{8-z}}^{\sqrt{8-z}} \int_{-\sqrt{8-z-y^2}}^{\sqrt{8-z-y^2}} dx dy dz.$$

# Examples

- Find the average value of  $F(x, y, z) = x^2 + 9$  over the cube in the first octant bounded by the coordinate planes and the planes  $x = 2$ ,  $y = 2$  and  $z = 2$ . (31/3)
- Solve for  $a$ :

$$\int_0^1 \int_0^{4-a-x^2} \int_a^{4-x^2-y} dz dy dx = \frac{4}{15}. \quad (a = 13/3, 3)$$

Solved examples from Thomas' Calculus (please review the figures for guidance on how to sketch the region.)

- Find the volume of the region  $D$  enclosed by the surface  $z = x^2 + 3y^2$  and  $z = 8 - x^2 - y^2$ .
- Set up the limits of integration for evaluating the triple integral of a function  $F(x, y, z)$  over the tetrahedron  $D$  with vertices  $(0, 0, 0)$ ,  $(1, 1, 0)$ ,  $(0, 1, 0)$ , and  $(0, 1, 1)$ . Use the order of integration  $dy dz dx$  and  $dz dy dx$ .