

Extreme Values & Saddle Points

Devika S

Department of Mathematics
BITS Pilani, K K Birla Goa Campus

October 26, 2024



Extreme Values

Let $f(x, y)$ be defined on a region R containing the point (a, b) .

- ① $f(a, b)$ is a **local maximum** value of f if $f(a, b) \geq f(x, y)$ for all domain points (x, y) in an open disk centered at (a, b) .
- ② $f(a, b)$ is a **local minimum** value of f if $f(a, b) \leq f(x, y)$ for all domain points (x, y) in an open disk centered at (a, b) .
- ③ An **interior point** of the domain of a function f is called a **critical point** of the function if **either both f_x, f_y vanish or at least one of f_x and f_y does not exist** at the point.
- ④ A differentiable function $f(x, y)$ has a **saddle point** at a critical point (a, b) if in **every open disk** centered at (a, b) there are domain points (x, y) where $f(x, y) > f(a, b)$ and domain points (x, y) where $f(x, y) < f(a, b)$.

First Derivative Test for Local Extreme Values

Theorem (First Derivative Test for Local Extreme Values)

If $f(x, y)$ has a *local maximum or minimum value at an interior point* (a, b) of its domain and if the *first partial derivatives exist there*, then

$$f_x(a, b) = 0 \text{ and } f_y(a, b) = 0.$$

First Derivative Test for Local Extreme Values

Theorem (First Derivative Test for Local Extreme Values)

If $f(x, y)$ has a *local maximum or minimum value at an interior point* (a, b) of its domain and if the *first partial derivatives exist there*, then

$$f_x(a, b) = 0 \text{ and } f_y(a, b) = 0.$$

The **extreme values** of $f(x, y)$ can occur only at

- **boundary points** of the domain of f
- **critical points** (interior points where $f_x = f_y = 0$ or points where f_x or f_y fails to exist).

First Derivative Test for Local Extreme Values

Theorem (First Derivative Test for Local Extreme Values)

If $f(x, y)$ has a *local maximum or minimum value at an interior point* (a, b) of its domain and if the *first partial derivatives exist there*, then

$$f_x(a, b) = 0 \text{ and } f_y(a, b) = 0.$$

The **extreme values** of $f(x, y)$ can occur only at

- **boundary points** of the domain of f
- **critical points** (interior points where $f_x = f_y = 0$ or points where f_x or f_y fails to exist).

Example 1. Find local extreme values of $f(x, y) = x^2 + y^2$.

First Derivative Test for Local Extreme Values

Theorem (First Derivative Test for Local Extreme Values)

If $f(x, y)$ has a *local maximum or minimum value at an interior point* (a, b) of its domain and if the *first partial derivatives exist there*, then

$$f_x(a, b) = 0 \text{ and } f_y(a, b) = 0.$$

The **extreme values** of $f(x, y)$ can occur only at

- **boundary points** of the domain of f
- **critical points** (interior points where $f_x = f_y = 0$ or points where f_x or f_y fails to exist).

Example 1. Find local extreme values of $f(x, y) = x^2 + y^2$.

f is defined and differentiable for all x and y and its domain has no boundary points. The extreme values can occur only at the points where f_x and f_y are simultaneously zero. Solving $2x = 0$ and $2y = 0$, $(0, 0)$ is the only point **where f may take on an extreme value.** **CHECK!**

Examples

Example 2. Find the local extreme values of $f(x, y) = y^2 - x^2$.

Examples

Example 2. Find the local extreme values of $f(x, y) = y^2 - x^2$.

Solution: Since the domain of f is entire plane (so there are no boundary points) and the partial derivatives exist everywhere, the local extreme values can occur only where the first derivatives vanish.

Examples

Example 2. Find the local extreme values of $f(x, y) = y^2 - x^2$.

Solution: Since the domain of f is entire plane (so there are no boundary points) and the partial derivatives exist everywhere, the local extreme values can occur only where the first derivatives vanish.

Both the partial derivatives vanish only at the origin.

Examples

Example 2. Find the local extreme values of $f(x, y) = y^2 - x^2$.

Solution: Since the domain of f is entire plane (so there are no boundary points) and the partial derivatives exist everywhere, the local extreme values can occur only where the first derivatives vanish.

Both the partial derivatives vanish only at the origin.

Hence the only critical point is the origin. But at this point we see that along y axis,

$$f(0, y) = y^2 > 0.$$

$$\text{Along } x \text{ axis, } f(x, 0) = -x^2 < 0.$$

Hence, the function has a saddle point at the origin and no local extreme values.

Examples

Example 2. Find the local extreme values of $f(x, y) = y^2 - x^2$.

Solution: Since the domain of f is entire plane (so there are no boundary points) and the partial derivatives exist everywhere, the local extreme values can occur only where the first derivatives vanish.


Both the partial derivatives vanish only at the origin.

Hence the only critical point is the origin. But at this point we see that along y axis,

$$f(0, y) = y^2 > 0.$$

$$\text{Along } x \text{ axis, } f(x, 0) = -x^2 < 0.$$

Hence, the function has a saddle point at the origin and no local extreme values.



Examples

Example 3. Find the critical points of $f(x, y) = x^2 + 2xy$.

Examples

Example 3. Find the critical points of $f(x, y) = x^2 + 2xy$.

Solution: The partial derivatives are $f_x = 2x + 2y$ and $f_y = 2x$.

Examples


Example 3. Find the critical points of $f(x, y) = x^2 + 2xy$.

Solution: The partial derivatives are $f_x = 2x + 2y$ and $f_y = 2x$. Solving $2x + 2y = 0$ and $2x = 0$, we get the point $(0, 0)$.

Examples

Example 3. Find the critical points of $f(x, y) = x^2 + 2xy$.

Solution: The partial derivatives are $f_x = 2x + 2y$ and $f_y = 2x$. Solving $2x + 2y = 0$ and $2x = 0$, we get the point $(0, 0)$.



Examples

Example 4. Find the critical points of $f(x, y) = y \sin x$.

Examples

Example 4. Find the critical points of $f(x, y) = y \sin x$.

Solution: First we find the partial derivatives.

$$f_x = y \cos x, \quad f_y = \sin x.$$

Examples

Example 4. Find the critical points of $f(x, y) = y \sin x$.

Solution: First we find the partial derivatives.

$$f_x = y \cos x, \quad f_y = \sin x.$$

Find the solution of the system: $y \cos x = 0, \sin x = 0$. We get the points $(n\pi, 0)$ for integers n .


Examples

Example 4. Find the critical points of $f(x, y) = y \sin x$.

Solution: First we find the partial derivatives.

$$f_x = y \cos x, \quad f_y = \sin x.$$

Find the solution of the system: $y \cos x = 0, \sin x = 0$. We get the points $(n\pi, 0)$ for integers n .



Second Derivative Test

Note that $f_x = f_y = 0$ at an interior point (a, b) of R does not guarantee f has a local extreme value.

Theorem (Second Derivative Test for Local Extreme Values)

Suppose that $f(x, y)$ and its first and second partial derivatives are *continuous* throughout a disk centered at (a, b) and that $f_x(a, b) = f_y(a, b) = 0$. Then

- 1 f has a *local maximum* at (a, b) if $f_{xx} < 0$ and $f_{xx}f_{yy} - f_{xy}^2 > 0$ at (a, b) .
- 2 f has a *local minimum* at (a, b) if $f_{xx} > 0$ and $f_{xx}f_{yy} - f_{xy}^2 > 0$ at (a, b) .
- 3 f has a *saddle point* at (a, b) if $f_{xx}f_{yy} - f_{xy}^2 < 0$ at (a, b) .
- 4 the test is *inconclusive* at (a, b) if $f_{xx}f_{yy} - f_{xy}^2 = 0$ at (a, b) . In this case, we must find some other way to determine the behavior of f at (a, b) .

Examples

The expression $f_{xx}f_{yy} - f_{xy}^2$ is called **the discriminant or Hessian** of f . It is sometimes easier to remember it in determinant form,

$$f_{xx}f_{yy} - f_{xy}^2 = \begin{vmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{vmatrix}.$$

Examples

The expression $f_{xx}f_{yy} - f_{xy}^2$ is called **the discriminant or Hessian** of f . It is sometimes easier to remember it in determinant form,

$$f_{xx}f_{yy} - f_{xy}^2 = \begin{vmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{vmatrix}.$$

- 1 Find all local maxima, local minima, and saddle points.
 - $f(x, y) = x^2 + 2xy$.

Examples

The expression $f_{xx}f_{yy} - f_{xy}^2$ is called **the discriminant or Hessian** of f . It is sometimes easier to remember it in determinant form,

$$f_{xx}f_{yy} - f_{xy}^2 = \begin{vmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{vmatrix}.$$

- 1 Find all local maxima, local minima, and saddle points.
 - $f(x, y) = x^2 + 2xy$. ((0, 0) - saddle point)

Examples

The expression $f_{xx}f_{yy} - f_{xy}^2$ is called **the discriminant or Hessian** of f . It is sometimes easier to remember it in determinant form,

$$f_{xx}f_{yy} - f_{xy}^2 = \begin{vmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{vmatrix}.$$

① Find all local maxima, local minima, and saddle points.

- $f(x, y) = x^2 + 2xy$. ($(0, 0)$ - saddle point)
- $f(x, y) = x^4 + y^4 + 4xy$.

Examples

The expression $f_{xx}f_{yy} - f_{xy}^2$ is called **the discriminant or Hessian** of f . It is sometimes easier to remember it in determinant form,

$$f_{xx}f_{yy} - f_{xy}^2 = \begin{vmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{vmatrix}.$$

① Find all local maxima, local minima, and saddle points.

- $f(x, y) = x^2 + 2xy$. ((0, 0) - saddle point)
- $f(x, y) = x^4 + y^4 + 4xy$. ((0, 0) - saddle point, (-1, 1) and (1, -1) - local minima, $f(0, 0) = 0$, $f(1, -1) = f(-1, 1) = -2$.)

Examples

The expression $f_{xx}f_{yy} - f_{xy}^2$ is called **the discriminant or Hessian** of f . It is sometimes easier to remember it in determinant form,

$$f_{xx}f_{yy} - f_{xy}^2 = \begin{vmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{vmatrix}.$$

① Find all local maxima, local minima, and saddle points.

- $f(x, y) = x^2 + 2xy$. ((0, 0) - saddle point)
- $f(x, y) = x^4 + y^4 + 4xy$. ((0, 0) - saddle point, (-1, 1) and (1, -1) - local minima, $f(0, 0) = 0$, $f(1, -1) = f(-1, 1) = -2$.)
- $f(x, y) = y \sin x$.

Examples

The expression $f_{xx}f_{yy} - f_{xy}^2$ is called **the discriminant or Hessian** of f . It is sometimes easier to remember it in determinant form,

$$f_{xx}f_{yy} - f_{xy}^2 = \begin{vmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{vmatrix}.$$

① Find all local maxima, local minima, and saddle points.

- $f(x, y) = x^2 + 2xy$. ((0, 0) - saddle point)
- $f(x, y) = x^4 + y^4 + 4xy$. ((0, 0) - saddle point, (-1, 1) and (1, -1) - local minima, $f(0, 0) = 0$, $f(1, -1) = f(-1, 1) = -2$.)
- $f(x, y) = y \sin x$. (($n\pi$, 0) - saddle point)

Examples

The expression $f_{xx}f_{yy} - f_{xy}^2$ is called **the discriminant or Hessian** of f . It is sometimes easier to remember it in determinant form,

$$f_{xx}f_{yy} - f_{xy}^2 = \begin{vmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{vmatrix}.$$

① Find all local maxima, local minima, and saddle points.

- $f(x, y) = x^2 + 2xy$. ((0, 0) - saddle point)
- $f(x, y) = x^4 + y^4 + 4xy$. ((0, 0) - saddle point, (-1, 1) and (1, -1) - local minima, $f(0, 0) = 0$, $f(1, -1) = f(-1, 1) = -2$.)
- $f(x, y) = y \sin x$. (($n\pi$, 0) - saddle point)
- $f(x, y) = e^{2x} \cos(y)$.

Examples

The expression $f_{xx}f_{yy} - f_{xy}^2$ is called **the discriminant or Hessian** of f . It is sometimes easier to remember it in determinant form,

$$f_{xx}f_{yy} - f_{xy}^2 = \begin{vmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{vmatrix}.$$

① Find all local maxima, local minima, and saddle points.

- $f(x, y) = x^2 + 2xy$. ((0, 0) - saddle point)
- $f(x, y) = x^4 + y^4 + 4xy$. ((0, 0) - saddle point, (-1, 1) and (1, -1) - local minima, $f(0, 0) = 0$, $f(1, -1) = f(-1, 1) = -2$.)
- $f(x, y) = y \sin x$. (($n\pi$, 0) - saddle point)
- $f(x, y) = e^{2x} \cos(y)$. (No local extrema and no saddle point)

Examples

The expression $f_{xx}f_{yy} - f_{xy}^2$ is called **the discriminant or Hessian** of f . It is sometimes easier to remember it in determinant form,

$$f_{xx}f_{yy} - f_{xy}^2 = \begin{vmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{vmatrix}.$$

① Find all local maxima, local minima, and saddle points.

- $f(x, y) = x^2 + 2xy$. ((0, 0) - saddle point)
- $f(x, y) = x^4 + y^4 + 4xy$. ((0, 0) - saddle point, (-1, 1) and (1, -1) - local minima, $f(0, 0) = 0$, $f(1, -1) = f(-1, 1) = -2$.)
- $f(x, y) = y \sin x$. (($n\pi$, 0) - saddle point)
- $f(x, y) = e^{2x} \cos(y)$. (No local extrema and no saddle point)
- $f(x, y) = \ln(x + y) + x^2 - y$.

Examples

The expression $f_{xx}f_{yy} - f_{xy}^2$ is called **the discriminant or Hessian** of f . It is sometimes easier to remember it in determinant form,

$$f_{xx}f_{yy} - f_{xy}^2 = \begin{vmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{vmatrix}.$$

① Find all local maxima, local minima, and saddle points.

- $f(x, y) = x^2 + 2xy$. ((0, 0) - saddle point)
- $f(x, y) = x^4 + y^4 + 4xy$. ((0, 0) - saddle point, (-1, 1) and (1, -1) - local minima, $f(0, 0) = 0$, $f(1, -1) = f(-1, 1) = -2$.)
- $f(x, y) = y \sin x$. (($n\pi$, 0) - saddle point)
- $f(x, y) = e^{2x} \cos(y)$. (No local extrema and no saddle point)
- $f(x, y) = \ln(x + y) + x^2 - y$. ($\left(\frac{-1}{2}, \frac{3}{2}\right)$ - saddle point)

Examples

The expression $f_{xx}f_{yy} - f_{xy}^2$ is called **the discriminant or Hessian** of f . It is sometimes easier to remember it in determinant form,

$$f_{xx}f_{yy} - f_{xy}^2 = \begin{vmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{vmatrix}.$$

- Find all local maxima, local minima, and saddle points.
 - $f(x, y) = x^2 + 2xy$. ((0, 0) - saddle point)
 - $f(x, y) = x^4 + y^4 + 4xy$. ((0, 0) - saddle point, (-1, 1) and (1, -1) - local minima, $f(0, 0) = 0$, $f(1, -1) = f(-1, 1) = -2$.)
 - $f(x, y) = y \sin x$. ($(n\pi, 0)$ - saddle point)
 - $f(x, y) = e^{2x} \cos(y)$. (No local extrema and no saddle point)
 - $f(x, y) = \ln(x + y) + x^2 - y$. ($(\frac{-1}{2}, \frac{3}{2})$ - saddle point)
- Find the maxima, minima, and saddle points of f , if any, given that $f_x(x, y) = 2x - 4y$ and $f_y = 2y - 4x$.

Examples

The expression $f_{xx}f_{yy} - f_{xy}^2$ is called **the discriminant or Hessian** of f . It is sometimes easier to remember it in determinant form,

$$f_{xx}f_{yy} - f_{xy}^2 = \begin{vmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{vmatrix}.$$

① Find all local maxima, local minima, and saddle points.

- $f(x, y) = x^2 + 2xy$. ((0, 0) - saddle point)
- $f(x, y) = x^4 + y^4 + 4xy$. ((0, 0) - saddle point, (-1, 1) and (1, -1) - local minima, $f(0, 0) = 0$, $f(1, -1) = f(-1, 1) = -2$.)
- $f(x, y) = y \sin x$. (($n\pi$, 0) - saddle point)
- $f(x, y) = e^{2x} \cos(y)$. (No local extrema and no saddle point)
- $f(x, y) = \ln(x + y) + x^2 - y$. ($\left(\frac{-1}{2}, \frac{3}{2}\right)$ - saddle point)

② Find the maxima, minima, and saddle points of f , if any, given that $f_x(x, y) = 2x - 4y$ and $f_y = 2y - 4x$. ((0, 0) - saddle point)

Examples

- ③ Find the critical point of $xy + 2x - \ln x^2y$ in the open first quadrant and show that f takes on a minimum there.

Examples

- ③ Find the critical point of $xy + 2x - \ln x^2y$ in the open first quadrant and show that f takes on a minimum there. ($f(1/2, 2)$ - local minimum)

Examples

- ③ Find the critical point of $xy + 2x - \ln x^2y$ in the open first quadrant and show that f takes on a minimum there. ($f(1/2, 2)$ - local minimum)
- ④ Find the maximum value of $s = xy + yz + xz$ where $x + y + z = 6$.

Examples

- ③ Find the critical point of $xy + 2x - \ln x^2y$ in the open first quadrant and show that f takes on a minimum there. ($f(1/2, 2)$ - local minimum)
- ④ Find the maximum value of $s = xy + yz + xz$ where $x + y + z = 6$. ($s(2, 2, 2) = 12$)

Examples

- ③ Find the critical point of $xy + 2x - \ln x^2y$ in the open first quadrant and show that f takes on a minimum there. ($f(1/2, 2)$ - local minimum)
- ④ Find the maximum value of $s = xy + yz + xz$ where $x + y + z = 6$. ($s(2, 2, 2) = 12$)
- ⑤ Find three positive numbers whose sum is 3 and whose product is a maximum.

Examples

- ③ Find the critical point of $xy + 2x - \ln x^2y$ in the open first quadrant and show that f takes on a minimum there. ($f(1/2, 2)$ - local minimum)
- ④ Find the maximum value of $s = xy + yz + xz$ where $x + y + z = 6$. ($s(2, 2, 2) = 12$)
- ⑤ Find three positive numbers whose sum is 3 and whose product is a maximum. ($(1, 1, 1)$)

Examples

- ③ Find the critical point of $xy + 2x - \ln x^2y$ in the open first quadrant and show that f takes on a minimum there. ($f(1/2, 2)$ - local minimum)
- ④ Find the maximum value of $s = xy + yz + xz$ where $x + y + z = 6$. ($s(2, 2, 2) = 12$)
- ⑤ Find three positive numbers whose sum is 3 and whose product is a maximum. ($(1, 1, 1)$)
- ⑥ Find the minimum distance from the point $(2, -1, 1)$ to the plane $x + y - z = 2$.

Examples

- ③ Find the critical point of $xy + 2x - \ln x^2y$ in the open first quadrant and show that f takes on a minimum there. ($f(1/2, 2)$ - local minimum)
- ④ Find the maximum value of $s = xy + yz + xz$ where $x + y + z = 6$. ($s(2, 2, 2) = 12$)
- ⑤ Find three positive numbers whose sum is 3 and whose product is a maximum. ($(1, 1, 1)$)
- ⑥ Find the minimum distance from the point $(2, -1, 1)$ to the plane $x + y - z = 2$. ($(x, y, z) = (8/3, -1/3, 1/3)$ and distance: $2/\sqrt{3}$)

Examples

- ③ Find the critical point of $xy + 2x - \ln x^2y$ in the open first quadrant and show that f takes on a minimum there. ($f(1/2, 2)$ - local minimum)
- ④ Find the maximum value of $s = xy + yz + xz$ where $x + y + z = 6$. ($s(2, 2, 2) = 12$)
- ⑤ Find three positive numbers whose sum is 3 and whose product is a maximum. ($(1, 1, 1)$)
- ⑥ Find the minimum distance from the point $(2, -1, 1)$ to the plane $x + y - z = 2$. ($(x, y, z) = (8/3, -1/3, 1/3)$ and distance: $2/\sqrt{3}$)
- ⑦ Can you conclude anything about $f(a, b)$ if its first and second partial derivatives are continuous throughout a disk centered at a critical point (a, b) and $f_{xx}(a, b)$ and $f_{yy}(a, b)$ differ in sign? Give reason for your answer.

Examples

- ③ Find the critical point of $xy + 2x - \ln x^2y$ in the open first quadrant and show that f takes on a minimum there. ($f(1/2, 2)$ - local minimum)
- ④ Find the maximum value of $s = xy + yz + xz$ where $x + y + z = 6$. ($s(2, 2, 2) = 12$)
- ⑤ Find three positive numbers whose sum is 3 and whose product is a maximum. ($(1, 1, 1)$)
- ⑥ Find the minimum distance from the point $(2, -1, 1)$ to the plane $x + y - z = 2$. ($(x, y, z) = (8/3, -1/3, 1/3)$ and distance: $2/\sqrt{3}$)
- ⑦ Can you conclude anything about $f(a, b)$ if its first and second partial derivatives are continuous throughout a disk centered at a critical point (a, b) and $f_{xx}(a, b)$ and $f_{yy}(a, b)$ differ in sign? Give reason for your answer. (This can happen only if the point (a, b) is a saddle point. Since opposite sign here implies the Hessian $f_{xx}f_{yy} - f_{xy}^2$ is negative at this point.)

Absolute Maxima and Minima on Closed Bounded Regions

We organize the search for the absolute extrema of a continuous function $f(x, y)$ on a closed and bounded region R into three steps.

Absolute Maxima and Minima on Closed Bounded Regions

We organize the search for the **absolute extrema** of a **continuous function** $f(x, y)$ on a **closed and bounded region** R into three steps.

Steps to find absolute maxima and minima

- 1 *List the interior points* of R where f may have local maxima and minima and evaluate f at these points. These are the **critical points** of f .
- 2 *List the boundary points* of R where f has local maxima and minima and evaluate f at these points.
- 3 *Look through the lists* for the maximum and minimum values of f . These will be the absolute maximum and minimum values of f on R .

Examples

Example 1. Find the absolute maximum and absolute minimum of the function $f(x, y) = x^2 + y^2$ on the closed triangular plate bounded by the lines $x = 0, y = 0, y + 2x = 2$ in the first quadrant.

Examples

Example 1. Find the absolute maximum and absolute minimum of the function $f(x, y) = x^2 + y^2$ on the closed triangular plate bounded by the lines $x = 0, y = 0, y + 2x = 2$ in the first quadrant.

Solution. First let us find the critical points in the interior points of the given domain.

$$f_x = 2x, \quad f_y = 2y.$$

Examples

Example 1. Find the absolute maximum and absolute minimum of the function $f(x, y) = x^2 + y^2$ on the closed triangular plate bounded by the lines $x = 0, y = 0, y + 2x = 2$ in the first quadrant.

Solution. First let us find the critical points in the interior points of the given domain.

$$f_x = 2x, \quad f_y = 2y.$$

So the critical point is $(0, 0)$ which is not an interior point of the given domain. So there is no critical points in the interior of the given domain.

Examples

Example 1. Find the absolute maximum and absolute minimum of the function $f(x, y) = x^2 + y^2$ on the closed triangular plate bounded by the lines $x = 0, y = 0, y + 2x = 2$ in the first quadrant.

Solution. First let us find the critical points in the interior points of the given domain.

$$f_x = 2x, \quad f_y = 2y.$$

So the critical point is $(0, 0)$ which is not an interior point of the given domain. So there is no critical points in the interior of the given domain.

In the boundary, let's check first on the line segment $x = 0, 0 \leq y \leq 2$:

Examples

Example 1. Find the absolute maximum and absolute minimum of the function $f(x, y) = x^2 + y^2$ on the closed triangular plate bounded by the lines $x = 0, y = 0, y + 2x = 2$ in the first quadrant.

Solution. First let us find the critical points in the interior points of the given domain.

$$f_x = 2x, \quad f_y = 2y.$$

So the critical point is $(0, 0)$ which is not an interior point of the given domain. So there is no critical points in the interior of the given domain.

In the boundary, let's check first on the line segment

$$x = 0, \quad 0 \leq y \leq 2:$$

Here the function takes the form $f(x, y) = y^2$ and the only local extremum is at the boundary points $y = 0, 2$ and the function values are $0, 4$ respectively.

Examples

A similar situation occurs with the line segment $y = 0$, $0 \leq x \leq 1$ and local extremum occurs at the boundary points $x = 0, 1$ and the function values are $0, 1$.

Examples

A similar situation occurs with the line segment $y = 0, 0 \leq x \leq 1$ and local extremum occurs at the boundary points $x = 0, 1$ and the function values are 0, 1.

On the line segment $y + 2x = 2, 0 \leq x \leq 1$, the function becomes $f(x) = 5x^2 - 8x + 4$.

Examples

A similar situation occurs with the line segment $y = 0, 0 \leq x \leq 1$ and local extremum occurs at the boundary points $x = 0, 1$ and the function values are 0, 1.

On the line segment $y + 2x = 2, 0 \leq x \leq 1$, the function becomes $f(x) = 5x^2 - 8x + 4$. Here, f has extrema at $f'(x) = 10x - 8 = 0$ or $x = 4/5$ where $y = 2 - 8/5 = 2/5$ and $f(4/5, 2/5) = 4/5$.

Examples

A similar situation occurs with the line segment $y = 0, 0 \leq x \leq 1$ and local extremum occurs at the boundary points $x = 0, 1$ and the function values are 0, 1.

On the line segment $y + 2x = 2, 0 \leq x \leq 1$, the function becomes $f(x) = 5x^2 - 8x + 4$. Here, f has extrema at $f'(x) = 10x - 8 = 0$ or $x = 4/5$ where $y = 2 - 8/5 = 2/5$ and $f(4/5, 2/5) = 4/5$.

Table: Points and corresponding $f(x, y)$

$(0,0)$	0
$(0,2)$	4
$(1,0)$	1
$(4/5, 2/5)$	$4/5$

So the absolute maximum is 4 and the absolute minimum is 0.

Examples

Example 2. Find the absolute maximum and minimum of the function $f(x, y) = (4x - x^2) \cos y$ in the region $1 \leq x \leq 3$, $-\pi/4 \leq y \leq \pi/4$.

Examples

Example 2. Find the absolute maximum and minimum of the function $f(x, y) = (4x - x^2) \cos y$ in the region $1 \leq x \leq 3$, $-\pi/4 \leq y \leq \pi/4$.

Solution. First let us deal with the **interior points**.

$$f_x = (4 - 2x) \cos y = 0$$

$$f_y = -(4x - x^2) \sin y = 0$$

Solving these two gives us $(2, 0)$. The value at this point is 4.

Examples

Example 2. Find the absolute maximum and minimum of the function $f(x, y) = (4x - x^2) \cos y$ in the region $1 \leq x \leq 3$, $-\pi/4 \leq y \leq \pi/4$.

Solution. First let us deal with the **interior points**.

$$f_x = (4 - 2x) \cos y = 0$$

$$f_y = -(4x - x^2) \sin y = 0$$

Solving these two gives us $(2, 0)$. The value at this point is 4.

Boundary points:

Along the line $1 \leq x \leq 3$, $y = -\pi/4$. The function becomes $g(x) = \frac{4x-x^2}{\sqrt{2}}$ the critical point is $x = 2$ and the value is $2\sqrt{2}$.

Along the line $1 \leq x \leq 3$, $y = \pi/4$. $\cos y$ being an even function we have the same situation as above.

Examples

Along the line $-\pi/4 \leq y \leq \pi/4, x = 1$. The function becomes $h(y) = 3 \cos y$ it has a critical point at $y = 0$ and the value is 3

Along the line $-\pi/4 \leq y \leq \pi/4, x = 3$. The function becomes $3 \cos y$, so it is the same situation as above and the value is same 3.

Now the boundary of the lines $(1, -\pi/4), (1, \pi/4), (3, -\pi/4), (3, \pi/4)$ the values are all equal to $3/\sqrt{2}$. Let us now build the table:

Table: Points and corresponding $f(x, y)$

$(2,0)$	4
$(2, -\pi/4)$	$2\sqrt{2}$
$(2, \pi/4)$	$2\sqrt{2}$
$(1,0)$	3
$(3,0)$	3
$(1, -\pi/4), (1, \pi/4), (3, -\pi/4), (3, \pi/4)$	$3/\sqrt{2}$

So the absolute maximum is at $(2,0)$ and the value is 4, and absolute minimum is $3/\sqrt{2}$ at $(1, -\pi/4), (1, \pi/4), (3, -\pi/4), (3, \pi/4)$.

Examples

- ③ Find the absolute maximum and minimum of the function $T(x, y) = x^2 + xy + y^2 - 6x + 2$ in the region given by $0 \leq x \leq 5$ and $-3 \leq y \leq 0$.

Examples

- ③ Find the absolute maximum and minimum of the function $T(x, y) = x^2 + xy + y^2 - 6x + 2$ in the region given by $0 \leq x \leq 5$ and $-3 \leq y \leq 0$. (The absolute maximum is 11 attained at the point $(0, -3)$ and the absolute minimum is -10 attained at the point $(4, -2)$.)

Examples

- ③ Find the absolute maximum and minimum of the function $T(x, y) = x^2 + xy + y^2 - 6x + 2$ in the region given by $0 \leq x \leq 5$ and $-3 \leq y \leq 0$. (The absolute maximum is 11 attained at the point $(0, -3)$ and the absolute minimum is -10 attained at the point $(4, -2)$.)
- ④ Among all closed rectangular boxes of volume 27 cm^3 , what is the smallest surface area?

Examples

- ③ Find the absolute maximum and minimum of the function $T(x, y) = x^2 + xy + y^2 - 6x + 2$ in the region given by $0 \leq x \leq 5$ and $-3 \leq y \leq 0$. (The absolute maximum is 11 attained at the point $(0, -3)$ and the absolute minimum is -10 attained at the point $(4, -2)$.)
- ④ Among all closed rectangular boxes of volume 27 cm^3 , what is the smallest surface area? (Volume $= xyz = 27$ and Surface area $S = 2(xy + yz + xz)$. Eliminate z and find the local minima for $S(x, y)$. Local minimum of $S(3, 3, 3) = 54$.)