

HOMWORK 4 AND 5

MATH 578 NUMERICAL ANALYSIS
DUE IN CLASS TUES NOV 17

This is a double HW so you have extra time

Engineering students: do (1) and (2) and choose 4 problems out of 3 - 8 from the theory section. Math students: do all theory problems.

Theory. References for this section are Boyd, Convex Optimization, and Barvinok, A course in convexity

- (1) (Reference: Strang Linear Algebra p209-210) Find the formula for the projection of a point b in \mathbb{R}^n onto the subspace $c^T x = 0$. More generally, find the formula for the projection of a point b on the subspace spanned by the columns of an $n \times m$ matrix M . Relate this formula to the least squares solution of $Mx = b$.
- (2) (Reference Cheney, Approximation Theory). Consider the inconsistent equations $x = 5, x = -5, x = 1, x = -1$, write these equations as $Mx = b$ where M is 4 by 1, and b is 4 by 1. Define $r = Mx - b$. Find by hand, the solution which minimizes $|r|_2$ and $|r|_\infty$.
- (3) Prove that the permutation matrices are extreme points of the Birkhoff Polytope, which is the set of all double stochastic matrices, $M = (m_{i,j})$ where the rows and columns of M sum to 1. (This is the easy part of the Birkhoff-von Neumann Theorem, p57 of Barvinok).
- (4) Write the matrix M as a convex combination of permutation matrices (you can do this by hand or use CVX).

$$M = \frac{1}{4} \begin{bmatrix} 2 & 2 & 0 \\ 1 & 0 & 3 \\ 1 & 2 & 1 \end{bmatrix}$$

- (5) Show that the set of continuous functions $f : [0, 1] \rightarrow \mathbb{R}$ with $|f(x)| \leq 1$ for all $x \in [0, 1]$ is a convex set.
- (6) Suppose $A \subset \mathbb{R}^d$ is a convex set, and $T : \mathbb{R}^d \rightarrow \mathbb{R}^m$ is an affine transformation, $T(x) = Mx + b$. Prove that $T(A)$ is a convex set as well.
- (7) Recall that the d-cube, is defined to be the set $C = \{x \in \mathbb{R}^d \mid |x|_\infty \leq 1\}$ and the hyper-octahedron is defined by $O = \{x \in \mathbb{R}^d \mid |x|_1 \leq 1\}$. Write each of these sets in two ways. (i) as a polyhedron $Mx \leq b$, for some M, b , (ii) as a polytope, the convex hull of $V = v_1, \dots, v_N$
- (8) Recall the definition of the polar of a set A ,

$$A^\circ = \{y \mid y^T x \leq 1 \text{ for all } x \in A\}$$

- (i) Prove that when A is a polytope, we can restrict to $A^\circ = \{y \mid y^T v_i \leq 1 \text{ for } i = 1, \dots, N\}$ (ii) Find the polar of C, O .

- (9) Prove the easy half of the bipolar theorem, $A \subset (A^\circ)^\circ$. (see Barvinok Ch IV).

Numerical Problems. Use the example code provided to solve these problems in MATLAB, using CVX. Download CVX from <http://cvxr.com/>, and type `cvx_setup` in the MATLAB command line (making sure the CVX directory is in the path) to install CVX.

- (10) Use the forward Euler, backward Euler, and Crank-Nicolson methods to solve the following ODEs: (i) $(x'(t), y'(t)) = (-y(t), x(t))$. (ii) $(x'(t), y'(t)) = (-y(t), x(t))/r(t)$, where $r(t) = \sqrt{x(t)^2 + y(t)^2}$. Plot the trajectories. What happens numerically to $r(t)$ in each case?
- (11) Run the code `ProblemPoly.m` to find best fit polynomials in the norms $p = 1, 2, \infty$. Modify the code to also consider the cases $p = 1.5, 2.5, 4, 8$. What is the difference in the maximum norm between the interpolated function values using these values of p and the other cases?
- (12) Modify the code `ProblemLines.m` to solve problem (2) above.
- (13) Write your own CVX code to solve the facility location problem. Given points $(x_1, y_1), \dots, (x_n, y_n)$ in the plane, find the point (x, y) which minimizes the norm of the vectors of the distances to the points, using the (i) 1-norm, (ii) 2-norm (iii) ∞ norm. Plot some interesting looking examples with small values of n , and compare the solutions in the different norms.
- (14) Write your own CVX code to the linear discrimination problem. (Ref Boyd 8.6.1). Use the data given in the file `discrim.mat`. Find a linear function $p(x) = c^T x - b$ which discriminates the data.