

Comagnetometers as Probes for Ultralight Dark Matter

May 2021



Itay M. Bloch
Tel Aviv University
(Work done in the Rafael Quantum Optics Lab)



Collaborators:
Y. Hochberg, O. Katz, O. Katz
E. Kuflik, G. Ronen, R. Shaham, T. Volansky.
Based on work presented in arxiv:1907.03767, and arXiv:2105.XXXX

Outline

- **Axion Like Particles (ALPs)**
 - ALPs Brief Overview
 - Coherent Interactions
- **Noble-Alkali Comagnetometers**
 - Spin-Based Magnetometry
 - Why Noble-Alkali?
 - Old Results
 - New Results
- **Conclusions**

Axion Like Particles

Axion Like Particles

- Axions were originally a solution to the strong CP problem [Peccei, Quinn 1977; Weinberg 1978; Wilczek 1978]. ALPs however don't have to be related to the strong CP.

Axion Like Particles

- Axions were originally a solution to the strong CP problem [Peccei, Quinn 1977; Weinberg 1978; Wilczek 1978]. ALPs however don't have to be related to the strong CP.
- Pseudo-scalars.

Axion Like Particles


- Axions were originally a solution to the strong CP problem [Peccei, Quinn 1977; Weinberg 1978; Wilczek 1978]. ALPs however don't have to be related to the strong CP.
- Pseudo-scalars.
- Can be a CDM component (we assume all).



Axion Like Particles



- Axions were originally a solution to the strong CP problem [Peccei, Quinn 1977; Weinberg 1978; Wilczek 1978]. ALPs however don't have to be related to the strong CP.
- Pseudo-scalars.
- Can be a CDM component (we assume all).
- Can be very light and remain CDM candidate:

$$m_a(\text{relevant to talk}) < 4 \times 10^{-12} \text{ eV}$$

ALP-SM Interactions







Coherent Interactions (1)

$$n_a = \frac{0.4 \text{ GeV}}{m_a \cdot \text{cm}^3}$$

Coherent Interactions (1)

For light ALPs ($m_a \lesssim 30$ eV), $n_a = \frac{0.4 \text{ GeV}}{m_a \cdot \text{cm}^3} > 1/\lambda_{\text{de-Broglie}}^3$

Coherent Interactions (1)

For light ALPs ($m_a \lesssim 30$ eV), $n_a = \frac{0.4 \text{ GeV}}{m_a \cdot \text{cm}^3} > 1/\lambda_{\text{de-Broglie}}^3$



$$a = a_0 e^{ix^\mu p_\mu}$$

Coherent Interactions (1)

For light ALPs ($m_a \lesssim 30$ eV), $n_a = \frac{0.4 \text{ GeV}}{m_a \cdot \text{cm}^3} > 1/\lambda_{\text{de-Broglie}}^3$



$$a = a_0 e^{ix^\mu p_\mu}$$

In the non-relativistic limit...

Coherent Interactions (2)

$$H_{a\psi\psi} = -g_{a\psi\psi} \vec{b}_a \cdot \vec{S}_\psi = -\vec{b}_{a-\psi} \cdot \vec{S}_\psi$$

$$\vec{b}_{a-\psi} = g_{a\psi\psi} \sqrt{2\rho_a} \cos(m_a t) \cdot \vec{v}_{a-\psi} \quad [\text{astro-ph/9501042}]$$

Coherent Interactions (2)

$$H_{a\psi\psi} = -g_{a\psi\psi} \vec{b}_a \cdot \vec{S}_\psi = -\vec{b}_{a-\psi} \cdot \vec{S}_\psi$$

$$\vec{b}_{a-\psi} = g_{a\psi\psi} \sqrt{2\rho_a} \cos(m_a t) \cdot \vec{v}_{a-\psi} \quad [\text{astro-ph/9501042}]$$

This is an effect linear in $g_{a\psi\psi}$!

Coherent Interactions (2)

$$H_{a\psi\psi} = -g_{a\psi\psi} \vec{b}_a \cdot \vec{S}_\psi = -\vec{b}_{a-\psi} \cdot \vec{S}_\psi$$


$$\vec{b}_{a-\psi} = g_{a\psi\psi} \sqrt{2\rho_a} \cos(m_a t) \cdot \vec{v}_{a-\psi} \quad [\text{astro-ph/9501042}]$$

This is an effect linear in $g_{a\psi\psi}$!

But how to measure it?


Larmor Precession and Zeeman Splitting

Interaction of a classical magnetic field \vec{B} with a spin \vec{S} :




Larmor Precession and Zeeman Splitting

Interaction of a classical magnetic field \vec{B} with a spin \vec{S} :


$$\dot{\vec{S}} = \gamma \vec{S} \times \vec{B} \quad \longrightarrow \quad H = -\gamma \vec{B} \cdot \vec{S}$$

Larmor Precession and Zeeman Splitting


Interaction of a classical magnetic field \vec{B} with a spin \vec{S} :


$$\dot{\vec{S}} = \gamma \vec{S} \times \vec{B} \quad \longrightarrow \quad H = -\gamma \vec{B} \cdot \vec{S}$$

Reminder: $H_{a\psi\psi} = -\vec{b}_{a-\psi} \cdot \vec{S}_\psi$

Larmor Precession and Zeeman Splitting

Interaction of a classical magnetic field \vec{B} with a spin \vec{S} :


$$\dot{\vec{S}} = \gamma \vec{S} \times \vec{B} \quad \longrightarrow \quad H = -\gamma \vec{B} \cdot \vec{S}$$

Reminder: $H_{a\psi\psi} = -\vec{b}_{a-\psi} \cdot \vec{S}_\psi$

Is there a known way to measure magnetic fields?

Outline

- ✓. **Axion Like Particles (ALPs)**
 - ALPs Brief Overview
 - Coherent Interactions
- **Noble-Alkali Comagnetometers**
 - Spin-Based Magnetometry
 - Why Noble-Alkali?
 - Old Results
 - New Results
- **Conclusions**

Outline



Axion Like Particles (ALPs)

- ALPs Brief Overview
- Coherent Interactions
- Noble-Alkali Comagnetometers

Ultra Light ALP DM
generates magnetic-like
fields.

- Spin-Based Magnetometry
- Why Noble-Alkali?
- Old Results
- New Results
- Conclusions

Noble-Alkali Comagnetometers

Bloch Equations

$$\dot{\vec{S}} =$$

Bloch Equations

- * To leading order in important stuff

$$\dot{\vec{S}} =$$

Bloch Equations

- * To leading order in important stuff

$$\dot{\vec{S}} = \gamma \left(\vec{B} + \frac{\vec{b}}{\gamma} \right) \times \vec{S}$$

Torque

(generates transverse from
longitudinal)

Bloch Equations

- * To leading order in important stuff

$$\dot{\vec{S}} = \gamma \left(\vec{B} + \frac{\vec{b}}{\gamma} \right) \times \vec{S} - \Gamma \vec{S}$$

Torque
(generates transverse from longitudinal)

Decaying excitations
(causes stabilization)

Bloch Equations

- * To leading order in important stuff

Creating macroscopic polarization
(generates a non-trivial steady state solution)

$$\dot{\vec{S}} = \gamma \left(\vec{B} + \frac{\vec{b}}{\gamma} \right) \times \vec{S} - \Gamma \vec{S} + R \hat{z}$$

Torque
(generates transverse from longitudinal)

Decaying excitations
(causes stabilization)

Transverse EOMs

We usually assume $\dot{S}_z = 0$ (also that $|\vec{S}| \approx |S_z|$),
And care only about $S_{\perp} = S_x + iS_y$

Transverse EOMs

We usually assume $\dot{S}_z = 0$ (also that $|\vec{S}| \approx |S_z|$),
And care only about $S_\perp = S_x + iS_y$

The transverse EOMs become

$$\dot{\vec{S}} = \gamma \left(\vec{B} + \frac{\vec{b}}{\gamma} \right) \times \vec{S} - \Gamma \vec{S} + R \hat{z}$$

Transverse EOMs

We usually assume $\dot{S}_z = 0$ (also that $|\vec{S}| \approx |S_z|$),
And care only about $S_\perp = S_x + iS_y$

The transverse EOMs become

$$\dot{\vec{S}} = \gamma \left(\vec{B} + \frac{\vec{b}}{\gamma} \right) \times \vec{S} - \Gamma \vec{S} + R \hat{z}$$

\downarrow

$$\dot{S}_\perp = i\gamma \left(B_z + \frac{b_z}{\gamma} \right) S_\perp - i\gamma \left(B_\perp + \frac{b_\perp}{\gamma} \right) S_z - \Gamma S_\perp$$

Transverse EOMs

We usually assume $\dot{S}_z = 0$ (also that $|\vec{S}| \approx |S_z|$),
And care only about $S_\perp = S_x + iS_y$

The transverse EOMs become

$$\dot{\vec{S}} = \gamma \left(\vec{B} + \frac{\vec{b}}{\gamma} \right) \times \vec{S} - \Gamma \vec{S} + R \hat{z}$$

↓

$$\dot{S}_\perp = i\gamma \left(B_z + \cancel{\frac{b_z}{\gamma}} \right) S_\perp - i\gamma \left(B_\perp + \frac{b_\perp}{\gamma} \right) S_z - \Gamma S_\perp$$

Solving the EOMs

$$\vec{b}_{a-\psi} = g_{a\psi\psi} \sqrt{2\rho_a} \cos(m_a t) \cdot \vec{v}_{a-\psi}$$

Solving the EOMs

$$\vec{b}_{a-\psi} = g_{a\psi\psi} \sqrt{2\rho_a} \cos(m_a t) \cdot \vec{v}_{a-\psi}$$

$$\dot{S}_\perp = i\gamma B_z S_\perp - i\gamma \left(B_\perp + \frac{b_\perp}{\gamma} \right) S_z - \Gamma S_\perp$$

Solving the EOMs

$$\vec{b}_{a-\psi} = g_{a\psi\psi} \sqrt{2\rho_a} \cos(m_a t) \cdot \vec{v}_{a-\psi}$$

$$\dot{S}_\perp = i\gamma B_z S_\perp - i\gamma \left(B_\perp + \frac{b_\perp}{\gamma} \right) S_z - \Gamma S_\perp$$

If B_z is constant



(Fourier. From now on I'm
going to ignore subtleties
regarding $\cos(m_a t) \neq e^{im_a t}$)

Solving the EOMs

$$\vec{b}_{a-\psi} = g_{a\psi\psi} \sqrt{2\rho_a} \cos(m_a t) \cdot \vec{v}_{a-\psi}$$

$$\dot{S}_\perp = i\gamma B_z S_\perp - i\gamma \left(B_\perp + \frac{b_\perp}{\gamma} \right) S_z - \Gamma S_\perp$$

If B_z is constant



(Fourier. From now on I'm
going to ignore subtleties
regarding $\cos(m_a t) \neq e^{im_a t}$)

$$S_\perp(\omega = m_a) = \frac{b_\perp + \gamma B_\perp(\omega = m_a)}{(\gamma B_z - m_a) + i\Gamma} S_z$$

The Result

$$S_{\perp} = \frac{b_{\perp} + \gamma B_{\perp}}{i\Gamma + (\gamma B_z - m_a)} S_z$$

The Result

The transverse
spin

$$S_{\perp} = \frac{b_{\perp} + \gamma B_{\perp}}{i\Gamma + (\gamma B_z - m_a)} S_z$$

The transverse spin:

Everything is encoded in the spin projections in the
directions perpendicular to the pump

The Result

$$S_{\perp} = \frac{b_{\perp} + \gamma B_{\perp}}{i\Gamma + (\gamma B_z - m_a)} S_z$$

The transverse spin 

Signal 

Signal:

The thing we want to measure that an ALP generates

The Result

$$S_{\perp} = \frac{b_{\perp} + \gamma B_{\perp}}{i\Gamma + (\gamma B_z - m_a)} S_z$$

The transverse spin S_{\perp} is given by the equation above. The terms in the numerator represent the signal and transverse magnetic fields, while the denominator represents the decay rate and the difference between the magnetic field and the atomic mass. Red annotations with arrows point to each term: "The transverse spin" points to S_{\perp} , "Signal" points to $b_{\perp} + \gamma B_{\perp}$, and "Transverse Magnetic fields" points to γB_{\perp} .

Transverse Magnetic fields:

Can either be noise, or (as we will see) the effect of one atom species on the other. Note that it is proportional to γ .

The Result

$$S_{\perp} = \frac{b_{\perp} + \gamma B_{\perp}}{i\Gamma + (\gamma B_z - m_a)} S_z$$

The transverse spin S_{\perp} is given by the equation above. The components are labeled as follows:

- Signal: $b_{\perp} + \gamma B_{\perp}$
- Transverse Magnetic fields: $i\Gamma + (\gamma B_z - m_a)$
- Spin in the z direction: S_z

Spin in the z direction
Main demand: Don't be tiny

The Result

$$S_{\perp} = \frac{b_{\perp} + \gamma B_{\perp}}{i\Gamma + (\gamma B_z - m_a)} S_z$$

The transverse spin S_{\perp} is given by the equation above. The components of the magnetic field are b_{\perp} and γB_{\perp} . The signal is proportional to $b_{\perp} + \gamma B_{\perp}$. The denominator is $i\Gamma + (\gamma B_z - m_a)$. The spin in the z direction is S_z . The ALP mass is m_a . Transverse magnetic fields are also indicated.

ALP Masses

Our experiments can only probe ultralight ALPs. To date we can probe up to ~ 5 peV, can probably be extended to neV, theoretically μ eV, though that is unlikely.

The Result

$$S_{\perp} = \frac{b_{\perp} + \gamma B_{\perp}}{i\Gamma + (\gamma B_z - m_a)} S_z$$

The transverse spin S_{\perp} is determined by the signal $b_{\perp} + \gamma B_{\perp}$ and the resonance frequency $i\Gamma + (\gamma B_z - m_a)$. The spin in the z direction S_z is influenced by the transverse magnetic fields γB_{\perp} and the ALP mass m_a .

Resonance Frequency

Determined mostly by external magnetic fields (which we can control with coils). Note that it is proportional to γ .

The Result

$$S_{\perp} = \frac{b_{\perp} + \gamma B_{\perp}}{i\Gamma + (\gamma B_z - m_a)} S_z$$

The transverse spin
Decoherence Rate
Signal
Transverse Magnetic fields
Resonance Frequency
ALP mass
Spin in the z direction

Decoherence Rate:

The decoherence rate determines the width of the atomic response to ALPs. Can be mHz-kHz (though exceptions exist). A small decoherence rate can be problematic due to slow response time.

Why Alkali?

$$S_{\perp} = \frac{b_{\perp} + \gamma B_{\perp}}{(\gamma B_z - m_a) + i\Gamma} S_z$$

Why Alkali?

$$S_{\perp} = \frac{b_{\perp} + \gamma B_{\perp}}{(\gamma B_z - m_a) + i\Gamma} S_z$$

We need a way to generate S_z , and to measure S_{\perp} .

Why Alkali?

$$S_{\perp} = \frac{b_{\perp} + \gamma B_{\perp}}{(\gamma B_z - m_a) + i\Gamma} S_z$$

We need a way to generate S_z , and to measure S_{\perp} .

Both can be achieved with optical lasers for Alkali metals

Why Alkali?

$$S_{\perp} = \frac{b_{\perp} + \gamma B_{\perp}}{(\gamma B_z - m_a) + i\Gamma} S_z$$

We need a way to generate S_z , and to measure S_{\perp} .

Both can be achieved with optical lasers for Alkali metals

“Pump” laser polarizes the spins,
“Probe” laser measures them

Why Alkali?

$$S_{\perp} = \frac{b_{\perp} + \gamma B_{\perp}}{(\gamma B_z - m_a) + i\Gamma} S_z$$

We need a way to generate S_z , and to measure S_{\perp} .

Both can be achieved with optical lasers for Alkali metals

“Pump” laser polarizes the spins,
“Probe” laser measures them

Problem:

$SNR \propto \gamma$, and the gyromagnetic ratio of alkali metals is large.

Why Noble?

$$S_{\perp} = \frac{b_{\perp} + \gamma B_{\perp}}{(\gamma B_z - m_a) + i\Gamma} S_z$$

Why Noble?

$$S_{\perp} = \frac{b_{\perp} + \gamma B_{\perp}}{(\gamma B_z - m_a) + i\Gamma} S_z$$

Our problem was that γ_{Alk} was too large

Why Noble?

$$S_{\perp} = \frac{b_{\perp} + \gamma B_{\perp}}{(\gamma B_z - m_a) + i\Gamma} S_z$$

Our problem was that γ_{Alk} was too large

Noble gasses have a gyromagnetic ratio which is smaller by 2-3 orders of magnitude!

Why Noble?

$$S_{\perp} = \frac{b_{\perp} + \gamma B_{\perp}}{(\gamma B_z - m_a) + i\Gamma} S_z$$

Our problem was that γ_{Alk} was too large

Noble gasses have a gyromagnetic ratio which is smaller by 2-3 orders of magnitude!


Noble gasses do not interact with the lasers

but


They can be both polarized, and measured by Alkali spins.

Spin Exchange: Polarizing $S_{\text{Nob},z}$


Spin Exchange: Polarizing $S_{\text{Nob},z}$



Spin Exchange: Polarizing $S_{\text{Nob},z}$




Spin Exchange: Polarizing $S_{\text{Nob},z}$



Spin passed from one atom to the other during collision:




$$R_{\text{induced on Alk}} \propto S_{\text{Nob}}$$




$$R_{\text{induced on Nob}} \propto S_{\text{Alk}}$$


Spin Exchange: Polarizing $S_{\text{Nob},z}$



Spin passed from one atom to the other during collision:



Spin Exchange: “Measuring” $S_{\perp,\text{Nob}}$



Magnetic field from
(quantum) point-like
interactions

$$B_{\text{induced on Alk}} \propto -S_{\text{Nob}}$$


$$B_{\text{induced on Nob}} \propto -S_{\text{Alk}}$$

Comagnetometer

Ingredients List


Comagnetometer Ingredients List

Glass Cell




Comagnetometer Ingredients List

Glass Cell
Alkali Vapor




Comagnetometer Ingredients List

Glass Cell
Alkali Vapor
Noble Gas



Comagnetometer Ingredients List

Glass Cell
Alkali Vapor
Noble Gas




Comagnetometer
Ingredients List


Polarization measurement

Glass Cell
Alkali Vapor
Noble Gas
Lasers

Probe Laser



Pump Laser




Comagnetometer
Ingredients List

Glass Cell
Alkali Vapor
Noble Gas
Lasers

Polarization measurement

Misc:

- **Oven**
- **Magnetic Shields**
- **Magnetic Coils**
- **Optical Components**



Outline

- ✓ • Axion Like Particles (ALPs)
 - ALPs Brief Overview
 - Coherent Interactions
- Noble-Alkali Comagnetometers
 - ✓ • Spin-Based Magnetometry
 - Why Noble-Alkali?
 - Old Results
 - New Results
- Conclusions

Outline

- ✓ • Axion Like Particles (ALPs)
 - ALPs Brief Overview
 - Coherent Interactions
- Noble-Alkali Comagnetometers
 - ✓ • Spin-Based Magnetometry
 - Why Noble-Alkali?
 - Old Results
 - New Results
- Conclusions

Ultra Light ALP DM
generates magnetic-like
fields.

Outline



Axion Like Particles (ALPs)

- ALPs Brief Overview
- Coherent Interactions
- Noble-Alkali Comagnetometers

Ultra Light ALP DM
generates magnetic-like
fields.



- Spin-Based Magnetometry
- Why Noble-Alkali?
- Old Results
- New Results
- Conclusions

Magnetometers can
measure ALPs. Alkali
magnetometers are
easy to work with, while
Noble magnetometers
are more sensitive.

“Compensation Point” Comagnetometer

[IMB, Y. Hochberg, E. Kuflik, T. Volansky. arxiv:1907.03767]

Response to Magnetic Noise

Response to Magnetic Noise

$$S_{\text{Alk}}(\omega = m_a) = \frac{\text{signal} + \gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}}$$

Response to Magnetic Noise

$$S_{\text{Alk}}(\omega = m_a) = \frac{\text{signal} + \gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}}$$

$$B_{\perp,\text{Alk}} = B_{\perp,\text{noise}} + 2\lambda M_{\text{Nob}} S_{\perp,\text{Nob}} / S_{\text{Nob},z}$$

Response to Magnetic Noise

$$S_{\text{Alk}}(\omega = m_a) = \frac{\text{signal} + \gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}}$$

$$B_{\perp,\text{Alk}} = B_{\perp,\text{noise}} + 2\lambda M_{\text{Nob}} S_{\perp,\text{Nob}} / S_{\text{Nob},z}$$

(Ignoring backreaction of Alkali on Noble)

Response to Magnetic Noise

$$S_{\text{Alk}}(\omega = m_a) = \frac{\text{signal} + \gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}}$$

$$B_{\perp,\text{Alk}} = B_{\perp,\text{noise}} + 2\lambda M_{\text{Nob}} S_{\perp,\text{Nob}} / S_{\text{Nob},z}$$

(Ignoring backreaction of Alkali on Noble)

$$\frac{\partial S_{\text{Alk}}}{\partial B_{\perp,\text{noise}}} = \frac{\gamma_{\text{Alk}} S_{z,\text{Alk}}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}} \left(1 + \frac{2\gamma_{\text{Nob}} \lambda M_{\text{Nob}}}{(\gamma_{\text{Nob}} B_{z,\text{Nob}} - m_a) + i\Gamma_{\text{Nob}}} \right)$$

Response to Magnetic Noise

$$S_{\text{Alk}}(\omega = m_a) = \frac{\text{signal} + \gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}}$$

$$B_{\perp,\text{Alk}} = B_{\perp,\text{noise}} + 2\lambda M_{\text{Nob}} S_{\perp,\text{Nob}} / S_{\text{Nob},z}$$

(Ignoring backreaction of Alkali on Noble)

$$\frac{\partial S_{\text{Alk}}}{\partial B_{\perp,\text{noise}}} = \frac{\gamma_{\text{Alk}} S_{z,\text{Alk}}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}} \left(1 + \frac{2\gamma_{\text{Nob}} \lambda M_{\text{Nob}}}{(\gamma_{\text{Nob}} B_{z,\text{Nob}} - m_a) + i\Gamma_{\text{Nob}}} \right)$$

For $\Gamma_{\text{Nob}} \approx 0, m_a \approx 0$, $B_{z,\text{Nob}}$ is tunable
such that $\partial_{B_{\perp,\text{noise}}} S_{\text{Alk}} = 0$

The Compensation Point

The Compensation Point

At the compensation point, any magnetic noise (at low frequencies) has no effect on the alkali spins!

The Compensation Point

At the compensation point, any magnetic noise (at low frequencies) has no effect on the alkali spins!

Additionally, the two species are “in resonance”, allowing for a fast response of the system to sudden changes.

The Compensation Point


At the compensation point, any magnetic noise (at low frequencies) has no effect on the alkali spins!

Additionally, the two species are “in resonance”, allowing for a fast response of the system to sudden changes.

Let's illustrate


Compensation Point Illustration (B)

* A 2D heuristic illustration, so some artistic freedom was taken.




Compensation Point Illustration (B)

* A 2D heuristic illustration, so some artistic freedom was taken.




Compensation Point Illustration (B)

* A 2D heuristic illustration, so some artistic freedom was taken.



Compensation Point Illustration (B)


* A 2D heuristic illustration, so some artistic freedom was taken.



$$S_K^\perp \propto \gamma_K B_{tot}^\perp = \gamma_K B_{ind}^\perp + \gamma_K B_{noise}^\perp = 0!$$


Compensation Point Illustration (b_e)

* A 2D heuristic illustration, so some artistic freedom was taken.




Compensation Point Illustration (b_e)

* A 2D heuristic illustration, so some artistic freedom was taken.




Compensation Point Illustration (b_e)

* A 2D heuristic illustration, so some artistic freedom was taken.




Compensation Point Illustration (b_n)

* A 2D heuristic illustration, so some artistic freedom was taken.




Compensation Point Illustration (b_n)

* A 2D heuristic illustration, so some artistic freedom was taken.




Compensation Point Illustration (b_n)

* A 2D heuristic illustration, so some artistic freedom was taken.





Compensation Point Illustration (b_n)

* A 2D heuristic illustration, so some artistic freedom was taken.




Measurable, Enhanced Signal!

The Romalis Group Comags




[Gergoios Vasilakis Dissertation 2011], [Justin M. Brown Dissertation 2011], [Thomas W. Kornack Dissertation 2005]

Old Data




Results (e)




[Y. Hochberg, E. Kuflik, T. Volansky, IMB 1907.03767. W. A. Terrano, *et al.*: 1508.02463, LUX Collaboration: 1704.02297, M. M. M. Bertolami, *et al.*: 1406.7712, W. A. Terrano, *et al.*: 1902.04246, G. Vasilakis, Dissertation: 2011, J. M. Brown, Dissertation: 2011, T. W. Kornack, Dissertation: 2005].

Results (n)




[Y. Hochberg, E. Kuflík, T. Volansky, IMB 1907.03767. T. Wu *et al.*:1901.10843, C. Abel *et al.*:1708.06367, M. V. Beznogov *et al.*:1806.07991, CASPER Collaboration: 1902.04644, P. W. Graham *et al.*:1709.07852, G. Vasilakis, Dissertation: 2011, J. M. Brown, Dissertation: 2011, T. W. Kornack Dissertation: 2005]

Results (n)




[Y. Hochberg, E. Kuflik, T. Volansky, IMB 1907.03767. T. Wu *et al.*:1901.10843, C. Abel *et al.*:1708.06367, M. V. Beznogov *et al.*:1806.07991, CASPER Collaboration: 1902.04644, P. W. Graham *et al.*:1709.07852, G. Vasilakis, Dissertation: 2011, J. M. Brown, Dissertation: 2011, T. W. Kornack Dissertation: 2005]

Results (n)



Results (p)




Noble and Alkali Spin Detectors for Ultralight Coherent darK matter

NASDUCK collaboration

[**IMB**, Y. Hochberg, O. Katz, O. Katz
E. Kuflik, G. Ronen, R. Shaham, T. Volansky. 2105.XXXX, and a bit of 21YY.XXXX]

Five experiments should occur in the “near” future



Five experiments should occur in the “near” future



Five experiments should occur in the “near” future



1. NASDUCK Floquet

Wait a few slides

Five experiments should occur in the “near” future




1. NASDUCK Floquet

Wait a few slides

2. NASDUCK High Frequencies

Data is waiting to be analyzed.

Five experiments should occur in the “near” future



1. NASDUCK Floquet

Wait a few slides

2. NASDUCK High Frequencies

Data is waiting to be analyzed.

3. NASDUCK Floquet Low

**Experiment will start running in
the next few months**

Five experiments should occur in the “near” future



1. NASDUCK Floquet

Wait a few slides

2. NASDUCK High Frequencies

Data is waiting to be analyzed.


3. NASDUCK Floquet Low

**Experiment will start running in
the next few months**

4. NASDUCK Modulated

**Some data already taken, more
data will be taken soon.**

Five experiments should occur in the “near” future



1. NASDUCK Floquet

Wait a few slides

2. NASDUCK High Frequencies

Data is waiting to be analyzed.

3. NASDUCK Floquet Low

**Experiment will start running in
the next few months**

4. NASDUCK Modulated

**Some data already taken, more
data will be taken soon.**

5. NASDUCK CoComag

**Data exists for Proof of concept.
Needs to be analyzed. No clear
schedule for a full experiment.**

NASDUCK

Floquet

Response to Signal

$$S_{\text{Alk}}(\omega = m_a) = \frac{\gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}} =$$

Response to Signal

$$S_{\text{Alk}}(\omega = m_a) = \frac{\gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}} =$$

$$\frac{\gamma_{\text{Alk}} \lambda M_{\text{Nob}} S_{z,\text{Alk}}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}} \frac{b_{\perp,\text{ALP-Nob}}}{(\gamma_{\text{Nob}} B_{z,\text{Nob}} - m_a) + i\Gamma_{\text{Nob}}}$$

Response to Signal

$$S_{\text{Alk}}(\omega = m_a) = \frac{\gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}} =$$

$\gamma_{\text{Alk}} \lambda M_{\text{Nob}} S_{z,\text{Alk}}$

$b_{\perp,\text{ALP-Nob}}$

$$\boxed{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}} \quad \boxed{(\gamma_{\text{Nob}} B_{z,\text{Nob}} - m_a) + i\Gamma_{\text{Nob}}}$$

Alkali response

Noble response

Response to Signal

$$S_{\text{Alk}}(\omega = m_a) = \frac{\gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}} =$$

$$\frac{\gamma_{\text{Alk}} \lambda M_{\text{Nob}} S_{z,\text{Alk}}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}} b_{\perp,\text{ALP-Nob}}$$

$(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}$
Alkali response

$(\gamma_{\text{Nob}} B_{z,\text{Nob}} - m_a) + i\Gamma_{\text{Nob}}$
Noble response

$$|\gamma_{\text{Alk}}| \gg |\gamma_{\text{Nob}}|, B_{z,\text{Alk}} = B_{z,\text{Nob}} + c$$

Response to Signal

$$S_{\text{Alk}}(\omega = m_a) = \frac{\gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}} =$$

$$\frac{\gamma_{\text{Alk}} \lambda M_{\text{Nob}} S_{z,\text{Alk}}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}} b_{\perp,\text{ALP-Nob}}$$

Alkali response
Noble response

$$|\gamma_{\text{Alk}}| \gg |\gamma_{\text{Nob}}|, B_{z,\text{Alk}} = B_{z,\text{Nob}} + c$$

For large magnetic fields (=high frequencies), we cannot be in resonance for both the alkali and the noble simultaneously!

$$\left| \frac{\gamma_{\text{Alk}} B_{z,\text{Alk}}}{\gamma_{\text{Nob}} B_{z,\text{Nob}}} \right| \gg 1$$

Floquet Fields

Floquet Fields

$$B_z = B_{z,0} + B_F \cos(\omega_F t)$$

Floquet Fields

$$B_z = B_{z,0} + B_F \cos(\omega_F t)$$

$$S_{\text{Alk}}(t) = \frac{\gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}(\omega = m_a) \cdot e^{im_a t}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}}$$

Floquet Fields

$$B_z = B_{z,0} + B_F \cos(\omega_F t)$$

$$S_{\text{Alk}}(t) = \frac{\gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}(\omega = m_a) \cdot e^{im_a t}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}} \rightarrow \sum_n \eta_F^{(n)} \frac{\gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}(\omega = m_a) \cdot e^{im_a t + n\omega_F t}}{(\gamma_{\text{Alk}} B_{z,\text{Alk},0} - m_a - n\omega_F) + i\Gamma_{\text{Alk}}}$$

Floquet Fields

$$B_z = B_{z,0} + B_F \cos(\omega_F t)$$

$$S_{\text{Alk}}(t) = \frac{\gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}(\omega = m_a) \cdot e^{im_a t}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}} \rightarrow \sum_n \eta_F^{(n)} \frac{\gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}(\omega = m_a) \cdot e^{im_a t + n\omega_F t}}{(\gamma_{\text{Alk}} B_{z,\text{Alk},0} - m_a - n\omega_F) + i\Gamma_{\text{Alk}}}$$

For $\omega_F = \gamma_{\text{Alk}} B_{z,\text{Alk},0} - \gamma_{\text{Nob}} B_{z,\text{Nob},0}$, we get that around the floquet frequency

Floquet Fields

$$B_z = B_{z,0} + B_F \cos(\omega_F t)$$

$$S_{\text{Alk}}(t) = \frac{\gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}(\omega = m_a) \cdot e^{im_a t}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}} \rightarrow \sum_n \eta_F^{(n)} \frac{\gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}(\omega = m_a) \cdot e^{im_a t + n\omega_F t}}{(\gamma_{\text{Alk}} B_{z,\text{Alk},0} - m_a - n\omega_F) + i\Gamma_{\text{Alk}}}$$

For $\omega_F = \gamma_{\text{Alk}} B_{z,\text{Alk},0} - \gamma_{\text{Nob}} B_{z,\text{Nob},0}$, we get that around the floquet frequency

$$S_{\text{Alk}}(\omega = m_a + \omega_F) = \eta_F^{(1)} \frac{\gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}(\omega = m_a)}{(\gamma_{\text{Nob}} B_{z,\text{Nob},0} - m_a) + i\Gamma_{\text{Alk}}}$$

Floquet Fields

$$B_z = B_{z,0} + B_F \cos(\omega_F t)$$

$$S_{\text{Alk}}(t) = \frac{\gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}(\omega = m_a) \cdot e^{im_a t}}{(\gamma_{\text{Alk}} B_{z,\text{Alk}} - m_a) + i\Gamma_{\text{Alk}}} \rightarrow \sum_n \eta_F^{(n)} \frac{\gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}(\omega = m_a) \cdot e^{im_a t + n\omega_F t}}{(\gamma_{\text{Alk}} B_{z,\text{Alk},0} - m_a - n\omega_F) + i\Gamma_{\text{Alk}}}$$

For $\omega_F = \gamma_{\text{Alk}} B_{z,\text{Alk},0} - \gamma_{\text{Nob}} B_{z,\text{Nob},0}$, we get that around the floquet frequency

$$S_{\text{Alk}}(\omega = m_a + \omega_F) = \eta_F^{(1)} \frac{\gamma_{\text{Alk}} S_{z,\text{Alk}} B_{\perp,\text{Alk}}(\omega = m_a)}{(\gamma_{\text{Nob}} B_{z,\text{Nob},0} - m_a) + i\Gamma_{\text{Alk}}}$$

So that for $m_a = \gamma_{\text{Nob}} B_{z,\text{Nob},0}$, we can now have both the species in resonance!

NASDUCK Floquet

NASDUCK Floquet

For each measurement, we only get bounds on

$$|m_a - \gamma_{\text{Nob}} B_{z,\text{Nob},0}| < \mathcal{O}(1) \Gamma_{\text{Nob}}$$

Therefore, nearly 3000 measurements were taken during a 5-months period
(active run time ~2 months)

NASDUCK Floquet

For each measurement, we only get bounds on

$$|m_a - \gamma_{\text{Nob}} B_{z,\text{Nob},0}| < \mathcal{O}(1) \Gamma_{\text{Nob}}$$

Therefore, nearly 3000 measurements were taken during a 5-months period
(active run time ~2 months)

Sensitivity was limited by noise of probe beam (i.e. OOM larger than magnetic noise)

NASDUCK Floquet

For each measurement, we only get bounds on

$$|m_a - \gamma_{\text{Nob}} B_{z,\text{Nob},0}| < \mathcal{O}(1) \Gamma_{\text{Nob}}$$

Therefore, nearly 3000 measurements were taken during a 5-months period
(active run time ~2 months)

Sensitivity was limited by noise of probe beam (i.e. OOM larger than magnetic noise)



An improvement by an order of magnitude should not be too difficult*

NASDUCK Floquet

For each measurement, we only get bounds on

$$|m_a - \gamma_{\text{Nob}} B_{z,\text{Nob},0}| < \mathcal{O}(1) \Gamma_{\text{Nob}}$$

Therefore, nearly 3000 measurements were taken during a 5-months period
(active run time ~2 months)


Sensitivity was limited by noise of probe beam (i.e. OOM larger than magnetic noise)




An improvement by an order of magnitude should not be too difficult*

Analysis procedure was determined pre-unblinding, final stages of the processing.

NASDUCK Floquet Results (n)



NASDUCK Floquet Results (p)



Conclusions

Conclusions

- Comagnetometers offer unprecedented sensitivity for Ultralight ALPs

Conclusions

- Comagnetometers offer unprecedented sensitivity for Ultralight ALPs
- The NASDUCK collaboration has many experiments it can do/has already done.

Conclusions

- Comagnetometers offer unprecedented sensitivity for Ultralight ALPs
- The NASDUCK collaboration has many experiments it can do/has already done.
- With creativity, one can think of new experiments to run! We already have several ideas for how to utilize existing experiments for other things.

Noble and Alkali Spin Detectors for Ultralight Coherent darK-matter



DUCK-matter



NASDUCK-matter

Thanks For Listening!