

Modeling the mitigation of dengue fever, Chikungunya and Zika by infecting mosquitoes with *Wolbachia* bacteria

Zhuolin Qu

Mathematics Department
Tulane University

James (Mac) Hyman, Tulane University, USA
Ling Xue, University of Manitoba, Canada

Outlines

- 1 Mosquito-borne diseases v.s. *Wolbachia*
- 2 Vertical transmission *Wolbachia* model
- 3 Numerical simulations of *Wolbachia* mitigation

“Mosquitoes cause more human suffering than any other organism.”


– American Mosquito Control Association

- **dengue fever, Chikungunya**: high fever, muscle and joint pains, may be life-threatening
- **Zika virus**: no or only mild symptoms; infection during pregnancy can cause birth defect of brain

“Mosquitoes cause more human suffering than any other organism.”

– American Mosquito Control Association

- **dengue fever, Chikungunya**: high fever, muscle and joint pains, may be life-threatening
- **Zika virus**: no or only mild symptoms; infection during pregnancy can cause birth defect of brain



Aedes aegypti – primary vector for the transmission

Aedes aegypti – primary vector for the transmission



Aedes aegypti, the
yellow fever mosquito

Aedes aegypti – primary vector for the transmission



Aedes aegypti, the
yellow fever mosquito

mitigation approaches:

- remove breeding habitats
 - water tank/scrap tires
- natural predators
 - fish to control larvae
- spraying of insecticide (most used)
 - financial cost can be prohibitively high
 - logistically difficult in very urban/remote areas

Aedes aegypti – primary vector for the transmission



Aedes aegypti, the
yellow fever mosquito

mitigation approaches:

- remove breeding habitats
 - water tank/scrap tires
- natural predators
 - fish to control larvae
- spraying of insecticide (most used)
 - financial cost can be prohibitively high
 - logistically difficult in very urban/remote areas

- ☹ Eradication failed: lost political importance, no financial support, etc.
- ☹ The re-infestation of *Aedes aegypti* keeps happening.

Wolbachia – a promising strategy to stop diseases at source

- Recently proposed by public health researchers.

Wolbachia – a promising strategy to stop diseases at source

- Recently proposed by public health researchers.

reducing population of the mosquitoes



create a *Wolbachia* epidemic in the population to stop an epidemic

Wolbachia – a promising strategy to stop diseases at source

- Recently proposed by public health researchers.

reducing population of the mosquitoes



create a *Wolbachia* epidemic in the population to stop an epidemic

- a natural parasitic microbe, found in 60% insects, but not in the wild *Aedes aegypti* mosquitoes (reproductive number < 1)
- stops the proliferation of harmful viruses inside the mosquito
→ reduces the disease transmissions


Wolbachia – complex vertical transmission

- *Wolbachia* can be **maternally transmitted** from infected mothers to offspring within the mosquito population.

Wolbachia – complex vertical transmission

- Wolbachia* can be **maternally transmitted** from infected mothers to offspring within the mosquito population.

Schematic of the complex vertical transmission mating



Green: uninfected/natural; Red: *Wolbachia*-infected.

Q: How many *Wolbachia*-infected mosquitoes need to be released?

- release infected mosquitoes into the wild mosquito population

Q: How many *Wolbachia*-infected mosquitoes need to be released?

- release infected mosquitoes into the wild mosquito population
- difficult to *sustain* a stable infection in a wild mosquito population

Q: How many *Wolbachia*-infected mosquitoes need to be released?

- release infected mosquitoes into the wild mosquito population
- difficult to *sustain* a stable infection in a wild mosquito population

fitness-cost:

- ☹ cytoplasmic incompatibility: no viable offspring
- ☹ female lifespan ↓
- ☹ number of eggs produced ↓

Q: How many *Wolbachia*-infected mosquitoes need to be released?

- release infected mosquitoes into the wild mosquito population
- difficult to *sustain* a stable infection in a wild mosquito population




fitness-cost:

- ☹ cytoplasmic incompatibility: no viable offspring
 - ☹ female lifespan ↓
 - ☹ number of eggs produced ↓
- small infection will be wiped out (infection not naturally found in *Aedes aegypti* mosquitoes)

Q: How many *Wolbachia*-infected mosquitoes need to be released?

- release infected mosquitoes into the wild mosquito population
- difficult to *sustain* a stable infection in a wild mosquito population

fitness-cost:

-  cytoplasmic incompatibility: no viable offspring
-  female lifespan ↓
-  number of eggs produced ↓
- small infection will be wiped out (infection not naturally found in *Aedes aegypti* mosquitoes)

If there is a threshold condition that the *Wolbachia* endemic can take off?

Our approach:

- develop an ODE model to describe the complex transmission cycle
- analyze the critical threshold condition for a sustained *Wolbachia* infected population



Outlines


- 1 Mosquito-borne diseases v.s. *Wolbachia*
- 2 Vertical transmission *Wolbachia* model
- 3 Numerical simulations of *Wolbachia* mitigation

Our new model captures the complex transmission cycle

by accounting for ...

- ▶ heterosexual transmission
- ▶ multiple pregnant states for females
- ▶ aquatic-life stage with carrying capacity





ν_w : vertical transmission

σ : mating rate

ϕ_w/ϕ_u : egg laying rates

ψ : egg developing rate

single males $\begin{cases} \dot{M}_u = b_m \psi A_u - \mu_{mu} M_u \\ \dot{M}_w = b_m \psi A_w - \mu_{mw} M_w \end{cases}$

single females $\begin{cases} \dot{F}_u = b_f \psi A_u - \sigma F_u - \mu_{fu} F_u \\ \dot{F}_w = b_f \psi A_w - \sigma F_w - \mu_{fw} F_w \end{cases}$

pregnant females $\begin{cases} \dot{F}_{pu} = \sigma F_u \frac{M_u}{M_u + M_w} - \mu_{fu} F_{pu} \\ \dot{F}_{pw} = \sigma F_w \frac{M_u}{M_u + M_w} + \sigma F_w \frac{M_w}{M_u + M_w} - \mu_{fw} F_{pw} \end{cases}$

aquatic stage $\begin{cases} \dot{A}_u = (\phi_u F_{pu} + \nu_u \phi_w F_{pw}) \left(1 - \frac{A_u + A_w}{K_a}\right) - (\mu_a + \psi) A_u \\ \dot{A}_w = \nu_w \phi_w \left(1 - \frac{A_u + A_w}{K_a}\right) F_{pw} - (\mu_a + \psi) A_w \end{cases}$

Basic Reproduction Number R_0

In epidemiology, the basic reproduction number R_0 of an infection

- is the number of cases one case generates within its infectious period, in a totally susceptible population

Basic Reproduction Number R_0

In epidemiology, the basic reproduction number R_0 of an infection

- is the number of cases one case generates within its infectious period, in a totally susceptible population
- threshold condition for the disease transmission
 - $R_0 > 1$ the infection spreads in a population
 - $R_0 < 1$ the infection dies out

Basic Reproduction Number R_0

In epidemiology, the basic reproduction number R_0 of an infection

- is the number of cases one case generates within its infectious period, in a totally susceptible population
- threshold condition for the disease transmission
 - $R_0 > 1$ the infection spreads in a population
 - $R_0 < 1$ the infection dies out

By *next generation matrix* approach ... (messy calculations) ...

$$R_0 = \nu_w \frac{\mu_{fu} \phi_w (\sigma + \mu_{fu})}{\mu_{fw} \phi_u (\sigma + \mu_{fw})}$$

Basic Reproduction Number R_0

In epidemiology, the basic reproduction number R_0 of an infection

- is the number of cases one case generates within its infectious period, in a totally susceptible population
- threshold condition for the disease transmission
 - $R_0 > 1$ the infection spreads in a population
 - $R_0 < 1$ the infection dies out

By *next generation matrix* approach ... (messy calculations) ...

$$R_0 = \nu_w \frac{\mu_{fu} \phi_w (\sigma + \mu_{fu})}{\mu_{fw} \phi_u (\sigma + \mu_{fw})}$$

$$= \frac{\text{\# of new infected eggs generated by an infected egg}}{\text{\# of new uninfected eggs generated by an uninfected egg}} \\ \text{(within one life cycle)}$$

$R_0 \sim$ Stability analysis

$$R_0 = \frac{\text{\# of new infected eggs generated by an infected egg}}{\text{\# of new uninfected eggs generated by an uninfected egg}} \\ \text{(within one life cycle)}$$

$R_0 \sim$ Stability analysis

$$R_0 = \frac{\text{\# of new infected eggs generated by an infected egg}}{\text{\# of new uninfected eggs generated by an uninfected egg}} \\ \text{(within one life cycle)}$$

The stability analysis on ODE system shows that ...

- $R_0 > 1$ (“new infected > new uninfected”)
 - *infection spreads* \rightarrow system approach to endemic state

$R_0 \sim$ Stability analysis

$$R_0 = \frac{\text{\# of new infected eggs generated by an infected egg}}{\text{\# of new uninfected eggs generated by an uninfected egg}} \\ \text{(within one life cycle)}$$

The stability analysis on ODE system shows that ...

- $R_0 > 1$ (“new infected > new uninfected”)
 - *infection spreads* \rightarrow system approach to endemic state
 - too good to be true (baseline $R_0 \approx 0.7$)

$R_0 \sim$ Stability analysis

$$R_0 = \frac{\text{\# of new infected eggs generated by an infected egg}}{\text{\# of new uninfected eggs generated by an uninfected egg}} \\ \text{(within one life cycle)}$$

The stability analysis on ODE system shows that ...

- $R_0 > 1$ (“new infected > new uninfected”)
 - *infection spreads* \rightarrow system approach to endemic state
 - too good to be true (baseline $R_0 \approx 0.7$)
- $R_0 < 1$ (“new infected < new uninfected”) \leftarrow the real world
 - *infection dies out? Not necessarily!*



$R_0 \sim$ Stability analysis

$$R_0 = \frac{\text{\# of new infected eggs generated by an infected egg}}{\text{\# of new uninfected eggs generated by an uninfected egg}} \\ \text{(within one life cycle)}$$

The stability analysis on ODE system shows that ...

- $R_0 > 1$ (“new infected > new uninfected”)
 - *infection spreads* \rightarrow system approach to endemic state
 - too good to be true (baseline $R_0 \approx 0.7$)
- $R_0 < 1$ (“new infected < new uninfected”) \leftarrow the real world
 - *infection dies out? Not necessarily!*
 - There is a **critical threshold** to maintain *Wolbachia* infection!

Critical threshold: backward bifurcation





- disease-free equilibrium ($R_0 < 1$): stable
- endemic equilibrium (bifurcated)
 - upper branch ($R_0 > 4\nu_u\nu_w$): stable
 - lower branch ($4\nu_u\nu_w < R_0 < 1$): unstable

Outlines

- 1 Mosquito-borne diseases v.s. *Wolbachia*
- 2 Vertical transmission *Wolbachia* model
- 3 Numerical simulations of *Wolbachia* mitigation

It's more effective if we remove some natural population before the release.

It's more effective if we remove some natural population before the release.





Dashed lines: natural; Solid lines: infected

(The percentage is relative to the initial natural population.)

It's more effective to split a big release into several smaller ones.

It's more effective to split a big release into several smaller ones.



Dashed lines: natural; Solid lines: infected

Acknowledgment

This research was partially supported by



- National Science Foundation MPS/DMS/Mathematical Biology Program
- National Institutes of Health NIGMS/MIDAS Program

Thank you!

Releases of infected males only (to wipe out the population)



- requires repetitive releases to wipe out thoroughly
- -25% adult, -50% aquatic, release 80%



Releasing infected males only is not a reliable mitigation strategy

► large release quantity ► local effect only; the environment is not isolated