

# Modeling the transmission of *Wolbachia* in mosquitoes for controlling mosquito-borne diseases

*Zhuolin Qu*

Mathematics Department  
Tulane University

Ling Xue, University of Manitoba, Canada  
James (Mac) Hyman, Tulane University, USA

# Outlines

- 1 Mosquito-borne diseases v.s. *Wolbachia*
- 2 Maternal transmission *Wolbachia* model
- 3 Numerical simulations of *Wolbachia* mitigation

“Mosquitoes cause more human suffering than any other organism.”


– American Mosquito Control Association

- **dengue fever, chikungunya**: high fever, muscle and joint pains, may be life-threatening
- **Zika virus**: no or only mild symptoms; infection during pregnancy can cause birth defect of brain

“Mosquitoes cause more human suffering than any other organism.”

– American Mosquito Control Association

- **dengue fever, chikungunya**: high fever, muscle and joint pains, may be life-threatening
- **Zika virus**: no or only mild symptoms; infection during pregnancy can cause birth defect of brain



<http://www.healthmap.org/zika/#timeline>


# *Aedes aegypti* – primary vector for the transmission

# *Aedes aegypti* – primary vector for the transmission



*Aedes aegypti*, the  
yellow fever mosquito

# *Aedes aegypti* – primary vector for the transmission




*Aedes aegypti*, the  
yellow fever mosquito

## mitigation approaches:

- remove breeding habitats
  - water tank/scrap tires
- introduce natural predators
  - fish to control larvae
- spraying of insecticide (most used)
  - financial cost can be prohibitively high
  - logistically difficult in urban/remote areas
  - evolution of resistance

# *Aedes aegypti* – primary vector for the transmission



*Aedes aegypti*, the  
yellow fever mosquito

## mitigation approaches:

- remove breeding habitats
  - water tank/scrap tires
- introduce natural predators
  - fish to control larvae
- spraying of insecticide (most used)
  - financial cost can be prohibitively high
  - logistically difficult in urban/remote areas
  - evolution of resistance

- ☹ Eradication failed: lost political importance, no financial support, etc.
- ☹ The re-infestation of *Aedes aegypti* keeps happening.



## *Wolbachia* – fight an epidemic with an epidemic

- Recently proposed by public health researchers.

## *Wolbachia* – fight an epidemic with an epidemic

- Recently proposed by public health researchers.

reducing population of the mosquitoes



create a *Wolbachia* epidemic in the mosquito population

## *Wolbachia* – fight an epidemic with an epidemic

- Recently proposed by public health researchers.

reducing population of the mosquitoes



create a *Wolbachia* epidemic in the mosquito population

- a natural parasitic microbe, found in 60% insects, but not in the wild *Aedes aegypti* mosquitoes (reproductive number  $< 1$ )
- stops the proliferation of dengue/chikungunya/Zika viruses inside the mosquitoes → reduces the disease transmission to human


## *Wolbachia* – complex maternal transmission

- *Wolbachia* is **maternally transmitted** from infected mothers to offspring within the mosquito population.

## *Wolbachia* – complex maternal transmission

- Wolbachia* is **maternally transmitted** from infected mothers to offspring within the mosquito population.

### Schematic of the complex maternal transmission mating



■ uninfected mosquitoes


■ *Wolbachia*-infected mosquitoes

♀ female    ♂ male

## *Wolbachia* – complex maternal transmission

- Wolbachia* is **maternally transmitted** from infected mothers to offspring within the mosquito population.

### Schematic of the complex maternal transmission mating



Q: How many *Wolbachia*-infected mosquitoes need to be released?

- release infected mosquitoes into the wild (uninfected) mosquito population

## Q: How many *Wolbachia*-infected mosquitoes need to be released?

- release infected mosquitoes into the wild (uninfected) mosquito population
- difficult to *sustain* a stable infection in a wild mosquito population



## Q: How many *Wolbachia*-infected mosquitoes need to be released?

- release infected mosquitoes into the wild (uninfected) mosquito population
- difficult to *sustain* a stable infection in a wild mosquito population
  - small infection will be wiped out
  - infection is not naturally found in *Aedes aegypti* mosquitoes

## Q: How many *Wolbachia*-infected mosquitoes need to be released?

- release infected mosquitoes into the wild (uninfected) mosquito population
- difficult to *sustain* a stable infection in a wild mosquito population
  - small infection will be wiped out
  - infection is not naturally found in *Aedes aegypti* mosquitoes

fitness cost due to *Wolbachia* infection:

- ☹ female lifespan ↓
- ☹ number of eggs produced ↓

## Q: How many *Wolbachia*-infected mosquitoes need to be released?

- release infected mosquitoes into the wild (uninfected) mosquito population
- difficult to *sustain* a stable infection in a wild mosquito population
  - small infection will be wiped out
  - infection is not naturally found in *Aedes aegypti* mosquitoes

fitness cost due to *Wolbachia* infection:

- ☹ female lifespan ↓
- ☹ number of eggs produced ↓

If there is a tipping point, beyond which the infection could take off?

- develop an ODE model to describe the complex transmission cycle
- analyze the threshold condition for having a stable endemic *Wolbachia*



# Outlines


- 1 Mosquito-borne diseases v.s. *Wolbachia*
- 2 Maternal transmission *Wolbachia* model
- 3 Numerical simulations of *Wolbachia* mitigation

# Our new model captures the complex transmission cycle

by accounting for ...

- two-sex transmission
- multi-stage for female life cycle
- carrying capacity for aquatic stage





$\sigma$ : mating rate

$m_w$ : prop. infected males

$\eta_w/\eta_u$ : egg laying rates

$v_w$ : maternal transmission

$\psi$ : egg developing rate

single males  $\begin{cases} \dot{M}_u = b_m \psi A_u - \mu_{mu} M_u \\ \dot{M}_w = b_m \psi A_w - \mu_{mw} M_w \end{cases}$

single females  $\begin{cases} \dot{F}_u = b_f \psi A_u - (\sigma + \mu_{fu}) F_u \\ \dot{F}_w = b_f \psi A_w - (\sigma + \mu_{fw}) F_w \end{cases}$

pregnant females  $\begin{cases} \dot{F}_{pu} = \sigma \frac{M_u}{M_u + M_w} F_u - \mu_{fu} F_{pu} \\ \dot{F}_{pw} = \sigma F_w - \mu_{fw} F_{pw} \end{cases}$

aquatic stage  $\begin{cases} \dot{A}_u = (\phi_u F_{pu} + v_u \phi_w F_{pw}) \left(1 - \frac{A_u + A_w}{K_a}\right) - (\mu_a + \psi) A_u \\ \dot{A}_w = v_w \phi_w \left(1 - \frac{A_u + A_w}{K_a}\right) F_{pw} - (\mu_a + \psi) A_w \end{cases}$

# Basic Reproductive Number $\mathbb{R}_0$

In epidemiology, the basic reproductive number  $\mathbb{R}_0$  of an infection

- is the number of cases one case generates within its infectious period, in a totally susceptible population

# Basic Reproductive Number $\mathbb{R}_0$

In epidemiology, the basic reproductive number  $\mathbb{R}_0$  of an infection

- is the number of cases one case generates within its infectious period, in a totally susceptible population
- threshold condition for the disease transmission
  - $\mathbb{R}_0 > 1$  the infection spreads out in a population
  - $\mathbb{R}_0 < 1$  the infection dies out



# Basic Reproductive Number $\mathbb{R}_0$

In epidemiology, the basic reproductive number  $\mathbb{R}_0$  of an infection

- is the number of cases one case generates within its infectious period, in a totally susceptible population
- threshold condition for the disease transmission
  - $\mathbb{R}_0 > 1$  the infection spreads out in a population
  - $\mathbb{R}_0 < 1$  the infection dies out

By *next generation matrix* approach ... (messy calculations) ...

$$\mathbb{R}_0 = v_w \frac{\mu_{fu} \phi_w (\sigma + \mu_{fu})}{\mu_{fw} \phi_u (\sigma + \mu_{fw})}$$

# Basic Reproductive Number $\mathbb{R}_0$

In epidemiology, the basic reproductive number  $\mathbb{R}_0$  of an infection

- is the number of cases one case generates within its infectious period, in a totally susceptible population
- threshold condition for the disease transmission
  - $\mathbb{R}_0 > 1$  the infection spreads out in a population
  - $\mathbb{R}_0 < 1$  the infection dies out

By *next generation matrix* approach ... (messy calculations) ...

$$\mathbb{R}_0 = v_w \frac{\mu_{fu} \phi_w (\sigma + \mu_{fu})}{\mu_{fw} \phi_u (\sigma + \mu_{fw})}$$

$$= \frac{\# \text{ of new infected eggs generated by an infected egg}}{\# \text{ of new uninfected eggs generated by an uninfected egg}} \\ \text{(within one life cycle)}$$

# $\mathbb{R}_0 \sim$ Stability analysis

$$\mathbb{R}_0 = \frac{\text{\# of new infected eggs generated by an infected egg}}{\text{\# of new uninfected eggs generated by an uninfected egg}} \\ \text{(within one life cycle)}$$

## $\mathbb{R}_0 \sim$ Stability analysis

$$\mathbb{R}_0 = \frac{\# \text{ of new infected eggs generated by an infected egg}}{\# \text{ of new uninfected eggs generated by an uninfected egg}} \\ \text{(within one life cycle)}$$

The stability analysis on ODE system shows that ...

- $\mathbb{R}_0 > 1$  (“new infected > new uninfected”)
  - *infection spreads*  $\rightarrow$  system approach to endemic state

# $\mathbb{R}_0 \sim$ Stability analysis

$$\mathbb{R}_0 = \frac{\# \text{ of new infected eggs generated by an infected egg}}{\# \text{ of new uninfected eggs generated by an uninfected egg}} \\ \text{(within one life cycle)}$$

The stability analysis on ODE system shows that ...

- $\mathbb{R}_0 > 1$  (“new infected > new uninfected”)
  - *infection spreads*  $\rightarrow$  system approach to endemic state
  - too good to be true (real world  $\mathbb{R}_0 < 1$ )

## $\mathbb{R}_0 \sim$ Stability analysis

$$\mathbb{R}_0 = \frac{\# \text{ of new infected eggs generated by an infected egg}}{\# \text{ of new uninfected eggs generated by an uninfected egg}} \\ \text{(within one life cycle)}$$

The stability analysis on ODE system shows that ...

- $\mathbb{R}_0 > 1$  (“new infected > new uninfected”)
  - *infection spreads*  $\rightarrow$  system approach to endemic state
  - too good to be true (real world  $\mathbb{R}_0 < 1$ )
- $\mathbb{R}_0 < 1$  (“new infected < new uninfected”)
  - *infection dies out?*


# $\mathbb{R}_0 \sim$ Stability analysis

$$\mathbb{R}_0 = \frac{\# \text{ of new infected eggs generated by an infected egg}}{\# \text{ of new uninfected eggs generated by an uninfected egg}} \\ \text{(within one life cycle)}$$

The stability analysis on ODE system shows that ...


- $\mathbb{R}_0 > 1$  (“new infected > new uninfected”)
  - *infection spreads*  $\rightarrow$  system approach to endemic state
  - too good to be true (real world  $\mathbb{R}_0 < 1$ )
- $\mathbb{R}_0 < 1$  (“new infected < new uninfected”)
  - *infection dies out? Not necessarily!*
  - There is a **critical threshold** for *Wolbachia* infection to spread out.

# Critical threshold: bifurcation analysis







# Critical threshold: bifurcation analysis



# Critical threshold: bifurcation analysis




# Critical threshold: bifurcation analysis




# Outlines

- 1 Mosquito-borne diseases v.s. *Wolbachia*
- 2 Maternal transmission *Wolbachia* model
- 3 Numerical simulations of *Wolbachia* mitigation

# Threshold condition for *Wolbachia* transmission



Not enough infected population  
are released  $\Rightarrow$  infection dies out



Stable endemic *Wolbachia* is  
established (90% around day 261)


( $X$  = number of natural pregnant female mosquitoes at disease-free state)

# Integrated mosquito control with pre-release mitigation

- 1 Remove some of the natural population before releases
- 2 Release *Wolbachia*-infected mosquitoes

# Integrated mosquito control with pre-release mitigation

- 1 Remove some of the natural population before releases
- 2 Release *Wolbachia*-infected mosquitoes



Without pre-release mitigation,  
90% infection  $\sim$  day 261

Without pre-release mitigation,  
90% infection  $\sim$  day 203

# Integrated mosquito control with pre-release mitigation

- ① Remove some of the natural population before releases
- ② Release *Wolbachia*-infected mosquitoes

Comparison using different pre-release mitigations

Approach	Target	$T_{90\% \text{ in F}}$
None	N/A	261
Residual spraying	Adults	<b>52</b>
Larval control	Aquatic-stage	203
Sticky ovitrap	Pregnant females	105
Acoustic attraction	Males	215

residual spraying > sticky trap > larval control  $\approx$  acoustic attraction



# Acknowledgment

This research was partially supported by

- NSF/MPS/DMS-NIH/NIGMS award
- NIH-NIGMS Models of Infectious Disease Agent Study (MIDAS) award

# Thank you!



# Sensitivity Analysis

