

partitions-leanblueprint

Carter Hastings

Quinn Oveson

September 7, 2026

0.1 Definitions

Definition 1 (Integer Modular Form). An integer modular form of weight $k \in \mathbb{N}$ is a sequence $a : \mathbb{N} \rightarrow \mathbb{Z}$ such that $\sum_{n=0}^{\infty} a(n)q^n$ is a modular form of weight k , where $q = e^{2\pi iz}$.

Definition 2 (Modular Form Mod ℓ). A modular form mod ℓ of weight $k \in \mathbb{Z}/(\ell-1)\mathbb{Z}$ is a sequence $a : \mathbb{N} \rightarrow \mathbb{Z}/\ell\mathbb{Z}$ such that there exists an integer modular form b of weight k' where $b \equiv a \pmod{\ell}$ and $k' \equiv k \pmod{\ell-1}$.

Definition 3 (Theta). Θ sends modular forms mod ℓ of weight k to weight $k+2$ by $(\Theta a)n = na(n)$.

Definition 4 (U Operator). The operator U sends modular forms mod ℓ of weight k to weight k by $(a|U)n = a(\ell n)$.

Definition 5 (hasWeight). A modular form mod ℓ called a has weight $j \in \mathbb{N}$ if there exists an integer modular form b of weight j such that $b \equiv a \pmod{\ell}$.

Definition 6 (Filtration). Let a be a modular form mod ℓ . The filtration of a , $\omega(a)$, is defined as the minimum natural number j such that a has weight j . The filtration of the zero function is 0.

Definition 7 (multiplication and exponentiation). It's worth stating how multiplication and exponentiation are defined here, because they are not defined in the normal way. The multiplication of two modular forms mod ℓ called a and b is defined as

$$(a \cdot b)n = \sum_{x+y=n} a(x)b(y).$$

The exponentiation of a modular form mod ℓ called a to the power of $j \in \mathbb{N}$ is defined as

$$(a^j)n = \sum_{x_1+\dots+x_j=n} \prod_{i=1}^j a(x_i).$$

Theorem 8 (Pow Prime). Let ℓ be a prime and a a modular form mod ℓ of any weight. Then

$$a^\ell(n) = \begin{cases} a(n/\ell) & \text{if } \ell \mid n \\ 0 & \text{otherwise} \end{cases}$$

Proof.

□

0.2 Theorems

Theorem 9. Let a be a modular form mod ℓ . Then $(a|U)^\ell = a - \Theta^{\ell-1}a$.

Proof.

□

Theorem 10.

If an integer modular form of a weight k reduces to a modular form mod ℓ of weight j then we must have $k \equiv j \pmod{\ell}$

Theorem 11. Let a be a modular form mod ℓ of weight k . Then $\omega(a) \equiv k \pmod{\ell-1}$.

Proof. □

Definition 12 (Eisenstein Series). For $k \geq 2 \in \mathbb{N}$, the Eisenstein series E_k is an integer modular form of weight $2k$ defined by

$$E_k = 1 - \frac{4k}{B_{2k}} \sum_{n=1}^{\infty} \sigma_{2k-1}(n) q^n$$

where B_k is the k th Bernoulli number and $\sigma_k(n)$ is the sum of the k th powers of the divisors of n .

Definition 13 (Delta).

Δ is the sequence obtained from $q(\prod_{n=1}^{\infty} (1 - q^n))^{24}$. It is an integer modular form of weight 12. The modular form mod ℓ , also called Δ , is defined to be the reduction of Δ .

Definition 14. For a prime $\ell \geq 5$, we define $\delta_\ell = \frac{\ell^2 - 1}{24} \in \mathbb{N}$. We define $f_\ell = \Delta^{\delta_\ell}$, which is an integer modular form of weight $12\delta_\ell$. We also define the reduction of f_ℓ as an modular form mod ℓ

Definition 15. For $k \in \mathbb{N}$, define $\dim(k) = \lfloor k/6 \rfloor + \begin{cases} 0 & \text{if } k \equiv 1 \pmod{6} \\ 1 & \text{else} \end{cases}$.

$\dim(k)$ is the dimension of the space of modular forms of weight $2k$.

Definition 16.

For $k \neq 1$ and $c < \dim(k)$, write k uniquely as $k = 6\ell + k'$, where $k' \in \{0, 2, 3, 4, 5, 7\}$. $G_{k,c}$ is a modular form of weight $2k$ defined as

$$G_{k,c} = E_2^a E_3^b E_2^{\ell-3c} \Delta^c$$

where a, b are the unique natural numbers with $2a + 3b = k'$.

We have that the set of all such $G_{k,c}$ for fixed k form a basis for the modular forms of weight $2k$.

Definition 17 (ord). Let b be an integer modular form. The order of b , $\text{ord}(b)$, is defined as the minimum $n \in \mathbb{N}$ such that $b(n) \neq 0$. This is the order of vanishing of b at infinity.

Theorem 18. $\text{ord}(a \cdot b) = \text{ord}(a) + \text{ord}(b)$.

Proof. □

Theorem 19. $(a \cdot b)(\text{ord}(a) + \text{ord}(b)) = a(\text{ord}(a)) \cdot b(\text{ord}(b))$.

Proof. □

Theorem 20. $\text{ord}(\Delta) = 1$.

Proof. □

Theorem 21. For any k , $\text{ord}(E_k) = 0$.

Proof. □

Theorem 22. $\text{ord}(G_{k,c}) = c$.

Proof. □

Theorem 23. $G_{k,c}(c) = 1$.

Proof. □

Theorem 24. *If the coefficients of a and b match up to $\dim(k)$, then $a = b$.*

Proof. □

Theorem 25. *Let a be a modular form of weight $2k$. If $\text{ord}(a) = \dim(k) - 1$ then $a = a(\dim(k) - 1)G_{k, \dim(k)-1}$.*

Proof. □

Theorem 26. *Let a be a modular form mod ℓ of weight k . If $\text{ord}(a) \geq m$, then there exists an integer modular form b of weight k such that*

- (1) $\text{ord}(b) \geq m$, and
- (2) the reduction of b is equal to a .

Proof. □

Lemma 27. $\omega(f_\ell) = 12\delta_\ell = \frac{\ell^2-1}{2}$.

Proof. □

Theorem 28. *This is part (1) of Lemma 2.1.*

Let a be a modular form mod ℓ . Then $\omega(\Theta a) \leq \omega(a) + \ell + 1$.

Theorem 29. *This is part (2) of Lemma 2.1.*

Let a be a modular form mod ℓ . Then $\omega(\Theta a) = \omega(a) + \ell + 1$ if and only if $\ell \nmid \omega(a)$.

Theorem 30. *This is Lemma 3.2.*

For all $m \in \mathbb{N}$, $\omega(\Theta^m f_\ell) \geq \omega(f_\ell) = \frac{\ell^2-1}{2}$.

Proof. □

Theorem 31. *The conclusion of Lemma 3.3*

If $f_\ell|U = 0$ then $\omega(\Theta^{\ell-1} f_\ell) = \frac{\ell^2-1}{2}$

Proof. □

Lemma 32. *Let a be a modular form mod ℓ . If $\ell \mid \omega(a)$ then there exists an $\alpha \in \mathbb{N}$ such that $\omega(\Theta a) = \omega(a) + \ell + 1 - (\alpha + 1)(\ell - 1)$.*

Proof. □

Theorem 33. *If $(f_\ell|U) = 0$ then $\ell \mid \omega(\Theta^{\ell-2} f_\ell)$.*

Proof. □

Lemma 34. *For all $m \in \mathbb{N}$ with $m \leq \frac{\ell+1}{2}$, $\omega(\Theta^m f_\ell) = \frac{\ell^2-1}{2} + m(\ell + 1)$.*

Proof. Induction. □

Theorem 35. $\ell \mid \omega(\Theta^{\frac{\ell+1}{2}} f_\ell)$.

Proof. □

Definition 36. We define α to be the natural number such that $\omega(\Theta^{\frac{\ell+3}{2}} f_\ell) = \frac{\ell^2-1}{2} + \frac{\ell+3}{2}(\ell + 1) - (\alpha + 1)(\ell - 1)$.

Such an α exists, because $\ell \mid \omega(\Theta^{\frac{\ell+1}{2}} f_\ell) = \frac{\ell^2-1}{2} + \frac{\ell+1}{2}(\ell + 1)$.

Definition 37. We define j to be the least natural number such that $\ell \mid \omega(\Theta^{\frac{\ell+3}{2}+j} f_\ell)$. Such a j exists, because $\ell \mid \omega(\Theta^{\ell-2} f_\ell)$.

Note : This definition requires that $(f_\ell|U) = 0$. We will assume this fact from now on.

Lemma 38. $\alpha \leq \frac{\ell+3}{2}$.

Proof.

□

Lemma 39. $j \leq \frac{\ell-7}{2}$.

Proof.

□

Lemma 40. For all $m \leq j$, $\omega(\Theta^{\frac{\ell+3}{2}+m} f_\ell) = \frac{\ell^2-1}{2} + (\frac{\ell+3}{2} + m)(\ell + 1) - (\alpha + 1)(\ell - 1)$.

Proof.

□

Lemma 41. $\ell \mid (j + 1) + (\alpha + 1)$.

Proof.

□

Lemma 42. $\alpha + 1 = \frac{\ell+5}{2}$.

Proof.

□

Theorem 43. $\omega(\Theta^{\frac{\ell+3}{2}} f_\ell) = \frac{\ell^2-1}{2} + 4$.

Proof.

□

Lemma 44. $f_\ell(\delta_\ell + 1) = 1$.

Proof.

□

Definition 45. Let p be the partion function. We say that there is a Ramanujan congruence mod a prime ℓ if $\forall n \in \mathbb{N}$, $\ell \mid p(\ell n - \delta_\ell)$.

Theorem 46. If $n > 0$ and $m \geq n$, then $p(n)$ is equal to the n th coefficient in the product exapansion of

$$\prod_{i=0}^m (1 - X^{i+1})^{-1}$$

Note : In lean, $p(0) = 0$.

Proof. This proof is almost entirely due to Archive.Wiedijk100Theorems.Partition

□

Definition 47. Let α be a commutative semiring, and let $f, g : \mathbb{N} \rightarrow \alpha[[X]]$ be two sequences of power series with coefficients in α . We say that f and g are eventually equal, and write $f \longrightarrow g$, if $\forall n \in \mathbb{N}$, $\exists m \in \mathbb{N}$, $\forall k \leq n$, $\forall j \geq m$, the k th coefficient of $f(j)$ is equal to the k th coefficient of $g(j)$. This is an equivalence relation.

Theorem 48. If $f \longrightarrow f'$ and $g \longrightarrow g'$, then $fg \longrightarrow f'g'$.

Proof.

□

Lemma 49. Let $n, N, \ell \in \mathbb{N}$ with $n < N$ and $\ell > 0$. Let $a : \mathbb{N} \rightarrow \alpha$ be a sequence with coefficients in a semiring α . Then the n th coefficient of

$$\sum_{i=0}^N a(n)X^{\ell i} = \begin{cases} a(n/\ell) & \text{if } \ell \mid n, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. □

Lemma 50. Let $j, \ell, N, M \in \mathbb{N}$ with $\ell > 0$ and $N, M > \ell j$. Let $a, b : \mathbb{N} \rightarrow \alpha$. Then the ℓj th coefficient of

$$\sum_{i=0}^N a(i)X^i \sum_{i=0}^M b(i)X^{\ell i} = \sum_{x+y=j} a(\ell x)b(y)$$

Proof. □

Lemma 51. Let $m, N \in \mathbb{N}$. Let $f : \mathbb{N} \rightarrow \alpha[[X]]$ be a sequence with $f(0) = 0$, the zero power series. Then

$$X^m \sum_{i=0}^N f(i)X^i = \sum_{i=0}^{N+m} f(i-m)X^i$$

Proof. □

Lemma 52. Let $M, \ell, k \in \mathbb{N}$ with $\ell \nmid k$. Then the k th coefficient of

$$\prod_{i=0}^M (1 - X^{\ell(i+1)})^\ell = 0$$

Proof. □

Lemma 53. Let $\ell, K \in \mathbb{N}$ with $\ell > 0$. Then there exists a $c : \mathbb{N} \rightarrow \alpha$ and $M \in \mathbb{N}$ such that

$$\prod_{i=0}^K (1 - X^{\ell(i+1)})^\ell = \sum_{i=0}^M c(i)X^{\ell i}$$

Proof. □

Theorem 54. The coefficients of the Delta product are eventually equal to the coefficients of the Delta function. In other words,

$$X \prod_{i \leq \cdot} (1 - X^{i+1})^{24} \longrightarrow \sum_{i \leq \cdot} \Delta(i)X^i$$

Proof. □

Theorem 55. If $\ell \geq 5$ is prime, then the coefficients of the f_ℓ product are eventually equal to the coefficients of the f_ℓ function. In other words,

$$(X^{\delta_\ell} \prod_{i \leq \cdot} (1 - X^{i+1})^{24\delta_\ell} \longrightarrow \sum_{i \leq \cdot} f_\ell(i)X^i$$

Proof. □

Theorem 56. Let $\ell \geq 5$ be prime. If there is a ramanujan congruence mod ℓ , then $(f_\ell|U) = 0$.

Proof. □

Theorem 57. $E_4(1) = 240$.

Proof. □

Theorem 58. $G_{6\delta_\ell+2, \delta_\ell}(\delta_\ell + 1) \equiv 241 \pmod{\ell}$.

Proof. □

Theorem 59.

$$\Theta^{\frac{\ell+3}{2}} f_\ell \equiv \delta_\ell^{\frac{\ell+3}{2}} G_{6\delta_\ell+2, \delta_\ell} \pmod{\ell}.$$

Proof. □

Theorem 60. $(\Theta^{\frac{\ell+3}{2}} f_\ell)(\delta_\ell + 1) = 241(\delta_\ell + 1)^{\frac{\ell+3}{2}}.$

Proof. □

Theorem 61. *If $\ell \geq 13$ is prime, then $(f_\ell|U) \neq 0$.*

Proof. □

Theorem 62. *If $\ell \geq 13$ is prime, then there does not exist a ramanujan congruence mod ℓ .*

Proof. □