



LightCOM

Lighthouse Consortium on
Obesity Management

Study name: **Lighthouse Consortium on Obesity Management**
Short name: **LightCOM**

SOP title: **Standard Operating Procedure for Clinical
Investigation Day**

SOP short title: **SOP on CID**

SOP Version: **1** Date: **09.04.2024**

Authored by: **Sabrina Wielsøe** Date: **September 2023**

TABEL OF CONTENTS

1	PURPOSE AND APPLICABILITY	3
2	PERSONNEL QUALIFICATIONS AND RESPONSIBILITIES	3
3	EQUIPMENT AND RESOURCES	3
3.1	Equipment.....	3
3.2	Forms.....	3
4	PREPARATION OF PARTICIPANT	3
5	ORDER AND SPECIFIC FOCUS POINT FOR EACH ASSESSMENT	3
6	EXPECTED RESULTS, TROUBLESHOOTING AND CORRECTIVE ACTION	4
7	DATA AND RECORDS MANAGEMENT	4
8	HISTORICAL RECORD of SOP	4

1 PURPOSE AND APPLICABILITY

This document outlines the procedure for clinical investigation days in LightCOM.

2 PERSONNEL QUALIFICATIONS AND RESPONSIBILITIES

This SOP should be referred to for training purposes. All practical training should be conducted by an experienced staff member.

3 EQUIPMENT AND RESOURCES

3.1 Equipment

- A stopwatch that measures in 1/100 seconds.
- Digital weighing scales
- A distance of at least 6 meters of free floor space, with measuring tape for gait speed test.
- Stadiometer
- Constant tension measure tape
- Blood pressure
- DXA scanner
- Equipment for blood and urine sampling

3.2 Forms

- Training checklist and staff training log

4 PREPARATION OF PARTICIPANT

When arriving at the CID the participant should:

- Be fasted from 10 PM the night before (water is allowed)
- Have abstained from smoking and exercise from 2 hours before
- Have abstained from alcohol from 12 hours before

If the participant has not been fasting since 10 PM the night before or has not abstained from alcohol 12 hours before the CID, the blood sample must be postponed to another day; however, it must be collected no later than a week after the CID. The remaining outcomes can still be collected. If the participant has smoked or exercised within two hours before data collection, the CID will continue as planned, but this information should still be noted on the CID checklist in OpenClinica. Please note the blood lipid profile and fasting blood glucose should be collected before randomisation in LightWAY and BAR. If the blood sample is postponed the randomisation must be so to.

5 ORDER AND SPECIFIC FOCUS POINT FOR EACH ASSESSMENT

- Ask the participant to empty their bladder and collect urine sample
- Collect faeces sample (LightBAR DK only)

- Blood pressure (requires 5 min of rest prior, and should be prior to gait speed and Sit-to-Stand tests)
- The following assessment can be performed in random order:
 - Assess anthropometry (Height, Weight, Waist circumference)
 - DXA Scan (including regional Bone mineral density CID 1 and 4) within 2 weeks of the remaining assessments
 - Blood sample
 - Investigator Questionnaire
 - Sociodemographic
 - Health economics
 - Safety endpoints
 - Gait-speed test
 - 30 sec Sit-to-Stand test
 - Self reported Questionnaires if not answered prior to CID
- Attach SENS motion® monitor on the leg.

For trial sites with sub-studies all assessments for sub-studies should be performed **after** primary and secondary outcomes.

Before the participant leaves, inform them of their next appointment (or timeframe for it).

6 EXPECTED RESULTS, TROUBLESHOOTING AND CORRECTIVE ACTION

- NA

7 DATA AND RECORDS MANAGEMENT

- All data will be recorded in OpenClinica at all CIDs.

HISTORICAL RECORD of SOP

Rev# / Date	Written/Revised By	Revision Description	Approval
0. Written	Sabrina Wielsøe		
1. Edited	Sabrina Wielsøe	Describing the action needed when a participant, turns up unfasted.	Sofus Larsen 11.10.2024
2.			
3.			
4.			
Removed / replaced:			