



LightCOM

Lighthouse Consortium on
Obesity Management

Study name: **Lighthouse Consortium on Obesity Management**
Short name: **LightCOM**

SOP title: **Standard Operating Procedure for the measures of
4m gait speed**

SOP short title: **SOP on gait speed**

SOP Version: **1** Date: **09.04.2024**

Authored by: **Sabrina Wielsøe** Date: **23.09.2023**

1	PURPOSE AND APPLICABILITY	3
2	PERSONNEL QUALIFICATIONS AND RESPONSIBILITIES	3
3	EQUIPMENT AND RESOURCES	3
3.1	Equipment	3
3.2	Forms	3
4	PROCEDURE	3
4.1	Instruction to be read to the participant (<i>in bold and italic</i>).....	4
4.3	Timing, Start and Stop	4
6.	EXPECTED RESULTS, TROUBLESHOOTING AND CORRECTIVE ACTION	5
7.	DATA AND RECORDS MANAGEMENT	5
8.	HISTORICAL RECORD of SOP	5

1 PURPOSE AND APPLICABILITY

This document outlines the procedure to follow whilst measuring the 4m gait speed of participants in the LightCOM trial.

The 4m gait speed is a measure of the time it takes for a person to walk four meters at their normal walking pace when starting from a "standing start".

2 PERSONNEL QUALIFICATIONS AND RESPONSIBILITIES

- 4m gait speed training

This SOP should be referred to for training purposes. All practical training should be conducted by an experienced staff member.

3 EQUIPMENT AND RESOURCES

3.1 Equipment


- A stopwatch that measures in 1/100 seconds.
- A distance of at least 6 meters of free floor space, preferably longer.
- Parallel tape lines or other markings on the floor, four meters apart, marking a start and a stop line.

3.2 Forms

- Training checklist and staff training log

4 PROCEDURE

- Measurements will be taken at all CIDs.
- The study participant stands with the feet up to the starting line, if walking assistance equipment is needed this should be placed in front of the participant ready for use (not locked).
- Walking canes can be used if necessary.
- A target (for example a door, chair or wall) which is at least one meter behind the stop line is designated by the tester.
- The tester stands at the starting line and accompanies the participant. The tester walks slightly behind the person.
- Cheering or providing physical support is not permitted.
- To ensure normal walking pace, the instructor shall remind the participant before each attempt to walk at their normal walking pace, as if they were going for a walk.



4.1 Instruction to be read to the participant (*in bold and italic*)

DK

- ***Vi skal nu måle dit normale ganghastighed. Du skal gå i en behageligt/normal hastighed, som hvis du gik en tur.***
- ***Kan du se "målet"?*** (det udpegede mål nævnes/vises).
- ***Du skal gå hen til "målet" (det udpegede mål nævnes) i din normale ganghastighed. Jeg vil tælle ned fra 3. "3-2-1-gå" også kan du begynde at gå.***
- ***Er du klar til at gå i dit normale/behagelige ganghastighed?***
- ***3-2-1-gå***
- Gentag testen efter minimum 20 sekunders pause. Husk at minde deltageren om det skal være i hans/hendes normale ganghastighed før hvert forsøg.

UK

- ***We will now measure your normal walking pace. You should walk at your comfortable/normal pace, as if you are going for a walk.***
- ***Can you see the "target"?*** (the designated target is mentioned/shown).
- ***You must walk to the "target" (the designated goal is mentioned) at your normal walking pace. I will count down "3-2-1 go", and then you can start walking.***
- ***Are you ready to walk at your comfortable/normal pace?***
- ***3-2-1 go***
- Repeat the test after a minimum of 20 seconds' rest. Remember to remind the participant that it must be at his/her normal walking speed before each attempt.

4.3 Timing, Start and Stop

- Look at the participants feet. The stopwatch is started the moment the participant moves a foot to take a step.
- Timing is stopped the moment the front foot touches or passes the stop line while the participant is walking towards the designated target.
- A test session should, as far as possible, consist of three attempts with approximately 20 seconds between each attempt. Note if the participant is unable to complete three attempts, the participant performs the test once or twice.

- All times will be recorded in seconds to 2 decimal places. OpenClinica will automatically calculate the gait speed the average gait time
- If the participant uses any walking equipment, this is to be noted in the eCRF.

6. EXPECTED RESULTS, TROUBLESHOOTING AND CORRECTIVE ACTION

- NA

7. DATA AND RECORDS MANAGEMENT

- All data will be recorded in OpenClinica at all CIDs.

8. HISTORICAL RECORD of SOP

Rev# / Date	Written/Revised By	Revision Description	Approval
0. Written	Sabrina Wielsøe		
1. Edited			
2.			
3.			
4.			
5.			
6.			
Removed / replaced:			