

Study name: **Lighthouse Consortium on Obesity Management**
Short name: **LightCOM**

SOP title: **Standard Operating Procedure for Collection of a
Routine Urine Sample**

SOP short title: **SOP on Routine Urine Sample**

SOP Version: **1** Date: **09.04.2024**

Authored by: Odette Dawkins Date: 21.09.2023

TABLE OF CONTENTS

1	PURPOSE	2
2	INTRODUCTION	2
3	SCOPE	2
4	RESPONSIBILITIES	2
4.1	Chief Investigator/Principle Investigator or delegate	2
5	SPECIFIC PROCEDURE.....	3
6	FORMS/TEMPLATES TO BE USED.....	3
7	INTERNAL AND EXTERNAL REFERENCES	3
7.1	Internal References	3
7.1	External References	3
8	HISTORICAL RECORD OF SOP	3

1 PURPOSE

The purpose of this Standard Operating Procedure (SOP) is to describe the standard procedures to be followed when asking a participant to collect a routine urine sample during the LightCOM studies which includes LightBAR, LightWAY and LightCARE.

2 INTRODUCTION

A routine urine sample will be collected from participants at four timepoints within the study. The routine urine sample will be for immediate analysis on receiving them at the clinic assessment visit for creatinine, albumin and for estimation of albumin/creatinine ratio.

3 SCOPE

This SOP applies to clinical research carried out under the LightCOM research studies which incorporates the individual studies of LightBAR, LightWay and LightCARE. The sponsor is Department of Medicine, Copenhagen University Hospital Amager and Hvidovre, Copenhagen, Denmark.

4 RESPONSIBILITIES

4.1 Chief Investigator/Principle Investigator or delegate

Will be responsible for ensuring the staff member assigned to see the participant in the clinical setting and to test the urine, is suitably qualified to undertake the procedure and has received adequate and up to date training.

5 SPECIFIC PROCEDURE

Ensure the participant is happy to give a urine sample by checking you have adequate signed consent on the outset of the study

Issue the participant with a sterile screw top container suitable for holding a urine sample and ensure the participant's study ID and barcode is recorded on the container.

- Give clear verbal instructions to the patient for giving a urine sample.
- Ensure the participant is instructed to provide a mid-stream urine sample. Immediately after the participant has handed you their urine sample, check that they have provided enough:
 - 5 mL for immediate analysis
 - 10 mL for bioresource

6 FORMS/TEMPLATES TO BE USED

Use local specified urine request forms for immediate analysis.

Use Study database for urine test results as per study protocol.

7 INTERNAL AND EXTERNAL REFERENCES

7.1 Internal References

Design and Development of SOP – CTRG Template SOP V3.2

7.2 External References

NHS England – Collection of routine urine sample.

8 HISTORICAL RECORD OF SOP

Rev# / Date	Written/Revised By	Revision Description	Approval
0. Written	Odette Dawkins		
1. Edited			
2.			
3.			
4.			
5.			
6.			
Removed / replaced:			