



LightCOM

Lighthouse Consortium on
Obesity Management

Study name: **Lighthouse Consortium on Obesity Management**
Short name: **LightCOM**

SOP title: **Standard Operating Procedure for the measures of 30
sec Sit-to-stand test**

SOP short title: **SOP on STS**

SOP Version: **1** Date: **09.04.2024**

Authored by: **Sabrina Wielsøe** Date: **September 2023**

TABLE OF CONTENTS

1	PURPOSE AND APPLICABILITY	3
2	PERSONNEL QUALIFICATIONS AND RESPONSIBILITIES	3
3	EQUIPMENT AND RESOURCES.....	3
3.1	Equipment.....	3
3.2	Forms.....	3
4	PROCEDURE	3
4.1	Preparation of the participant.....	3
4.2	Test Instruction to the participant (in bold and italic)	4
4.3	Timing and counting.....	5
5	EXPECTED RESULTS, TROUBLESHOOTING AND CORRECTIVE ACTION	5
6	DATA AND RECORDS MANAGEMENT.....	6
7	HISTORICAL RECORD of SOP	6

1 PURPOSE AND APPLICABILITY

This document outlines the procedure to follow whilst undertaking measures 30 sec Sit-to-stand (STS) test in participants of the LightCOM trial. The 30 second STS test measures how many times the participant can get up from a chair in 30 seconds.

2 PERSONNEL QUALIFICATIONS AND RESPONSIBILITIES

- 30 sec Sit- to-stand test training

This SOP should be referred to for training purposes. All practical training should be conducted by an experienced staff member.

3 EQUIPMENT AND RESOURCES

3.1 Equipment

- A high-backed chair, with seat height 43-44cm. The chair should have a wide seat.
- It is recommended that the chair be placed against a wall to prevent it from moving.
- A stopwatch for measuring 30 seconds.
- Optional: hand counter for counting the number of repetitions

3.2 Forms

- Training checklist and staff training log

4 PROCEDURE

- Measurements will be taken by a member of the research team at all CIDs.

4.1 Preparation of the participant

- Instruct the participant to sit in the middle of the chair (in an anterior-posterior direction) with their back straight, feet flat on the floor (side by side) and their arms crossed at the wrists on the chest. They cannot use the armrests during the test.
- First demonstrate the test slowly to show the correct technique, and then at a faster pace to show that the purpose is to do it as quickly as you can, but at a pace where you feel safe from falling over at the same time.
- The participant must complete as many "uprights" as possible within 30 seconds.
- Note the participant has to have fully extended their hips and knees at the top position of each repetition, and have made contact between the hamstrings and chair in the bottom position, but there is no requirement for the participant to be fully seated.
- Before the actual test, the participant can practice 1-2 uprights to ensure the correct technique.

- You may not assist or provide support during the test.
- The test is performed twice with a minimum of 2 minutes' rest in between. The result of both attempts should be noted in the eCRF.



4.2 Test Instruction to the participant (in bold and italic)

DK

- *Vi skal finde ud af, hvor stærk du er i dine ben, ved at se, hvor mange gange du kan rejse dig fra en stol og sætte dig på 30 sekunder. Jeg viser dig lige, hvordan testen skal udføres.*
- Vis testen (rejse sig-sætte sig), først i langsomt tempo for at demonstrere teknikken.
- *Sæt dig midt på stolen (med ret ryg) og fødderne ved siden af hinanden. Håndleddene krydser du og lægger hænderne fladt på brystet. Det er vigtigt, at du rejser dig helt op og sætter dig ned, for ellers tæller det ikke med. Når jeg siger **START** på kommandoen klar-parat-**START**, skal du rejse og sætte dig så mange gange som muligt i 30 sekunder.*

- Vis testen i hurtigt tempo, så deltageren er klar over, at man skal gøre det så hurtigt man kan.
- *Nu får du lov til at prøve 1-2 gange.*
- Før selve testen skal deltageren øve 1-2 oprejsninger for at sikre den korrekte teknik.
- Har du nogle spørgsmål? Eller er du klar?
- Okay Klar- Parat-START

UK

- *We will find out how strong you are in your legs by seeing how many times you can stand up from a chair and sit down in 30 seconds. I'll just show you how to do the test.*
- Show the test (stand up-sit down), first at a slow pace to demonstrate the technique.
- *Sit in the middle of the chair (with your back straight) and your feet next to each other. Cross your wrists and place your hands flat on your chest. It is important that you stand up completely and sit down, otherwise the repetition will not count.*
- *When I say Go on the command ready-set-GO, get up and sit down as many times as possible for 30 seconds.*
- Show the test at a fast pace so that the participant is aware that you have to do it as quickly as you can
- *Please now try 1-2 times.*
- Before the actual test, the participant must practice 1-2 uprights to ensure the correct technique.
- *Do you have any questions or are you ready?*
- *Okay, Ready- set - GO*

4.3 Timing and counting

- The stopwatch is started on the START/GO command, even if the test subject waits a little to get up.
- If the participant does not stand up completely during the test, the participant is asked to stand up completely or to slow down until she/he can perform the movement as prescribed. The timing is not stopped!
- Time is stopped after 30 seconds. During the 30 seconds, the number of full stands is counted. If the participant has risen more than halfway up at 30 seconds, it counts as full stand repetition.
- If you are in doubt as to whether the last repetition counts, ALWAYS choose the worst score.

5 EXPECTED RESULTS, TROUBLESHOOTING AND CORRECTIVE ACTION

- NA

6 DATA AND RECORDS MANAGEMENT

- All data will be recorded in OpenClinica at all CIDs.

7 HISTORICAL RECORD of SOP

Rev# / Date	Written/Revised By	Revision Description	Approval
0. Written	Sabrina Wielsøe		
1. Edited			
2.			
3.			
4.			
5.			
6.			
Removed / replaced:			