Psychiatric Emergencies

January GEMRU Training
Olivia Moumne & Cheleste Onorato

Psychiatric Disturbance & Excited Delirium

Excited Delirium

Symptoms: agitation, anxiety, hallucination, disorientation, violent and bizarre behavior, insensitivity to pain, elevated body temperature and superhuman strength, fast breathing (acidosis)

Can lead to cardiac/respiratory arrest, often from being tased

Commonly found in: males, alcohol withdrawal patients, drug use, head injury pts, mental illness

https://www.youtube.com/watch?v=k8eyWHf1y50

Psychiatric Disturbance

A psychiatric disturbance is defined by an individual who is presenting with acute mental distress or disability not associated with a medical condition.

- -can be mental or behavioral
- -interferes with daily behaviors and interactions with others
- -can be specific (anxiety attack, schizophrenic attack), or generalized unusual behavior

https://www.youtube.com/watch?v=T5bX5a8FZ9E

What to do

- -contact LEA immediately (if not already present)
- -assess whether the scene is safe
- -once scene is secure, attempt to deescalate the situation ..

-tips:

- -one person at a time
- -always have an exit
- -do not touch patient if not necessary, and explain everything are going to touch them

to pt if you

-assess patient as usual (vitals, SAMPLE, etc)

Attempted Suicide

Overview

MOST IMPORTANT: treat life threatening emergencies

If time: gather patient information, vitals, history, history of suicide attempts, etc.

- -Scene safety is vital
- -contact LEA
- -make the patient comfortable, listen to them, validate their concerns

Suicide Assessment

DEAD PIMP

D: disorder/depression

E: environmental stressors

A: access to firearms

D: disinhibition

P: previous attempts

I: ideation

M: male

P: plan

Method (success rate)

Firearms (82.5%)

Drowning (66%)

Hanging / Suffocation (61%)

Gas Poisoning (42%)

Jumping (34.5%)

Ingestion (1.5%)

Cutting (1.2%)

Scenarios!!