## Stats 110 Strategic Practice and Homework 1

#### Daniel Flannery

June 20, 2015

### 1 Strategic Practice

#### 1.1 Naive Definition of Probability

- 1. (a) The probability of rolling 21 with 4 fair dice is > The probability of rolling 22 with for 4 fair dice.
  - 4 fair dice have a sum in the range [4,24] with  $6^4 = 1296$  combinations. A 21 can be constructed from the partitioned into (6,6,6,3),(6,6,5,4),(6,5,5,5); while 22 can partitioned into (6,6,6,4),(6,6,5,5) For example, for (6,6,6,3) there are 4 possibilities: 4 dice, choose one to be 3, the rest are 6. For example, for (6,6,5,4) there are  $\frac{4!}{2!} = 12$  possibilities because there are 4 dice, you must choose one to be 5, one to be 4, and the rest to be 6. Divide by 2! to account for the possible orderings of 6 due to repetition. For example, for (6,6,5,5) there are  $\frac{4!}{2!2!} = 6$  possibilities because there are two possible values and order does not matter. Therefore, there are 4+4+12=20 ways to get 22. Therefore, there are 4+6=10 ways to get 21.
  - (b) The probability of a random 2 letter word being a palindrome = The probability of a random 3 letter word being a palindrome.

For a two letter random word, there are  $26^2$  outcomes. A two letter word is a palindrome when both letters are the same, so there are 26 palindrome and the probability is  $\frac{26}{26^2}$ . For a three letter random word, there are  $26^3$  outcomes. A three letter word is a palindrome when the first and third letter are the same, so there are 26 letters, then for each combination, the second letter can be any of the 26 letters. Then by the multiplication rule, there are 26\*26 palindrome and the probability is  $\frac{26^2}{26^3}$ 

$$P(2 \ letter \ palindrome) = P(3 \ letter \ palindrome) = \frac{26}{26^2} = \frac{26^2}{26^3}$$

A three letter palindrome is an expansion of a two letter palindrome.

- 2. Random 5 card poker hand from a standard deck
  - (a) A flush (all 5 cards of same suit, excluding royal flush)

There are 4 suits. For each suit there are 13 cards. There  $\binom{13}{5}$  ways to choose a 5 card hand for each suit. 1 hand is excluded, the royal flush. The probability is:

$$\frac{(\#suits*(allowable\ hands)}{all\ possible\ hands}$$
$$\frac{4(\binom{13}{5}-1)}{\binom{52}{5}}$$

(b) Two pair

Choose 2 ranks of 13 possible. Choose 2 of the 4 possible cards for each rank. Choose 1 of the remaining cards that is not of the 2 selected ranks. Divide by all possible hands.

$$\frac{\binom{13}{2} \cdot \binom{4}{2} \cdot \binom{4}{2} \cdot 44}{\binom{52}{5}}$$

- 3. Paths on a grid from (0,0) to (110,111) where the only moves are one unit up or one unit down.
  - (a) Encode the moves as such: U for up and R for right. There are 110-0 U's and 111-0 R's. Therefore the set of moves can be written as: URURURRRR...UR. There are then 110+111=221 symbols to be positioned. To determine the number of paths choose a symbol to position:

$$\binom{221}{110} = \binom{221}{111}$$

(b) Paths from (0,0) to (210,211) that go through (110,111)

From above there are  $\binom{221}{110}$  paths from (0,0) to (110,111). Now merely need to find the number of paths from (110,111) to (210,211). This is then 210-110=100 more R's and 211-111=100 more U's. Therefore  $\binom{200}{100}$  paths. By the multiplication rule then there are

$$\binom{221}{110}\cdot \binom{200}{100}$$

total paths.

4. A no-repeat-word is a sequence of 26 letters unique and with no repetition. Show that the probability that a no-repeat-word using all 26 letters selected at random with all no-repeat-word equally likely is very close to 1/e.

There are 26! total no-repeat-words of length 26. To create a no-repeat-word of length k, first select k letters from the 26.  $\binom{26}{k}$ . Then those letters can be arranged into k! words. Therefore there are  $\binom{26}{k}k$ ! no-repeat-words with k letters. To count all possible no-repeat-words, simply sum all the possible words of each possible length: [1,26]. Then, the probability of a no-repeat-word having all 26 letters is:

$$= \frac{26!}{\sum_{k=1}^{26} {26 \choose k} k!} = \frac{26!}{\sum_{k=1}^{26} 26 \frac{26!}{k!(26-k)} k!}$$

$$= \frac{26!}{26! \cdot \sum_{k=1}^{26} \frac{k!}{k!(26-k)!}} = \frac{1}{\sum_{k=1}^{26} \frac{1}{(26-k)!}}$$

$$= \frac{1}{\frac{1}{25!} + \frac{1}{24!} + \dots + \frac{1}{1!} + 1} = 1 + \frac{1}{1!} + \frac{1}{2!} + \dots + \frac{1}{25!}$$

which is equivalent to the Taylor series  $e^x = 1 + x + \frac{x^2}{2!} + \dots + \frac{x^{25}}{25!}$  evaluated at x = 1.

#### 1.2 Story Proofs

# 2 Homework