

UNIVERSITY OF CALIFORNIA, SAN DIEGO

**A Search for New Physics producing Jets, Large MT₂, and Disappearing Tracks in 13 TeV
Proton-Proton Collision at CERN's Large Hadron Collider**

A dissertation submitted in partial satisfaction of the
requirements for the degree
Doctor of Philosophy

in

Physics

by

Dylan Gilbert

Committee in charge:

Professor Avraham Yagil, Chair
Professor Frank Würthwein, Co-Chair
Professor Rommie Amaro
Professor Adam Burgasser
Professor George Fuller

2019

Copyright
Dylan Gilbert, 2019
All rights reserved.

The dissertation of Dylan Gilbert is approved, and it is acceptable in quality and form for publication on microfilm and electronically:

Co-Chair

Chair

University of California, San Diego

2019

TABLE OF CONTENTS

| | |
|---|------|
| Signature Page | iii |
| Table of Contents | iv |
| List of Figures | vi |
| List of Tables | viii |
| Acknowledgements | ix |
| Vita | x |
| Abstract of the Dissertation | xi |
| | |
| Chapter 1 Introduction | 1 |
| 1.1 The Standard Model of Particle Physics | 1 |
| 1.1.1 A Brief Description | 1 |
| 1.1.2 Feynman Diagrams | 1 |
| 1.1.3 Quantum Chromodynamics, Hadronization, and Jets | 2 |
| 1.1.4 Parton Distribution Functions | 2 |
| 1.2 Some Problems with the Standard Model | 2 |
| 1.2.1 Effective Field Theory | 2 |
| 1.2.2 Gravity | 2 |
| 1.2.3 The Hierarchy Problem | 2 |
| 1.2.4 Astrophysical Evidence of Dark Matter | 3 |
| 1.3 Supersymmetry | 3 |
| 1.3.1 Theoretical Appeal | 3 |
| 1.3.2 Experimental Signatures | 3 |
| 1.3.3 Simplified Models | 3 |
| 1.4 Other Models | 4 |
| | |
| Chapter 2 The Large Hadron Collider and the CMS Detector | 5 |
| 2.1 The Large Hadron Collider | 5 |
| 2.1.1 A Brief Description | 6 |
| 2.1.2 Luminosity Delivered | 7 |
| 2.2 The CMS Detector | 8 |
| 2.2.1 Magnet | 10 |
| 2.2.2 Tracker | 11 |
| 2.2.3 Electromagnetic Calorimeter | 17 |
| 2.2.4 Hadronic Calorimeter | 18 |
| 2.2.5 Muon System | 19 |
| 2.2.6 Missing Transverse Energy | 20 |

| | | |
|---------------------|---|------------|
| 2.3 | CMS Event Reconstruction | 25 |
| 2.3.1 | Challenges | 25 |
| 2.3.2 | Vertices | 25 |
| 2.3.3 | Particle Flow | 26 |
| 2.3.4 | Jet Clustering | 26 |
| 2.4 | Simulation | 27 |
| 2.5 | Objectives | 27 |
| 2.6 | Limitations and Challenges | 27 |
| 2.7 | The Simulation Pipeline | 27 |
| Chapter 3 | The Full 13 TeV M_{T2} Analysis | 28 |
| 3.1 | Classic Search | 29 |
| 3.1.1 | General Description | 29 |
| 3.1.2 | Signals | 30 |
| 3.1.3 | Backgrounds | 37 |
| 3.1.4 | Baseline Selection | 46 |
| 3.1.5 | Upgrades in 2019 | 46 |
| 3.1.6 | Signal Contamination | 52 |
| 3.1.7 | Results | 53 |
| 3.1.8 | Limits | 53 |
| 3.1.9 | Future of the Classic M_{T2} Search | 61 |
| 3.2 | Disappearing Tracks Search | 64 |
| 3.2.1 | General Description | 64 |
| 3.2.2 | Challenges | 64 |
| 3.2.3 | Signal Models | 65 |
| 3.2.4 | Backgrounds | 67 |
| 3.2.5 | The Short Track Selection and Data-Driven Background Estimate | 71 |
| 3.2.6 | Systematics | 77 |
| 3.2.7 | Results | 78 |
| 3.2.8 | Limits | 81 |
| 3.2.9 | Future | 89 |
| Chapter 4 | Conclusions | 91 |
| Appendix A | Full Classic Binning and Results | 92 |
| Bibliography | | 101 |

LIST OF FIGURES

| | | |
|--------------|--|----|
| Figure 1.1: | Theoretical gluino pair production cross section in simplified models. | 4 |
| Figure 2.1: | CMS integrated luminosity delivered and recorded over time. | 8 |
| Figure 2.2: | Image depicting the experimental signatures of different types of particle at CMS. | 9 |
| Figure 2.3: | Layout of the tracker up to 2016. | 11 |
| Figure 2.4: | Efficiency for a genuine particle to leave an expected hit in the CMS tracker. | 12 |
| Figure 2.5: | Track fake rate. | 14 |
| Figure 2.6: | Track reconstruction efficiency. | 14 |
| Figure 2.7: | Impact parameter resolution in the CMS tracker. | 15 |
| Figure 2.8: | Tracker material budget. | 15 |
| Figure 2.9: | Pion survival in the tracker. | 16 |
| Figure 2.10: | Pileup distribution in 2018. | 26 |
| Figure 3.1: | Diagrams for gluino and squark pair production, as predicted by supersymmetric extensions of the Standard Model. | 31 |
| Figure 3.2: | Diagram for the mono- ϕ model. | 35 |
| Figure 3.3: | Leptoquark pair production diagrams. | 36 |
| Figure 3.4: | M_{T2} distributions of backgrounds and an example signal point. | 39 |
| Figure 3.5: | $\Delta\phi_{min}^{1,2,3,4}$ distributions of backgrounds and an example signal point. | 40 |
| Figure 3.6: | (Left) The ratio of the number of same flavor to different flavor lepton pairs in data. (Right) A comparison of the M_{T2} shape between data Z to lepton events and simulated Z to neutrino events. | 44 |
| Figure 3.7: | Fit of r_ϕ as a function of M_{T2} | 47 |
| Figure 3.8: | Example plot of jet energy resolution template. | 49 |
| Figure 3.9: | Validation of the Rebalance and Smear jet mismeasurement background estimate in the $\Delta\phi_{min}^{1,2,3,4} < 0.3$ control region. | 50 |
| Figure 3.10: | Comparison of predicted background and observed data events in the classic search (upper) integrated over M_{T2} and (lower) for each of the medium H_T search regions. | 54 |
| Figure 3.11: | Exclusion limits for gluino pair production and decay to light quarks. | 57 |
| Figure 3.12: | Exclusion limits for gluino pair production and decay to bottom (left) and top (right) quarks. | 58 |
| Figure 3.13: | Exclusion limit at 95% CL _S for (upper left) light-flavor squark pair production, (upper right) bottom squark pair production, and (lower) top squark pair production. | 59 |
| Figure 3.14: | Exclusion limit at 95% CL _S for top squark pair production for various decay modes of the top squark. | 60 |
| Figure 3.15: | Exclusion limit at 95% CL _S for the mono- ϕ model. | 61 |
| Figure 3.16: | Upper limits at 95% CL _S on the leptoquark production cross sections as a function of leptoquark mass. | 62 |

| | |
|--|----|
| Figure 3.17: Lifetime of $\tilde{\chi}^\pm$ as a function of mass splitting. | 66 |
| Figure 3.18: Diagrams for (left) gluino, (center) light-flavor squark, and (right) top squark pair production, in which the gluinos and squarks can decay via a long-lived $\tilde{\chi}^\pm$ | 66 |
| Figure 3.19: Comparison of signal and background Short Track relative isolation distributions. | 68 |
| Figure 3.20: Signal efficiency with respect to the short track selection. | 73 |
| Figure 3.21: Validation of the disappearing tracks background estimate in (upper) 2016 and (lower) 2017–2018 data. | 76 |
| Figure 3.22: Comparison of the predicted background and observed data events in the disappearing track search for (upper) 2016 and (lower) 2017–2018. | 79 |
| Figure 3.23: Exclusion limits at 95% CL _S for gluino pair production, in the disappearing tracks search. | 82 |
| Figure 3.24: Exclusion limits at 95% CL _S for light squark pair production, in the disappearing tracks search. | 83 |
| Figure 3.25: Exclusion limits at 95% CL _S for top squark pair production, in the disappearing tracks search. | 84 |
| Figure 3.26: Exclusion limits at 95% CL _S on the $\tilde{\chi}_1^0$ mass as a function of $\tilde{\chi}^\pm$ proper decay length for gluino pair production and decay to light quarks, and light squark pair production, in the disappearing tracks search. | 85 |
| Figure 3.27: Exclusion limits at 95% CL _S on the $\tilde{\chi}_1^0$ mass as a function of $\tilde{\chi}^\pm$ proper decay length for top squark pair production, in the disappearing tracks search. | 86 |
| Figure 3.28: Exclusion limits at 95% CL _S on gluino, light squark, and top squark pair production cross sections as a function of $\tilde{\chi}^\pm$ proper decay length, in the disappearing tracks search. | 87 |

LIST OF TABLES

| | | |
|------------|--|-----|
| Table 3.1: | Table of systematic uncertainties affecting expected signal yields. | 36 |
| Table 3.2: | Table of uncertainties affecting the lost lepton background estimate. | 41 |
| Table 3.3: | Table summarizing the systematic uncertainties affecting the $Z \rightarrow vv$ background prediction. | 45 |
| Table 3.4: | Summary table of baseline event selection. | 45 |
| Table 3.5: | Systematic uncertainties affecting the Rebalance & Smear jet mismeasurement background estimate. | 51 |
| Table 3.6: | Table of ST and STC selections. | 72 |
| Table 3.7: | Table of uncertainties affecting the disappearing tracks background prediction. | 77 |
| | | |
| Table A.1: | Table of the classic monojet regions. | 92 |
| Table A.2: | Table of the Very Low H_T classic regions. | 93 |
| Table A.3: | Table of the Low H_T classic regions. | 94 |
| Table A.4: | Table of the Medium H_T classic regions, with $N_{\text{jet}} < 7$ | 95 |
| Table A.5: | Table of the Medium H_T classic regions, with $N_{\text{jet}} \geq 7$ | 96 |
| Table A.6: | Table of the High H_T classic regions, with $N_{\text{jet}} < 7$ | 97 |
| Table A.7: | Table of the High H_T classic regions, with $N_{\text{jet}} \geq 7$ | 98 |
| Table A.8: | Table of the Ultra-High H_T classic regions, with $N_{\text{jet}} < 7$ | 99 |
| Table A.9: | Table of the Ultra-High H_T classic regions, with $N_{\text{jet}} \geq 7$ | 100 |

ACKNOWLEDGEMENTS

Thousands of physicists, engineers, and technicians around the world have worked for decades to design, build, and operate CERN’s LHC and the CMS detector. My work would have been impossible without their efforts.

My advisors Avi Yagil and Frank Würthwein showed great patience and faith in me along the way, for which I am profoundly thankful.

Mario Masciovecchio deserves special thanks for providing the bulk of my technical training, and working with me at all hours of the day and night, transcending timezones, for years. He also developed the prototype short track definition.

Slava Krutelyov, Ryan Kelley, Ian MacNeill, Bobak Hashemi, Dominick Olivito, Mark Derdzinski, Giovanni Zevi Della Porta, and Daniel Klein also provided helpful training.

Bennett Marsh, as part of and in addition to maintaining the “classic” portion of the analysis, wrote a lot of helpful code, offered many useful tips and ideas, and in general managed much of the tedious technical work for years.

Claudio Campagnari provided useful instruction in statistical analysis, and suggested the idea that finalized the disappearing track background estimate.

More generally, I must thank the rest of the Surf and Turf (SNT) group for criticism and suggestions along the way, which helped to sharpen the analysis, and the team responsible for maintaining the CMS Tier 2 computing center at UCSD, which provided most of the computing resources.

Finally, the CMS SUSY group, like SNT, provided useful criticism during analysis development.

VITA

2013 B. A. in Physics, Williams College

2019 Ph. D. in Physics, University of California, San Diego

PUBLICATIONS

The CMS collaboration, Sirunyan, A.M., et al., *Searches for physics beyond the standard model with the M_{T2} variable in hadronic final states with and without disappearing tracks in proton-proton collisions at $\sqrt{s} = 13$ TeV*, Eur. Phys. J C (2019)

ABSTRACT OF THE DISSERTATION

A Search for New Physics producing Jets, Large MT₂, and Disappearing Tracks in 13 TeV Proton-Proton Collision at CERN's Large Hadron Collider

by

Dylan Gilbert

Doctor of Philosophy in Physics

University of California, San Diego, 2019

Professor Avraham Yagil, Chair
Professor Frank Würthwein, Co-Chair

This work presents two searches for new physics characterized by pair-production of strongly interacting particles, each decaying to hadronic jets and a particle that is not detectable. The searches use the full 13 TeV proton-proton collision dataset produced by CERN's Large Hadron Collider and recorded by the CMS detector from 2016 to 2018, with total integrated luminosity 137 fb^{-1} . The presence of particles interacting too weakly to be detected is inferred using imbalance in the transverse momentum of the collision products, and sensitivity to pair-production is enhanced by requiring large values of the kinematic variable M_{T2} in events with

at least two jets. The first search is inclusive, binning events using the total hadronic transverse energy, the total number of jets, the number of jets reconstructed as originating from a bottom quark, and either the value of M_{T2} in multijet events, or the transverse momentum of the jet in monojet events. The second search extends the first, by requiring the presence of a disappearing track in the event, and adds binning in the length and transverse momentum of the disappearing track. Both searches are sensitive to a variety of extensions to the Standard Model that include dark matter candidates. Of greatest interest, the results set constraints on pair production of squarks and gluinos as predicted by R-parity conserving supersymmetric extensions of the Standard Model, in which the lightest superysmmetric particle is a neutralino. The first search is sensitive to any decay chain terminating in Standard Model hadrons plus the neutralino, while the second specifically targets, with greatly enhanced sensitivity, decay chains containing an intermediate long-lived chargino. These constraints are the most stringent yet produced by any experiment.

Chapter 1

Introduction

Introduction

1.1 The Standard Model of Particle Physics

Briefly discuss the history of particle physics (eg Millikan, Rutherford, Planck, Bohr, Dirac... QFT), culminating in the Standard Model.

Figure ??

1.1.1 A Brief Description

Describe the particles of the Standard Model, and some of their experimental relevance.

1.1.2 Feynman Diagrams

Provide a few example Feynman diagrams (eg. ee -> mu mu, gg -> top loop -> higgs -> top loop -> gamma gamma) and describe how they represent particle physics processes, and are used to calculate amplitudes.

1.1.3 Quantum Chromodynamics, Hadronization, and Jets

Describe why final state quarks and gluons experimentally produce jets. Mention that it is difficult to make precise predictions in QCD (at low energy), leading to requirement for approximate models for jet formation (pythia...). In particular, ISR is challenging.

1.1.4 Parton Distribution Functions

Describe protons as a sea of quarks and gluons with varying momenta, and why we need a PDF to calculate production rates in a hadron collider.

1.2 Some Problems with the Standard Model

Standard Model is extremely successful (electron magnetic moment, Higgs prediction) but not entirely satisfactory.

1.2.1 Effective Field Theory

Includes explicit energy cutoff, as part of renormalization. Cannot, by construction, be the final theory.

1.2.2 Gravity

Doesn't include gravity, so Planck scale is maximum possible cutoff (but could be smaller).

1.2.3 The Hierarchy Problem

Why is the Higgs mass / weak scale so much smaller than the Planck scale? Is the cutoff much smaller?

1.2.4 Astrophysical Evidence of Dark Matter

SM has no (cold) dark matter candidate. Of course may be well out of reach, but look where we can. Good reason to believe that dark matter could be produced at LHC (WIMP miracle).

1.3 Supersymmetry

SUSY is one particular model that can patches up some of the SM's issues.

1.3.1 Theoretical Appeal

Solves up hierarchy problem if at the weak scale (so, potentially accessible to LHC) and provides a dark matter candidate. (Focus on RPC SUSY, due to proton stability.)

1.3.2 Experimental Signatures

SUSY produces MET, and in strong SUSY decays, potentially lots and lots of jets. Since SUSY

1.3.3 Simplified Models

SUSY parameter space is vast, so we consider one potential discovery channel at a time.




Figure 1.1: In simplified models, it is possible to calculate superpartner production cross sections from theory. Here, the theoretical gluino pair production cross section in 13 TeV proton-proton collisions is shown in black, with the one standard deviation uncertainty shown as a yellow band. The cross section drops rapidly with increasing mass. Based on cross section values used in [11], calculated in [9]. Compare Figure 5 (upper) from [9], of which this is a simplified reproduction.

1.4 Other Models

Introduce leptoquarks and mono-*phi*.

Chapter 2

The Large Hadron Collider and the CMS Detector

The searches for new physics discussed in the next chapter analyze data recorded by the Compact Muon Solenoid (CMS) experiment. The CMS detector itself straddles beams generated by the Large Hadron Collider (LHC), a particle collider located at CERN in Geneva, Switzerland. CMS directly measures the products of collisions between the protons in these beams, digitizes the measurements, and records a small subset of these events. The data is then processed by reconstruction algorithms and shared across the worldwide CMS computing grid. This chapter describes the LHC and CMS.

2.1 The Large Hadron Collider

A detailed description of the LHC is available in Reference [8]. This section provides a summary of the essential information.

2.1.1 A Brief Description

Although it sometimes collides atomic nuclei, typically Pb-208, the LHC is primarily a proton-proton collider, and that is the operational mode we shall consider in this work. The LHC is the highest energy and highest luminosity proton-proton collider ever constructed, thus far achieving collisions with center of momentum frame energy as large as 13TeV , produced by two counterrotating beams of 6.5TeV protons, at sustained rates on the order of one billion collisions per second.

Producing such high energy collisions at such high intensity requires that the LHC be a large ring, 27 km in circumference. The circular shape allows multiple opportunities for the protons in each beam to collide, rather than a single all-or-nothing pass, and additionally allows the beams to be circulated for as long as necessary to accelerate them up to the intended energy, at a rate of 0.5 MeV per revolution. The size makes it possible for the protons to be constrained to travel in a circle, even at very high energies. The beams are constrained to travel in the ring using powerful magnetic fields, and the maximum feasible magnetic field strength is the primary limit on the maximum achievable beam energy. The LHC beams require an 8.36 T field to stay on track, provided by superconducting niobium-titanium electromagnets carrying a current of $11,080\text{A}$. Superconductors stable at greater magnetic field strengths are rare and expensive, and the magnets must be superconducting to carry such enormous currents. Larger rings reduce the beams' curvature and allow for greater energies with the same magnetic field strength.

Protons are chosen for the beams rather than electrons, the only other obvious potential choice, for two reasons. First, protons are composite particles, so their total energies are distributed among their many subcomponents, referred to as partons in this context. Collisions between protons with energy 6.5 TeV each are in fact collisions between a parton in each proton with some fraction of the total energy. This naturally scans across the accessible energy range, providing sensitivity to any new state with mass on the TeV scale. Were the beams composed of elementary particles like electrons, the collider would need to scan the beam energy to provide sensitivity to

any possible mass. Therefore, electron colliders are poorly suited for new discoveries compared to proton colliders, and better suited for studying a state with known mass in greater detail. Second, and more importantly, the mass of the proton is much greater than that of the electron, reducing energy losses to synchrotron radiation. Because synchrotron losses are relatively small, circular colliders of protons are limited chiefly by the magnetic field strength, as discussed above. Electron colliders, on the other hand, are limited by synchrotron losses. Illustrating this, the Large Electron Positron (LEP) collider previously occupied the same tunnel that now houses the LHC, and managed a maximum collision energy of only 209 GeV ??.

The protons are not collided one-by-one, as this would be impossibly tedious. Instead, they are grouped in bunches of over 100 billion, spaced by 25 ns to make it possible for the detectors to distinguish the products of one bunch crossing from those of the preceding and following crossings. During each bunch crossing, a few dozen proton pairs (see Figure 2.10) collide simultaneously, of which either one or none are interesting and the rest are undesirable noise called pileup, discussed in Section 2.3.1. These bunch crossings occur at a few points around the LHC tunnel occupied by detectors. Among them are the CMS detector, described in Section 2.2, and its sister detector ATLAS, which are both general-purpose detectors designed to study any and all phenomena at the TeV scale, and operate in parallel to provide independent checks of any discoveries.

2.1.2 Luminosity Delivered

The sensitivity of searches performed in CMS scales straightforwardly with the total number of collisions provided by the LHC. Thus far, the LHC has provided CMS with nearly 200 fb^{-1} across all operating periods, with the rate accelerating as the machine is pushed to the limits of its design, as shown in Figure 2.1. The searches discussed in the next chapter use only the 137 fb^{-1} recorded by CMS during the 13 TeV runs from 2016 to 2018. As the total inelastic proton-proton collision cross section is approximately 69.2 mb at 13 TeV, the total number of




Figure 2.1: The integrated luminosity delivered to CMS over time by the LHC, in blue, and the portion CMS managed to record, in yellow. As can be seen, CMS is not 100% efficient. It is important to note that this inefficiency is a straightforward failure to observe the events, *not* the purposeful rejection of events by the trigger system discussed in Section 2.3.1. Taken from [2].

collisions observed by CMS from 2016 to 2018 is nearly 10^{16} . While this vast sample size makes very rare genuine physics processes potentially observable, it also presents a challenge in that very rare and difficult to model detector failures may pollute the dataset. It is prudent to adopt a conservative approach to any searches for rare new physics.

2.2 The CMS Detector

While the LHC provides the collisions, it is the Compact Muon Solenoid (CMS) detector that ultimately measures the products and reconstructs each event. A detailed description of all of the CMS hardware and software is available in Reference [7], and a description of the general event reconstruction technique used at CMS, called Particle Flow, is available in Reference [32].

This section serves as a general overview of how CMS works, sufficient to understand the searches for new physics presented later in this work.

The particles stable enough to be detected directly by CMS come in 5 distinct varieties, namely electrons, muons, photons, charged hadrons (largely protons, pions, and charged kaons), and neutral hadrons (largely neutrons and neutral kaons). Figure 2.2 depicts a generic detection of each kind of particle. All other particles are too unstable to be measured directly, except the neutrinos. Neutrinos are, uniquely among Standard Model particles, both stable and invisible to CMS. Their presence in the products of a collision is inferred by measuring all of the visible content of an event, and noting that some energy is missing, as discussed in Section 2.2.6. Any stable weakly interacting particles beyond the Standard Model produced in a collision at CMS would present a similar signature.




Figure 2.2: The design of CMS causes particles of different types to leave distinct experimental signatures, allowing them to be distinguished. The rest of this section references and explains this image. Taken from [32].

The CMS detector is a cylinder composed of a central barrel section with two flat endcaps,

each of which contain layers of different detection systems designed to provide the information necessary to distinguish the various particles that can be generated by collisions, and measure their positions and energies. Starting from the beam pipe, they are the tracker, the electromagnetic calorimeter, the hadronic calorimeter, and the muon system. The tracker and the calorimeters are inside a solenoid producing a magnetic field, while the muon system is subject only to the external, returning field of the solenoid. Altogether, the detector measures about 7.5 meters in radius and 21 meters in length. As suggested by the cylindrical detector shape, the experiment uses a cylindrical coordinate system, in which the beam axis is assigned to the z axis, and the angle in the transverse plane, ϕ , is measured starting from a line connecting CMS to the center of the LHC ring.

This section describes each of the subsystems, as well as the objects that are constructed using their measurements.

2.2.1 Magnet

As its name suggests, a major feature of the CMS detector is a superconducting solenoid, the largest ever built, which provides a constant 3.8 T magnetic field in the tracker and calorimeters. This magnetic field curves the trajectories of charged particles, allowing for a determination of their charges and assisting in the measurement of their masses and energies, contributing to highly accurate identification of particles.

CMS is compact in part to make it possible to fit the bulk of the detector inside this single solenoid. This maximizes the strength and consistency of the magnetic field, which in turn maximizes the precision with which particle positions, and the curvatures of their trajectories, can be measured.

In Figure 2.2, one can see the path of an electron (orange) curving sharply, in a direction indicating that it is negatively charged. Similarly, the path of a charged hadron (solid green) curves in the opposite direction, indicating a positive charge. A positively charged muon (blue)

curves first one way, then the reverse once the field flips after exiting the solenoid. Neutral particles (dotted lines) do not curve at all.

2.2.2 Tracker

The silicon tracker is the innermost detector system and is primarily responsible for measuring the positions of charged particles. In Figure 2.2, the trajectories of neutral particles are represented with dotted lines, to indicate that they do not actually leave those tracks in the detector. The tracker is critically important for the disappearing tracks search covered in Section 3.2, and so is described here in more detail than the other detector systems.




Figure 2.3: A slice of the CMS tracker in the $r - z$ plane, with the inner pixel tracker (red) in the 3-layer configuration used until 2016. In 2017 and 2018, the pixel tracker was upgraded with a 4th layer. The tracker is symmetric about the beam axis ($r = 0$); one should imagine rotating this image around the horizontal axis to form a cylinder, to visualize the tracker’s 3-dimensional form. The labels in green text denote tracker subsystems, the Tracker End Cap, the Tracker Inner Disk, the Pixel Tracker, the Tracker Inner Barrel, and Tracker Outer Barrel. The interior subsystems have more miniaturized detector elements and, therefore, better resolution. Taken from [13].

As shown in Figure 2.3, the tracker is composed of several discrete layers. When a charged particle passes through a layer, it usually causes a measurable current to flow in the silicon, and is said to have produced a “hit.” As shown in Figure 2.4, the probability for a genuine particle to leave a hit is near but not quite unity, so a missing hit where one would be expected is




Figure 2.4: The efficiency for a genuine charged particle to produce an expected hit in the various components of the CMS pixel tracker, in the configuration used in 2016 and earlier. Fake tracks may have many missing expected hits, but more than one or perhaps two is extremely rare in tracks produced by genuine particles. The probability for a neutral particle to produce a hit is small, and the probability to leave enough to produce a track is negligible. Taken from [13].

credible but cause for skepticism. Each event produces an intimidating array of these discrete hits throughout the tracker, which must then be connected by the track reconstruction algorithm into particle tracks. CMS uses an iterative procedure described in Reference [13], in which clean, high energy, isolated tracks are reconstructed first so that their hits may be removed from the pool, followed by the next most obvious tracks, and so on, dramatically simplifying the computational complexity of each step. The early iterations that produce the tracks judged to be of highest quality find only those tracks with transverse momenta at least 0.3 GeV, and no iteration can produce a track with transverse momentum below 0.1 GeV, so that particles of such low energy are effectively invisible to CMS. Note that, due to the strong magnetic field present in the tracker, particles with energies lower than around 0.1 GeV travel in such tight spirals that reconstruction

becomes infeasible. Although this is normally insignificant, since these particles are not typically important to an event’s overall characterization, this fact allows for the existence of disappearing tracks, in which a high energy charged particle decays to a trackless neutral particle and a charged particle of such low energy that it is not reconstructed. Additionally, all tracks must be composed of at least 3 hits.

The performance is outstanding, especially for tracks passing the sophisticated “high purity” selection applied by the vast majority of CMS analyses. Figure 2.5 shows the fake rate of this procedure, less than 5% for tracks in the intermediate energy range, and further suppressible with additional cleaning selections like those of the disappearing tracks search covered in Section 3.2. Figure 2.6 shows the efficiency to reconstruct tracks successfully for the three most common varieties of charged particles measured by CMS, namely muons (left), electrons (center), and pions (right), approaching unity in all cases for higher energy tracks in the barrel region of the tracker.

The tracker detection elements are smaller and consequently provide better localization in interior layers of the tracker, with the pixel detector components only approximately 100 μm across. As a general guideline, no element in the tracker is permitted to have an expected probability of a hit greater than 1% per bunch crossing, to maintain a low probability of tracks intersecting. This structure produces outstanding sub-millimeter resolution of a track’s origin, shown in Figure 2.7, allowing tracks originating from collisions only microns apart to be separated, and tracks originating from decays of particles with decay lengths on the order of a millimeter to be identified. The former is important for rejection of pileup, and the latter for identification of jets originating from bottom quark hadrons.

As a general rule, the tracker intends to interact with particles only to the extent absolutely necessary to measure their positions. Strong interactions between the tracker and the particles it is measuring modify the particles’ tracks and affect their energies, and are therefore undesirable. Unfortunately, electrons are so strongly affected by bremsstrahlung that the tracker is quite thick




Figure 2.5: The CMS track fake rate as a function of p_T , in the configuration used in 2016 and earlier. The sophisticated high purity selection aims to suppress the fake rate without badly affecting efficiency and is applied in the vast majority of CMS analyses. Taken from [13].




Figure 2.6: The CMS track reconstruction efficiency for (left) muons, (center) electrons, and (right) pions, split by detector region as a function of p_T , in the configuration used in 2016 and earlier. Muons are by far the cleanest objects at CMS as they are massive enough to avoid strong perturbations by tracker material, and lack nuclear interactions. Electrons are lighter and so are vulnerable to tracker interactions, experiencing strong bremsstrahlung losses, while pions can undergo nuclear interactions. Taken from [13].




Figure 2.7: The CMS tracker has micron-level impact parameter resolution, for both (left) the transverse plane and (right) the z direction. Solid markers indicate the half-width of the 68% confidence interval, while open markers indicate 90% confidence. With this resolution, a track reconstructed with an impact parameter of more than a few hundred microns is likely a fake, from a pileup vertex, or from a secondary vertex produced by a decaying particle like a bottom hadron. Taken from [13].




Figure 2.8: The CMS tracker material budget in (left) radiation lengths (ie electromagnetic) and (right) interaction lengths (ie nuclear), in the configuration used in 2016 and earlier. While the CMS tracker is not intended to shower particles, only to measure positions as gently as possible, some amount of interaction between tracker material and collision products is inevitable. Taken from [13].

in terms of radiation lengths, as shown in Figure 2.8 (left). Therefore, electrons at CMS are experimentally observed more as clouds of bremsstrahlung photons by the time they reach the electromagnetic calorimeter at CMS, and must be reconstructed with a special tracking procedure,




Figure 2.9: The probability for pions to undergo nuclear interactions in the CMS tracker by layer, in the configuration used in 2016 and earlier. In rare cases, the pion may shower in such a way that no charged daughters have high enough energy to be reconstructed, causing its track to disappear. Taken from [13].

the Gaussian-sum filter described in [5]. In terms of interaction lengths, the CMS tracker is much thinner, as shown in Figure 2.8 (right). The vast majority of hadrons therefore reach the calorimeters without having undergone a nuclear interaction. Figure 2.9 shows that over 80% of high energy charged pions reach the calorimeters intact. Even so, 80% is not 100%. Pion tracks occasionally terminate in a shower induced by a collision with a nucleus inside the tracker. In rare cases, these showers can contain *zero* visible particles of any kind. These disappearing pions constitute an important background for the disappearing tracks search.

2.2.3 Electromagnetic Calorimeter

The primary purpose of the electromagnetic calorimeter (ECAL) is to observe the energies of photons, which do not leave tracks. It is also important in the observation of electrons, which produce a jet of photons due to bremsstrahlung in the tracker. Other electromagnetically interacting particles will also leave small ECAL deposits, but much less due to their lesser tendency to produce photons. In Figure 2.2, the electron (red) and photon (dotted blue) produce strong showers in the ECAL (green), while the other particles leave negligible deposits.

The design prioritizes good resolution of photon energies and positions over all else, due to the experimental importance of the Higgs boson's decay to a pair of isolated photons. To accomplish this, the ECAL is constructed from tens of thousands of scintillating lead tungstate crystals, which has several desirable properties.

First, these crystals have a Moliere radius of only 2.2 cm [7]. Aside from providing generally good position resolution, this small Moliere radius most importantly allows a prompt isolated photon, perhaps from a Higgs decay, to be distinguished from two closely overlapping photons produced by the decay of a neutral pion, because the two separate showers produced by the photon pair are clearly separated in the ECAL cells. These two nearby showers may merge into a single large shower indistinguishable from that of a single photon in a material with a larger Moliere radius. Similarly, photons produced inside jets can be distinguished from truly isolated photons, since the showers of other particles in the jets can be discriminated from the photon's own shower. Although the size of the shower is restricted, a typical one will still extend over a few adjacent crystals as shown in Figure 2.2, so that a 5-by-5 cluster of crystals, or potentially a smeared supercluster for electrons that have radiated over a larger area, is the essential object of ECAL reconstruction. Studying the distribution of energy measurements across the crystals in a cluster, as described in [10], allows for better fake rejection and energy resolution of 1% [?].

Second, lead tungstate also has an extremely short radiation length, only 0.89 cm compared to the roughly 2 radiation lengths of the entire 1 m radius tracker, so it contains electrons very

efficiently [7].

Third, scintillation in lead tungstate happens quickly, with 80% of the light produced within 25 ns [7]. This speed is critical for coping with the 25 ns bunch spacing of the LHC’s proton beams, so that the ECAL deposits produced by adjacent bunch crossings can be distinguished. Due to this rapid shower production and careful calibration of the electronics, the ECAL achieves a timing resolution of roughly 0.2 ns for particles with momenta on the order of tens of GeV [15].

Finally, lead tungstate is resistant to radiation damage, so that the ECAL can continuously operate in the high flux environment of LHC proton collisions. Unfortunately, it is not immune to radiation damage, and so the performance of the ECAL degrades over time. By the end of data taking in 2018, roughly 20% of ECAL cells no longer consistently detected showers of photons and electrons. This leads to a background for the disappearing tracks search consisting of electrons that transform to photons inside the tracker, no longer leaving a track, but are not later detected as photons in the ECAL. Fortunately, 80% of the ECAL is still performing well despite years of high radiation flux, so this background can be almost entirely rejected by mapping and vetoing the poorly performing cells, using events in which a Z boson decayed to a pair of electrons, exactly one of which is lost due to this effect, as described in Section 3.2.

2.2.4 Hadronic Calorimeter

Charged hadrons are too massive to shower significantly in the ECAL, and neutral hadrons are invisible even to the tracker. However, both can be made to shower by capitalizing on their powerful nuclear interactions, and the hadronic calorimeter (HCAL) accomplishes this, as depicted in Figure 2.2 for a neutral hadron in dotted green and a charged hadron in solid green. The performance of the HCAL is very important to analyses selecting events containing jets and missing energy, as HCAL measurement errors lead directly to errors in jet energies, which in turn cause errors in the inferred missing energy in the event.

The essential plan of the CMS HCAL design is to induce hadron showers by putting a

high density of atomic nuclei in their paths in the form of 5 cm brass plates, then to measure the products of these showers using plastic scintillators, and from this reconstruct the energies of the original hadrons. To ensure that as few hadrons as feasible escape detection, whether from the primary interaction or from secondary showers, there are multiple layers of this shower-inducing metal plate and scintillator combination. Nevertheless, some hadrons do manage to “punch through” the HCAL, especially if they have extremely high energy, and even successful measurements are only accurate to within roughly a factor of 2 [14], unless that hadron has very high energy. On the whole, the CMS HCAL performance is the weakest of any detector system. However, the energy of a typical jet is not measured only with the HCAL, and errors made on different hadrons within a jet tend to cancel, so that CMS achieves a typical jet energy resolution of approximately 10%, for the jets of most interest [7]. Still, jets are occasionally badly mismeasured, most often due to HCAL measurement errors, and it is important for analyses selecting events with large missing energy to filter out these mismeasurement events, see Section 3.1.3.

Accuracy of energy measurements aside, the ability to detect neutral hadrons consistently in the HCAL is important to the disappearing tracks search discussed in Section 3.2. A charged hadron can decay to a neutral hadron inside the tracker, and thus allow known Standard Model physics to produce an apparent disappearing track. Unlike a true disappearing track, however, the neutral hadron is detectable in the HCAL, revealing what happened in the tracker and rejecting this Standard Model background. As with the ECAL, poorly performing regions of the HCAL can be vetoed to ensure this background is wholly rejected.

2.2.5 Muon System

The muon system is the outermost detector of CMS, and the only one outside the solenoid. Muons, like charged hadrons, are too heavy to produce significant showers in the ECAL, and unlike hadrons, do not have nuclear interactions allowing them to be trapped in the HCAL. Instead, they penetrate through the entire detector, leaving a track in the tracker, largely disappearing

inside the calorimeters, and then reappearing in the muon system. Unlike other particles, then, the energies of muons must be measured largely by analyzing the curvature of their tracks. As one of the design goals of the Compact Muon Solenoid experiment is ultra-precise measurements of muons, the muon system is large partly in order to provide a long look at the muon’s track for momentum measurement purposes, and partly to provide a positive identification of a particle as a muon, rather than a hadron that managed to escape the HCAL, as only muons can penetrate through so many layers of detector material. As installing a second tracker would be far too expensive and excessive, the muon system instead uses only four widely spaced layers of detectors operating based on ionization of gas by passing muons, which while generally worse than the tracker system in terms of space and time resolution, are much more economical for covering such a large volume [7]. The design of CMS has proven successful. Muons are by far the cleanest, most well-measured particles at CMS, with momentum resolution better than 1% and high efficiency, as we have already seen for tracker muon tracks in Figure ??.

While muons are the only charged Standard Model particle that can consistently penetrate to the muon system, the possibility exists that heavy long-lived charged particles beyond the Standard Model may be produced in LHC collisions and leave hits in the muon system. The disappearing tracks search in Section 3.2 targets models that could potentially produce such a signature. However, that search considers only tracks that disappear inside the tracker, and in fact explicitly vetoes muon-like tracks, so it is not sensitive to this potential signature. Adding sensitivity to this powerful signature could be an interesting extension in the future.

2.2.6 Missing Transverse Energy

Once muons are measured in the muon system, all Standard Model particles have been accounted for, except neutrinos. Unfortunately, the neutrino interaction length, even at LHC energies, is on the order of billions of meters at Earthly densities, so that the probability of a typical LHC neutrino experiencing even a single interaction anywhere inside the CMS detector is

negligible [16]. However, the presence of neutrinos can be inferred by measuring all other particles inside the event, and observing that the transverse momentum does not balance, indicating either the presence of at least one undetected particle, or a detector mismeasurement. At a hadron collider like the LHC, only the transverse momentum can be used for this procedure, because the exact center of momentum frame of the primary collision is unknown along the beam axis. The actual colliding particles are not the protons themselves, but rather subcomponents of the proton called partons. The partons are approximately at rest with respect to the detector in the transverse plane, so that their transverse momenta are known to sum nearly to zero, but have unknown momenta along the beam axis. Unbalanced momentum along this axis may indicate the presence of an invisible particle, but more often it indicates simply that the initial momentum of the collision along the beam axis was nonzero. The observable is not the missing energy, then, but the missing transverse energy, written as MET or \cancel{E}_T . This observable is tentatively identified with the unobservable transverse momentum of a neutrino.

Although the invisible momentum is only available in the transverse plane, it is still a very useful quantity. For instance, consider the leptonic decay of a W boson, $W^\pm \rightarrow \ell^\pm v$. The W boson has mass approximately 80 GeV with width approximately 2 GeV [36]. Even allowing for detector resolution effects around 10%, a lepton and neutrino system originating from an on-shell W decay can never have mass greater than approximately 100 GeV. It follows directly that transverse mass M_T , the mass calculated using energy and momentum in the transverse plane, also cannot exceed 100 GeV, since $E_T < E$ and $p_T < p$. The analyses discussed in the next Section ?? leverage this fact to determine whether a lepton or disappearing track may potentially have originated in a leptonic W decay.

Of course, while the missing transverse energy is sometimes the undetected energy of a single neutrino or unknown invisible particle, it can also be a consequence of mismeasurement, or even potentially the vector sum of the transverse momenta of *multiple* undetected particles. In fact, even events with relatively large amounts of missing energy at CMS are most often produced

by detector mismeasurement (see Figure 3.4, in which “Multijet” events in yellow have \cancel{E}_T due to detector mismeasurement), even though such large errors are rare, because neutrinos are rarer still. Searches for invisible particles naturally wish to eliminate these events with fake \cancel{E}_T using cleaning selections. One of the most powerful observables that can be used to reject events with fake \cancel{E}_T is M_{T2} [22], a generalization of M_T .

M_{T2}

Although the presence of invisible particles can be inferred via the observation of a nonzero \cancel{E}_T vector, it is not trivial to distinguish events in which the \cancel{E}_T corresponds to the transverse momentum of a single invisible particle, and events in which the \cancel{E}_T is the vector sum of the momenta of multiple invisible particles. It can also be difficult to distinguish the presence of genuine invisible particles from cases of detector mismeasurement. While no conclusions about invisible content can ever be drawn with perfect certainty for any single event, the situation is not hopeless. The M_{T2} observable is one tool used to make quantitative inferences concerning an event’s invisible content. Specifically, M_{T2} is designed to identify events consistent with the presence of two invisible particles in the final state, produced in symmetric decay chains of pair-produced heavy particles. This also tends to reject events in which the \cancel{E}_T was produced by detector mismeasurement, as these events tend to have no symmetry whatsoever.

The algorithm begins by dividing the visible portion of the event into two hemispheres. To do this, an event must have at least two visible objects, so in an event with only hadron jets, like the majority of the events at CMS, M_{T2} can be calculated only for multijet events. The method used to assign each visible object to a hemisphere varies. One popular method, and the one used in [11], begins by identifying the pair of objects with largest system mass, M_{12} ,

$$M_{12} = \sqrt{(E_1 + E_2)^2 - (\vec{p}_1 + \vec{p}_2)^2}, \quad (2.1)$$

and assigning each as the seed of a separate hemisphere. In an event containing two identical pair-produced particles undergoing similar decay chains, the pair of objects with largest mass is unlikely to originate from the same particle, as the mass of the full system is strictly greater than the mass of either particle.

Then, each object is associated to the seed of lesser Lund distance, D_L , which is defined for seed i as [33, 34],

$$D_L = (E_i - p_i \cos \theta) \frac{E_i}{(E_i + E)^2} \quad (2.2)$$

where E_i is the seed's energy, p_i is the seed's momentum, θ is the angle between the momenta of the seed and of the object under consideration, and E is that object's energy. Considering the massless limit of D_L provides some intuition,

$$D_L = \frac{1 - \cos \theta}{\left(1 + \frac{E}{E_h}\right)^2}. \quad (2.3)$$

The Lund distance is small if the hemisphere and object point in nearly the same direction so that $\cos \theta \sim 1$, and if E_i is small. Thus, selecting based on minimal Lund distance prefers to assign objects to the less energetic seed when possible to keep the hemispheres energetically balanced, so long as the object and the seed point in similar directions. Once all objects have been assigned to a seed, two new seeds are defined as the sum of all the four-momenta in each hemisphere, and the assignment procedure repeats using these seeds. This continues until no object changes hemisphere after an iteration. Ultimately, the algorithm assigns objects to the correct hemisphere with efficiency ranging from approximately 70–85%, with higher efficiencies obtained in events with fewer objects and therefore reduced combinatorics [12].

The next step of the calculation of M_{T2} takes these finalized hemispheres as inputs. Under the hypothesis that the observed missing energy is the vector sum of the unobserved momenta of

two invisible particles, the algorithm considers every possible vector decomposition,

$$\vec{p}_T^{\text{miss}} = \vec{v}_1 + \vec{v}_2 \quad (2.4)$$

and calculates M_T for hemisphere i with respect to \vec{v}_i . The maximum of these two values of M_T is taken as the *candidate* M_{T2} for this decomposition. This choice is motivated by a desire to infer the true mass of the decaying pair-produced particle, again, under the hypothesis that there are two pair-produced particles undergoing symmetric partially invisible decays. For the correct \vec{p}_T^{miss} decomposition and hemisphere assignments, the two calculated transverse masses are correct, and both are no larger than the true mass of the pair-produced particles. The larger, of course, is nearer to that true mass.

The only remaining step is to select which of the candidate values of M_{T2} , equivalently which \vec{p}_T^{miss} decomposition, is correct. The algorithm selects the smallest candidate M_{T2} value found, the “minimum maximum,” so that the final M_{T2} is

$$M_{T2} = \min_{\vec{p}_T^{\text{miss}} = \vec{v}_1 + \vec{v}_2} [\max(M_T^1, M_T^2)] \quad (2.5)$$

where hemispheres 1 and 2 are assembled as described above. Since the correct missing energy decomposition is among those considered and manifestly cannot produce a value of M_T greater than the true mass of the parent particle when hemisphere assignments are correct, selecting the “minimum maximum” guarantees that M_{T2} can be no larger than the mass of the parent particle. Just as M_T cannot exceed the mass, M_{T2} is unlikely to exceed the true M_T .

Of course, it is possible for M_{T2} to be small even in genuine signal events for which it is designed to be large, and it is possible for especially pathological mismeasurement events to have large M_{T2} despite M_{T2} being a powerful rejector of these backgrounds (see Section 3.1.3). No observable is so magical that it can accept 100% of signal and reject 100% of background. Still, M_{T2} is one of the most powerful discriminants available for analyses targeting pair-produced

particles decaying semi-invisibly, as can be seen for instance in Figure 3.4. Indeed, the selection of $M_{T2} > 200$ GeV applied by the analyses discussed in Section ?? is the single most important part of their baseline selection, dramatically reducing especially the detector mismeasurement background.

2.3 CMS Event Reconstruction

2.3.1 Challenges

Spray of thousands of particles, many in dense jets, from collisions distributed across a rather large beam crossing, millions of times per second. Pace is so high that travel of electrical impulses along the equipment is comparable to the collision frequency.

The Trigger System

High event rate needs to be suppressed due to computing limitations. High event rate is necessary to see rare things (how rare?).

Pileup

Obtaining a high enough event leads to many simultaneous collisions (not 1 proton collision at a time, but bunches crossing every 25 ns). Charged pileup contribution can be subtracted exactly using the tracker, but neutral pileup correction is heuristic. Effective area vs delta-beta.

2.3.2 Vertices

Use tracker to identify tracks and extrapolate them back to an origin point. These origins are vertices.




Figure 2.10: The distribution in 2018 of the number of simultaneous collisions per bunch crossing, so-called pileup, provided by the LHC and observed by CMS. The average bunch crossing produced 32 simultaneous collisions, with an inferred total inelastic collision cross section of 69.2 mb. Taken from [2].

B-Tagging

Some tracks will not extrapolate back to a proton collision point but instead point a few millimeters away: secondary vertices characteristic of b-tags. Describe other indicators used to identify b-jets and the efficiency.

2.3.3 Particle Flow

Describe particle flow algorithm and advantage relative to other techniques.

2.3.4 Jet Clustering

Jets are made from PF candidates using anti-kt. How accurately?

2.4 Simulation

The Standard Model and models of new physics predict what happens at the primary interaction, and this needs to be converted into a prediction for what happens in the detector.

2.5 Objectives

Used to study backgrounds and develop estimation techniques, and to design analyses to maximize sensitivity to expected signal signatures.

2.6 Limitations and Challenges

Ultra-detailed knowledge of the detector at the subatomic level at all locations is unachievable, and the detector's state evolves over time. QCD is irritating, and even electroweak physics is subject to theoretical uncertainties due partly to dependence on experimental inputs, and partly to calculating only to finite order. Can't run the simulation forever, and the computation is expensive.

2.7 The Simulation Pipeline

Matrix elements (MadGraph), observed particles (pythia), how those particles interact with detector components (GEANT). Mention also fastsim vs fullsim (ie replacing GEANT with approximate smearing).

Chapter 3

The Full 13 TeV M_{T2} Analysis

This section presents two searches for new physics in all-hadronic final states with substantial \cancel{E}_T as inferred through large M_{T2} in 13 TeV proton-proton collisions recorded by the CMS detector [11].

The \cancel{E}_T selection targets invisible particles, motivated by the evidence for dark matter discussed in Section 1.2.4. If dark matter is at least partially composed of a particle or particles associated with physics at the weak scale, then it may be produced in LHC collisions and, while not detectable itself, be inferred via an excess of events with imbalanced transverse momentum.

Considering only all-hadronic events with large M_{T2} is part of a divide and conquer strategy employed by the CMS collaboration. CMS analyses also search for dark matter in events containing leptons, e.g. [30], but there is no way to know whether dark matter will be found in either or both. By splitting the searches, each can optimize for its unique event characteristics. Likewise, the collaboration also considers all-hadronic events that do not necessarily have large M_{T2} [31], since while M_{T2} in many cases provides enhanced sensitivity, it can reduce sensitivity to some models, for instance in cases where the decay energy is very small, or those in which invisible particles are not produced in pairs.

The first search is inclusive, and is a continuation of previous analyses based on smaller

datasets, most recently using the 35.9 fb^{-1} dataset recorded in 2016 [27]. Accordingly, it is referred to as the classic search.

The second search is a new extension of the first, requiring additionally the presence of a disappearing track in a selected event. Disappearing tracks have been targeted as an observable of interest by both CMS [18, 28] and ATLAS [4, 3] previously, but with lower energy or smaller datasets, and different methods.

Both searches set the strongest constraints to date on a variety of hypothetical extensions to the Standard Model possessing pair-produced dark matter candidates, most notably R-parity conserving supersymmetry.

3.1 Classic Search

3.1.1 General Description

The classic search’s core, defining selections are for events with large missing transverse energy \cancel{E}_T as inferred via M_{T2} , large total hadronic transverse energy, called H_T , and no leptons.

The primary motivation for the analysis is either finding or setting constraints on particle dark matter. The experimental signature of dark matter is its undetectability; its presence can only be inferred through large imbalance of transverse energy. The M_{T2} analysis further focusses on the case of pair-produced dark matter using its eponymous observable. For reasons discussed in Section 1.3, R-parity conserving supersymmetry is a model of special interest to the theoretical community, and this model’s dark matter candidate would always appear in pairs. The M_{T2} variable tends to increase sensitivity to the pair-production scenario.

The selection of events with large H_T is motivated by the CMS trigger as discussed in Section 2.3.1, which is in turn motivated by the inferred large energy scale of new physics. The decays of heavy new particles to relatively light Standard Model particles will result in, typically, very energetic events. It is possible that the dark matter candidate is itself very heavy, nearly as

massive as the new particle that decays to it. In this scenario, while the events are indeed very energetic, much of the energy is lost to the invisible portion of the event, and the visible, hadronic energy can be relatively small. Accordingly, sensitivity in the compressed scenario is weaker, as many of the new physics events are too low energy to pass the kinematic selections. The M_{T2} selection is especially inefficient, in this scenario. However, sensitivity is not negligible, and the search is sensitive to any mass splitting in many cases.

The last selection, a veto of leptons, is chiefly motivated by CMS’s divide and conquer strategy. Final states including leptons are considered in other searches. However, the veto does allow for partial suppression of the Standard Model neutrino background, since neutrinos are often produced alongside a charged lepton in decays of the W boson, as discussed in Section 3.1.3.

Ideally, any observation of an event with large \cancel{E}_T and M_{T2} , large H_T , and no leptons, would constitute a discovery. Unfortunately, this is not the case, as the Standard Model is capable of producing events with all of these signatures. Neutrinos and simple mismeasurement of the event can produce large missing energy signatures. Events with large H_T are uncommon in the Standard Model but hardly impossible, and the vast majority of events in a proton-proton collider have no leptons. Some number of events will have all of these properties without any new physics occurring, and most of the analysis’ efforts are spent estimating how often these background processes occur, as discussed in Section 3.1.3.

3.1.2 Signals

The classic search generically targets any new physics that produces high H_T events with jets and missing energy from undetected particles. Diagrams for several candidate models are shown in Figures 3.1, 3.2, and 3.3. The diagrams in Figure 3.1 of gluino and squark pair production as predicted by supersymmetric extensions of the Standard Model are of greatest interest, and while the analysis is sensitive to a wide variety of similar hypothetical models,




Figure 3.1: Diagrams of gluino (upper five) and squark (lower six) pair production, as predicted by supersymmetric extensions of the Standard Model. The classic search considers five potential gluino decay chains. At upper left, the gluinos decay to light flavor quarks (up, down, strange, or charm), and the lightest neutralino, $\tilde{\chi}_1^0$. At upper center, the gluinos decay to light flavor quarks, but rather than decaying directly to $\tilde{\chi}_1^0$, the gluinos decay either to the second neutralino, $\tilde{\chi}_2^0$, which subsequently decays to a Z boson and $\tilde{\chi}_1^0$, or to the lightest chargino, $\tilde{\chi}^\pm$, which subsequently decays to the W boson and $\tilde{\chi}_1^0$, with equal probability. At upper right, both gluinos undergo the $\tilde{\chi}^\pm$ to W decay chain. On the left of the second row, both gluinos decay to bottom quark pairs and $\tilde{\chi}_1^0$. On the right of the second row, both gluinos decay to top quark pairs and $\tilde{\chi}_1^0$. Additionally, the classic search considers six potential modes of squark production and decay. On the left of the third row, light flavor squarks decay to light flavor quarks and $\tilde{\chi}_1^0$. In the center of the third row, bottom squarks decay to bottom quarks and $\tilde{\chi}_1^0$. On the right of the third row, top squarks decay to top quarks and $\tilde{\chi}_1^0$. At lower left, top quarks decay to bottom squarks and $\tilde{\chi}^\pm$, which subsequently decay to the W boson and $\tilde{\chi}_1^0$. At lower center, top squarks may undergo either the $\tilde{\chi}^\pm$ decay chain, or a direct decay to a bottom squark and $\tilde{\chi}_1^0$, with equal probability. At lower right, each top squark decays to a charm squark and $\tilde{\chi}_1^0$. Taken from [11].

it is optimized for these. As discussed in Section 1.3.3, these models simplify the enormous parameter space of supersymmetric extensions by assuming that all superpartners except those in the process are so massive that they can be neglected entirely. The upper five diagrams all depict pair production of gluinos, the supersymmetric partner of the gluon, the mediating gauge boson of the strong interaction. As each gluino decays to two quarks, each of which will typically produce a jet, gluino pair production events tend to have many jets. Many different decay chains are possible, and the analysis considers five representative benchmarks.

In the first benchmark, the gluinos decay to light flavor quarks (up, down, strange, or charm), and the lightest neutralino, $\tilde{\chi}_1^0$.

In the second benchmark, the gluinos decay to light flavor quarks, but rather than decaying directly to $\tilde{\chi}_1^0$, the gluinos can decay either to the second neutralino, $\tilde{\chi}_2^0$, which subsequently decays to a Z boson and $\tilde{\chi}_1^0$, or to the lightest chargino, $\tilde{\chi}^\pm$, which subsequently decays to the W boson and $\tilde{\chi}_1^0$. Each of these decays can occur with equal probability. Both the W and Z will themselves decay, usually to a pair of quarks, which will in turn produce jets. So, relative to the first signal model, the second tends to have greater jet multiplicity. The Z can also decay to neutrinos, and the W can decay to a neutrino and a lepton that is not reconstructed. In these scenarios, this benchmark trades some jets for an enhanced missing energy signature. If the W or Z decays leptonically and a lepton is successfully reconstructed, events from these signals may fail the lepton veto and instead end up in control regions, biasing the background prediction. The procedure used to handle such signal contamination of control regions is described in Section 3.1.6.

The third benchmark, in which both gluinos undergo the $\tilde{\chi}^\pm$ to W decay chain, is similar.

In the fourth benchmark, both gluinos decay to the bottom quark and $\tilde{\chi}_1^0$. As described in Section 2.3.2, it is possible to identify jets that originated from a bottom quark; such a jet is said to be “b-tagged.” The classic search bins in the number of b-tagged jets in order to enhance sensitivity to signals of this type.

In the fifth and final gluino pair-production benchmark, both gluinos decay to top quarks and $\tilde{\chi}_1^0$. The top quark decays with probability near unity to a bottom quark and a W boson. The extra W bosons, compared to the direct bottom decay model, can either add jets or leptons and neutrinos, as previously discussed. This signal tends to produce the most remarkable events of any signal model considered, with very large H_T , N_{jet} , and $N_{b\text{-tag}}$, but also loses many events to the lepton veto as any of the four W bosons is liable to produce a lepton.

The lower six diagrams of Figure 3.1 all depict pair production of squarks, the supersymmetric partners of quarks. Squark decays directly produce only one quark each, compared to two for gluino decays, so squark pair-production events tend to have fewer jets than gluino pair-production events.

In the first squark benchmark diagram, on the left of the third row, a pair of light flavor squarks (up, down, strange, or charm) is produced and each decays to a light flavor squark and $\tilde{\chi}_1^0$.

In the second benchmark, bottom squarks are produced and decay to bottom quarks and $\tilde{\chi}_1^0$. These events tend to have b-tagged jets, but fewer than in gluino decays to bottom squarks.

In the third benchmark, top squarks are pair produced and decay to top quarks and $\tilde{\chi}_1^0$. The relationship between this process and the previous one is similar to the relationship between the fifth and fourth gluino benchmarks, respectively.

The fourth benchmark is very similar to the third. Instead of the top squark decaying directly to a top quark, which then decays to a bottom quark and W boson, the squark decays to a bottom quark directly and $\tilde{\chi}^\pm$, which subsequently produces the W. While the final state particles are identical, their kinematics can be very different depending on the distribution of masses realized in nature. For instance, if the top squark and $\tilde{\chi}^\pm$ mass splitting is very small, the bottom quark jet in this benchmark may have such low p_T in a typical event that it is difficult to reconstruct.

In the fifth benchmark, each top squark may undergo the $\tilde{\chi}^\pm$ decay chain or decay directly to a bottom squark and $\tilde{\chi}_1^0$, with equal probability, mixing the two previous models.

In the last benchmark, each top squark decays to a charm quark and $\tilde{\chi}_1^0$. This decay chain could dominate when the top squark and $\tilde{\chi}_1^0$ mass splitting is too small to allow decay to on-shell top quarks, and the mass of $\tilde{\chi}^\pm$ is larger than that of the top squark.

In each of these models, the mass of the squark or gluino and the mass of $\tilde{\chi}_1^0$ are free parameters. Large squark and gluino masses cause low production rates, as the production cross sections drop rapidly with increasing mass, as discussed in Section 1.3.3 and shown in Figure 1.1. The mass of $\tilde{\chi}_1^0$ does not affect the production rate, except insofar as it must be lower than the masses of all other superpartners, but it does affect the character of the events. When the mass splitting between the gluino or squark and $\tilde{\chi}_1^0$ is small, only a small portion of the event’s energy ends up in the visible decay products, and most is lost to the rest energy of $\tilde{\chi}_1^0$. These events have low H_T and \cancel{E}_T , low N_{jet} , and when applicable, low $N_{\text{b-tag}}$, and more closely resemble background. As the mass splitting increases, more energy shifts to the visible portion of the event, making events less background-like and increasing sensitivity. The analysis considers a grid of potential mass points for each model, and this sensitivity pattern is evident in the curves shown in Section 3.1.8.

The classic search also considers non-supersymmetric models.

The first is referred to as the mono- ϕ model. In this model, a colored scalar boson much like a squark is produced singly, rather than pair-produced as in squark models in order to conserve R-parity, and decays to a quark and invisible fermion, as shown in Figure 3.2. This model has an especially low number of jets and low M_{T2} , and so would appear in more background-like bins than most supersymmetric models. In fact, it was originally proposed [1] to explain a potential excess in bins of this sort in the previous edition of the classic search, published based on 2016 data [27], and so there is some interest in whether such an excess persists in the larger dataset. While the analysis is not optimized for models of this sort, indeed M_{T2} is designed to target pair-production, it still has some residual sensitivity to any model characterized by jets and missing energy in the final state, including mono- ϕ .

The second non-supersymmetric model, and final model considered explicitly by the classic search, is leptoquark extensions of the Standard Model. As discussed in Section 1.4, while squarks can decay to a quark and $\tilde{\chi}_1^0$, leptoquarks decay to a quark and a neutrino, which is experimentally effectively indistinguishable from a low mass $\tilde{\chi}_1^0$. This last fact was first established in a reinterpretation of the 2016 edition of the classic search [29]. Thus, it is a relatively simple exercise to reinterpret squark squark results in the low mass $\tilde{\chi}_1^0$ limit to leptoquark results. A few leptoquark production diagrams are shown in Figure 3.3.




Figure 3.2: Diagram for the mono- ϕ model, in which a colored scalar ϕ is resonantly produced, and decays to an invisible massive Dirac fermion ψ and an SM quark. Note that ϕ is not pair-produced, in contrast to otherwise-similar squarks. Taken from [11].

By necessity and unlike backgrounds, the expected event yields for signal models are obtained directly from simulation. For this reason, while most uncertainties are shared with background, signal estimates are exposed to a few extra uncertainties, and some of the shared uncertainties increase in magnitude.

First, while background is normalized to control region counts and so automatically scaled to the correct integrated luminosity, signal counts must be scaled by an independent measurement




Figure 3.3: Diagrams of leptoquark pair production. Each leptoquark decays to a neutrino and a quark. Taken from [11].

Table 3.1: Systematic uncertainties in the signal yields for the simplified models of BSM physics. The large statistical uncertainties in the simulated signal sample come from a small number of bins with low acceptance, which are typically not among the most sensitive bins contributing to a given model benchmark point. Taken from [11].

| Source | Range [%] |
|---|-----------|
| Integrated luminosity | 2.3–2.5 |
| Limited size of MC samples | 1–100 |
| b -tagging efficiency, heavy flavors | 0–40 |
| b -tagging efficiency, light flavors | 0–20 |
| Lepton efficiency | 0–20 |
| Jet energy scale | 5 |
| Fast simulation \cancel{E}_T modeling | 0–5 |
| ISR modeling | 0–30 |
| μ_R and μ_F | 5 |

of the integrated luminosity. This measurement is only precise to within a few per cent, leading to a few per cent uncertainty on the expected event count.

Second, additional jets obtained from ISR must be modeled in simulation for signal, while for background, the control regions provide a precise measurement of the relevant ISR jet production. In bins with many jets, where ISR jet production rates are less well-known and simultaneously more important, this uncertainty is as large as 30%.

Finally, signal models must be simulated in the fast-simulation framework discussed in Section 2.7 to make the enormous array of signal scenarios considered computationally feasible.

A systematic is assessed to cover potential errors made by this approximation.

The remaining uncertainties are described in the next section, in the context of background estimates.

3.1.3 Backgrounds

All of the targeted signals are characterized by all-hadronic events with large missing energy, but observing such an event does not constitute discovery due to the existence of backgrounds that can produce the same basic signature. These backgrounds can be broadly divided into the detector mismeasurement background, in which apparent missing energy is generated not by genuine undetected particles but by an error in reconstruction, and the neutrino background, in which the missing energy is produced by genuine neutrinos as predicted in the Standard Model. The neutrino background can be subdivided into neutrinos originating from $W^\pm \rightarrow \ell^\pm v$, in which the presence of a charged lepton allows the neutrino to be rejected with good efficiency, and those originating from $Z \rightarrow vv$, in which the final state is entirely invisible and cannot be efficiently vetoed.

Mismeasurement

The most problematic background is that caused by detector mismeasurement. Nearly every mismeasured event is a QCD multijet event, because nearly every event at a proton-proton collider is a QCD multijet event and the probability of mismeasurement is roughly flat across events. Accordingly, the mismeasurement background is also referred to as the QCD multijet background. While the detector makes mistakes only very rarely, QCD events are so relatively common that this background is still dominant in the raw dataset, before any cleaning selections.

Estimating the QCD background is challenging because it requires highly detailed knowledge of the detector’s idiosyncrasies. Rather than risk falsely discovering a signal or failing to identify one that is present due to misprediction of this background, the analysis adopts selections

designed to suppress it, until the background is sufficiently minor that large relative error in its prediction is acceptable.

The first and most powerful of these selections uses the M_{T2} variable described in Section 2.2.6, namely $M_{T2} > 200$ GeV, where the value is chosen to achieve the desired suppression of the mismeasurement background. Although the M_{T2} selection is somewhat expensive in the sense that it eliminates a significant fraction of signal, especially for signals with a small mass splitting and signals like mono- ϕ that are not pair-produced, it is crucial for suppressing the mismeasurement background. At $H_T > 1500$ GeV, the mismeasurement background extends unacceptably beyond $M_{T2} \sim 200$ GeV, and the M_{T2} selection is tightened to $M_{T2} > 400$ GeV.

The second selection uses the observable

$$\Delta\phi_{min}^{1,2,3,4} = \text{Min} \left(\left| \phi_{\cancel{E}_T} - \phi_i \right| \right)$$

where ϕ_i indicates the ϕ coordinate of the i th p_T jet and $\phi_{\cancel{E}_T}$ is the ϕ coordinate of the missing energy vector. Stated simply, $\Delta\phi_{min}^{1,2,3,4}$ is the smallest angular separation in the transverse plane of the missing energy vector and any of the four highest p_T jets. Close overlap between a jet and the missing energy vector indicates a high probability that the jet was badly mismeasured, and is itself the source of the \cancel{E}_T in the event. The selection applied is $\Delta\phi_{min}^{1,2,3,4} > 0.3$, where the value is chosen to achieve strong background rejection without too great a loss of signal efficiency. Only the four highest energy jets are used because the probability of *some* jet overlapping the missing energy vector approaches unity as the number of jets increases, so the selection would nearly always veto high N_{jet} events, and the high N_{jet} bins are sufficiently signal-rich that aggressive background rejection is not as necessary. The effect of this selection is depicted in Figure 3.5, after an M_{T2} selection of only 100 GeV.

The last selection rejects events in which a suspiciously large fraction of the missing energy comes from very soft objects. The minimum p_T for selected jets is 30 GeV. The missing

energy vector from only jets is denoted \vec{H}_T . The missing energy vector used in the analysis, \vec{E}_T' , uses all PF candidates, including those outside jets or in jets with $p_T < 30$ GeV. If these two quantities are very different, it means that a large portion of the \vec{E}_T' in the event was generated by these low p_T objects, a sign that something may have gone wrong in reconstruction. Specifically, the selection is $|\vec{H}_T - \vec{E}_T'| / |\vec{E}_T'| < 0.5$.




Figure 3.4: The distributions in M_{T2} of the QCD background (filled yellow) and the neutrino backgrounds (filled blue and green) are stacked and overlaid with an example signal point (gluino pair production and decay to bottom quarks, in red). Even with other mismeasurement-suppression selections applied, the mismeasurement background still dominates without $M_{T2} > 200$ GeV. Taken from [19].

The residual mismeasurement background is estimated using a procedure called Rebalance and Smear that was newly implemented for this edition of the classic analysis, and described in Section 3.1.5.

Lost Lepton

The lepton veto rejects most events containing neutrinos originating from the decay of a W boson, $W^\pm \rightarrow \ell^\pm \nu$. However, the lepton is not always successfully reconstructed, usually because the lepton is a τ that decays hadronically and is mistaken for a meson, and this residual




Figure 3.5: The distributions in $\Delta\phi_{min}^{1,2,3,4}$ of the QCD background (filled yellow) and the neutrino backgrounds (filled blue and green) are stacked and overlaid with an example signal point scaled up by a factor of 5 (bottom squark pair production, in red), with the M_{T2} selection relaxed to 100 GeV. Note that most QCD lies below the cut value of 0.3. Taken from [19] supplementary materials.

so-called lost lepton background must be estimated.

A first attempt might be to simulate events containing W bosons, scale the simulation to the desired luminosity, and count the events in which the lepton is not reconstructed. This procedure would have reasonable accuracy, but producing accurate simulations is not trivial, and the result would be subject to an array of systematic errors. Schematically, if N_{LL}^{Data} is the actual number of lost lepton events in data and N_{LL}^{MC} is the number predicted by Monte Carlo simulation, the simulation will mispredict by some factor ε such that $N_{LL}^{\text{Data}} = \varepsilon N_{LL}^{\text{MC}}$.

It is possible to do better than ε with data driven techniques. In addition to the lost lepton events, one can also ask the simulation for its prediction of the number of W events in which the lepton is not lost, single lepton events, N_{SL}^{MC} , where we restrict the lepton to electrons and muons since τ reconstruction is much more difficult. Every part of this simulation is exactly identical to the lost lepton simulation, subject to almost the same errors, with the major exception being the predicted lepton reconstruction efficiency. Call this new error factor δ so that $N_{SL}^{\text{Data}} = \delta N_{SL}^{\text{MC}}$.

Then $\varepsilon/\delta = \gamma$ is the portion of the misprediction due to the simulation's imperfect knowledge of the lepton reconstruction efficiency and a few other more minor uncorrelated effects, a small portion of the total. The ratio $R_{\text{MC}}^{0\ell/1\ell} = N_{LL}^{\text{MC}}/N_{SL}^{\text{MC}}$ is subject only to this relatively small error γ , since the fully correlated errors cancel. The prediction of N_{LL}^{Data} follows directly,

$$N_{LL}^{\text{Data};\text{Est}} = R_{\text{MC}}^{0\ell/1\ell} N_{SL}^{\text{Data}}. \quad (3.1)$$

The input N_{SL}^{Data} is measured in a control region populated by single lepton events observed in data, in a kinematic region identical to the corresponding lost lepton signal region. The remaining systematic uncertainty is only about 15% in most signal regions.

Table 3.2: Summary of systematic uncertainties in the lost-lepton background prediction, together with their typical size ranges across the search bins. (†) In every topological region, bins along the M_{T2} axis with insufficient statistics to allow a fully data-driven estimate are assigned an M_{T2} shape from simulation, normalized based on the integral of the affected bins in the data control region. Taken from [11].

| Source | Range [%] |
|--|-----------|
| Limited size of data control samples | 5–100 |
| Limited size of MC samples | 0–50 |
| e/μ efficiency | 0–10 |
| τ efficiency | 0–3 |
| b -tagging efficiency | 0–3 |
| Jet energy scale | 0–5 |
| $M_T(\text{lepton}, \vec{E}_T)$ selection efficiency | 0–3 |
| M_{T2} shape uncertainty [†] | 0–40 |
| Renormalization and factorization scale variation | 0–5 |
| $t\bar{t}b\bar{b}/t\bar{t}jj$ weight | 0–25 |

Other uncertainties affecting the lost lepton background prediction are summarized in Table 3.2, together with their typical size ranges across the search bins. The first two uncertainties due to limited data control region and Monte Carlo simulation sample sizes are purely statistical. The remainder are systematic.

The efficiency to reconstruct leptons is, as mentioned, the primary uncertainty of the

simulation with respect to data, that does not drop out after taking a ratio of lost- and single-lepton events.

The impact of the τ reconstruction efficiency is smaller, as it is not used in the single lepton control region, however uncertainty on the rate at which the τ veto fails to reject events contributes to uncertainty in the expected yield.

The expected transfer factor from the single lepton to lost lepton regions varies slightly with respect to the binning variables, H_T , N_{jet} , $N_{b-\text{tag}}$, and M_{T2} , and so is recalculated in simulation for each bin. Since the transfer factor varies, they may be incorrectly estimated if simulation does not accurately reproduce the b -tagging efficiency and jet energy assignments in real data.

To restrict the single lepton control region to leptons consistent with originating from a leptonic W decay, the lepton and \cancel{E}_T system is required to satisfy $M_T(\text{lepton}, \cancel{E}_T) < 100 \text{ GeV}$, since an on-shell W can never decay to a lepton-neutrino system more massive than this, and $M_T \leq M$.

The renormalization and factorization scales are generic issues with theoretical calculations, discussed in Section 2.6.

Finally, the $t\bar{t}b\bar{b}/t\bar{t}jj$ weight is a special systematic assessed to cover an ad hoc procedure that corrects a particularly severe data-simulation discrepancy affecting the relative rates of $b\bar{b}$ ISR and light flavor ISR in simulated $t\bar{t}$ events. This simulation error is not significant for the majority of CMS analyses, however it is important for analyses like this search that include bins with large $N_{b-\text{tag}}$, for which $t\bar{t}b\bar{b}$ is the only meaningful background.

In closing, it should be noted that while the lost lepton background tends to be subdominant relative to the Invisible Z background discussed in the next section, it is the largest background in certain high N_{jet} , high $N_{b-\text{tag}}$, high H_T bins since Z events do not populate these bins efficiently, while $t\bar{t}$ pair-production events do, and all $t\bar{t}$ events contain two W bosons that may decay leptonically.

Invisible Z

The $Z \rightarrow vv$ background is predicted in a similar fashion to the lost lepton background, using a control region populated with $Z \rightarrow \ell^+\ell^-$ events. Again, the leptons are restricted to pairs of electrons and pairs of muons, since τ reconstruction is much more difficult. Figure 3.6 (right) shows the similarity in the M_{T2} distributions of simulated $Z \rightarrow vv$ events and observed $Z \rightarrow \ell^+\ell^-$ events in which one pretends that the leptons are invisible, demonstrating both that these events are kinematically very similar as expected, and that the simulation is accurate. The ratio $R_{MC}^{vv/\ell^+\ell^-}$ can be extracted from Monte Carlo simulation, and is dominated by only the lepton reconstruction efficiency uncertainty. In the Standard Model, this ratio is almost exactly 3, but it is significantly larger experimentally because it is possible for one of the leptons in $Z \rightarrow \ell^+\ell^-$ not to be well-reconstructed, causing the affected event not to be counted. The value $N_{\ell^+\ell^-}^{Data}$, unlike N_{SL}^{Data} , is not trivial to extract, since a non-negligible fraction of double lepton events come from sources other than a single Z boson, almost always one each from a pair of W bosons. $N_{\ell^+\ell^-}^{MC}$ and N_{vv}^{Data} are exclusively $Z \rightarrow \ell^+\ell^-$ and $Z \rightarrow vv$ events, respectively, and for maximal cancellation of systematic uncertainties, it is desirable that $N_{\ell^+\ell^-}^{Data}$ be purified to the greatest extent possible.

Fortunately, there is an experimental handle on the contamination. When a Z boson decays to a lepton pair, the flavor is always identical, either two electrons or two muons. When a pair of W bosons each decay to a lepton pair, their choices are uncorrelated. Half the time, the flavors will be identical as for a Z event, and half the time, one W will decay to a muon and the other to an electron. The first case is the undesired impurity in $N_{\ell^+\ell^-}^{Data}$, and events of the second type are used to populated a different-flavor control region used to predict the impurity, N_{DF}^{Data} . It is nearly sufficient simply to subtract N_{DF}^{Data} from $N_{\ell^+\ell^-}^{Data}$ since the different flavor and same flavor W events occur at the same rate. However, it is possible that the detector is slightly more or less efficient at reconstructing events where both leptons are the same flavor than events in which they are different flavors, so the different flavor count must be scaled slightly to compensate, by a factor $R^{SF/DF}$. $R^{SF/DF}$ is measured in data in $\ell^+\ell^-$ events that are kinematically inconsistent




Figure 3.6: (Left) The ratio of the number of same flavor to different flavor lepton pairs in data, $R^{SF/OF}$, which is a component of the $Z \rightarrow vv$ background estimate from $Z \rightarrow \ell^+\ell^-$ events. (Right) A comparison of the M_{T2} shape between data $Z \rightarrow \ell^+\ell^-$ events and simulated $Z \rightarrow vv$ events. The two processes should be kinematically identical, and the similarity of the distributions indicates that the Monte Carlo simulation models the processes well. Taken from [11].

with originating from a Z decay, chiefly due to a requirement that the invariant mass of the lepton pair be at least 20 GeV away from the Z mass. One finds that $R^{SF/OF} \approx 1.06 \pm 0.15$, as shown in Figure 3.6 (left). The final prediction is

$$N_{vv}^{Data;Est} = R_{MC}^{vv/\ell^+\ell^-} (N_{Data}^{\ell^+\ell^-} - R^{SF/OF} N_{DF}^{Data}) \quad (3.2)$$

Being irreducible, the $Z \rightarrow vv$ plus jets background is dominant in the vast majority of bins.

The uncertainties in the $Z \rightarrow vv$ background prediction are summarized in Table 3.3 together with their typical size ranges across the search bins. Only the uncertainty in $R^{SF/DF}$ is unique to $Z \rightarrow vv$, while the others are shared with the lost-lepton estimate.

Table 3.3: Summary of systematic uncertainties in the $Z \rightarrow vv$ background prediction, together with their typical size ranges across the search bins. (\dagger) In every topological region, bins along the M_{T2} axis with insufficient statistics to allow a fully data-driven estimate are assigned an M_{T2} shape from simulation, normalized based on the integral of the affected bins in the data control region. Take from [11].

| Source | Range [%] |
|---------------------------------------|-----------|
| Limited size of data control samples | 5–100 |
| Limited size of MC samples | 0–50 |
| Lepton efficiency | 0–5 |
| Jet energy scale | 0–5 |
| Uncertainty in $R^{\text{SF/DF}}$ | 0–5 |
| M_{T2} shape uncertainty † | 0–40 |

| Observable | Selection | Notes |
|---------------------------------|---------------------|---|
| M_{T2} | $> 200 \text{ GeV}$ | Only for multijet events. Increased to $M_{T2} > 400 \text{ GeV}$ for $H_T > 1500 \text{ GeV}$ to maintain QCD suppression. |
| p_T^{jet1} | $> 250 \text{ GeV}$ | Only for monojet events. |
| H_T | $> 250 \text{ GeV}$ | Motivated by available triggers. Background events at lower H_T are too common for these events to be always recorded. |
| E_T | $> 250 \text{ GeV}$ | Relaxed to $E_T > 30 \text{ GeV}$ for $H_T > 1200 \text{ GeV}$. Motivated by available triggers. |
| $\Delta\phi_{min}^{1,2,3,4}$ | > 0.3 | Auxiliary mismeasurement suppression. |
| $ \vec{H}_T - \vec{E}_T / E_T$ | < 0.5 | Auxiliary mismeasurement suppression. |
| N_{lep} | $= 0$ | The lepton veto; rejects the majority of $W^\pm \rightarrow \ell^\pm v$ background. |

Table 3.4: A summary of the baseline event selection for the classic M_{T2} search. Events are required to have large H_T , no leptons, and significant missing energy unlikely to be the product of detector mismeasurement or a single undetected particle.

3.1.4 Baseline Selection

The properties of these signals and backgrounds motivate the baseline event selection summarized in Table 3.4. The M_{T2} selection primarily suppresses the mismeasurement background, by many orders of magnitude. The $\Delta\phi_{min}^{1,2,3,4}$ and H_T selections also help to suppress this background further, until it is smaller than the genuine \cancel{E}_T backgrounds. The H_T and \cancel{E}_T selections are chosen primarily so that all of the selected events pass the online trigger. Backgrounds are so common at $H_T < 250$ GeV, and $\cancel{E}_T < 250$ GeV for $H_T < 1200$ GeV, that the experiment is unable to record all of the events observed, making this part of the parameter space a poor region to search for a rare signal in any case. The lepton veto rejects most $W^\pm \rightarrow \ell^\pm v$ events, so that only events in which the lepton is not reconstructed remain in the signal region, as described in Section 3.1.3, and serves to narrow the analysis’ focus to the all-hadronic final state as part of CMS’s larger research program.

In addition to this baseline selection, the analysis bins in M_{T2} , H_T , N_{jet} , and N_{b-tag} to enhance signal sensitivity as described in Section 3.1.5.

3.1.5 Upgrades in 2019

The classic search has been performed before at 13 TeV, once in 2015 [19] using a 2.3 fb^{-1} dataset, and again in 2016 [27] using 35.9 fb^{-1} . The update in 2019 uses the full dataset from 2016, 2017, and 2018, totaling 137 fb^{-1} . As the analysis is dominated by statistical uncertainties in its most sensitive bins, the increased statistical power is the primary improvement in the update.

The update includes two other major upgrades.

The first improves the estimate of the mismeasurement background using a technique called Rebalance and Smear.

The second leverages the increased statistics to expand the signal region binning, better targeting signal models with more extreme jet and b-tagged jet multiplicities, and M_{T2} .

Rebalance and Smear




Figure 3.7: The fit of $r_\phi = N_{\Delta\phi_{min}^{1,2,3,4} > 0.3} / N_{\Delta\phi_{min}^{1,2,3,4} < 0.3}$ as a function of M_{T2} obtained in the 2016 classic search, for the $1000 < H_T < 1500$ GeV H_T band. The fit is performed in events in the M_{T2} band $70 < M_{T2} < 100$ GeV and extrapolated to the $M_{T2} > 200$ GeV signal region. Black points represent raw data, while white points represent data after the non-multijet contribution is subtracted. Taken from [27].

In older versions of the classic search [19, 27], the mismeasurement background estimate used the $\Delta\phi_{min}^{1,2,3,4}$ observable and extrapolated across M_{T2} . Events at low M_{T2} and at low $\Delta\phi_{min}^{1,2,3,4}$ are both dominated by QCD mismeasurement. The suppression effect of M_{T2} is so strong that even events at low M_{T2} and *high* $\Delta\phi_{min}^{1,2,3,4}$ are QCD dominated. As essentially every low M_{T2} event is a QCD event, low M_{T2} events can be used to measure the ratio r_ϕ of QCD mismeasurement

events at high and low $\Delta\phi_{min}^{1,2,3,4}$. High M_{T2} events with low $\Delta\phi_{min}^{1,2,3,4}$ can then serve as a control region for estimating the QCD mismeasurement background, $N_{\Delta\phi_{min}^{1,2,3,4} > 0.3} = r_\phi N_{\Delta\phi_{min}^{1,2,3,4} < 0.3}$. Unfortunately, r_ϕ decreases with increasing M_{T2} , so that instead the dependence must be fit to a power law at low M_{T2} and extrapolated to high M_{T2} , as shown in Figure 3.7. This procedure has obvious potential for statistical errors in the fit, systematic error in extrapolating the fit to high M_{T2} , and potential non-multijet contamination of the low $\Delta\phi_{min}^{1,2,3,4}$ control region at high M_{T2} , producing total relative error at least 40% and as large as 180%. Although the impact of these errors is controlled by suppressing the mismeasurement background as described in Section 3.1.3, it is desirable to replace this procedure with a more robust one.

Rebalance and Smear (R&S) achieves the desired improvement. R&S begins with a sample of multijet events from data with H_T on the order of hundreds of GeV and at least two jets with $p_T > 10$ GeV. This is as nearly unbiased a sample of QCD events as is allowed by available triggers. These events will generically have some small but nonzero \cancel{E}_T due to imperfect measurement of the jet energies, dictated by the detector's resolution, jets from pileup, very low energy objects in the event, and potentially a very small contribution from genuine neutrinos produced in the hadron decay chains inside the jets. The p_T values assigned to the jets are then adjusted, finding the most likely assignment of p_T values subject to the hypothesis that the true \cancel{E}_T is very nearly zero, and that jet mismeasurements of a given size occur with probability given by jet response templates. As shown in Figure 3.8, large mismeasurements are rare. Thus, the Rebalancing step tries to get the p_T as close to zero as feasible without making more improbable adjustments to the jets than necessary. The output of the Rebalancing step is a large sample of real QCD events with maximally accurate jet p_T assignments.

Then, each of these events goes through a Smearing step. This step randomly assigns a new p_T to each jet in the event according to the same jet response templates. The vast majority of the time, the new event looks as unremarkable as the original event before Rebalancing. Rarely, the output event passes the baseline selection and represents a potential mismeasurement event




Figure 3.8: $p_{T,ptcl}$ is the “particle level” p_T , the total energy that the jet’s particle components actually had, while p_T is the energy measured by the detector. The horizontal axis is the ratio of these two quantities, and the vertical axis the probability that a jet’s measured energy will differ from its true energy by that ratio. The response curve shown here applies to jets with particle level p_T in the band $97 < p_T < 114 \text{ GeV}$, and measured in the CMS barrel. Large mismeasurements of jet energy are much less probable than small mismeasurements. The core is well-described by a Gaussian, but the tails are highly non-Gaussian and best described by a Crystal-Ball function. Taken from [20].

that might lie in the data signal region. The Smearing can be repeated as many times for each event as is desired, subject to computing limitations. The number of Smeared events falling into each signal region, from this sample of known equivalent integrated luminosity, can then be converted into the expected number of events that would fall into the signal region in data, due to jet mismeasurement.




Figure 3.9: The Rebalance and Smear jet mismeasurement background estimate is validated in the $\Delta\phi_{min}^{1,2,3,4} < 0.3$ control region. The multijet mismeasurement event count predicted by Rebalance and Smear is shown in yellow, and contributes most of the event counts in this control region. The combined background estimate (filled histogram) is consistent with the data observation (black data points). Taken from [11].

Of course, many of the output events will have $\Delta\phi_{min}^{1,2,3,4} < 0.3$, falling outside of the signal region into the QCD-dominated low $\Delta\phi_{min}^{1,2,3,4}$ control region. This allows for a validation in data of the R&S procedure, shown in Figure 3.9. The total background estimate is consistent with data in the $\Delta\phi_{min}^{1,2,3,4} < 0.3$ control region across all of the analysis bins, with the R&S estimated counts contributing most of the predicted events.

R&S achieves a significant improvement over the old r_ϕ based system, effectively eliminating statistical error, and improving the worst case relative error on the mismeasurement estimate from 180% to less than 50%. Systematic uncertainties are summarized in Table 3.5 together with their typical size ranges across the search bins. The sources of these uncertainties are described either in the text above, or are shared with the uncertainties affecting the genuine neutrino backgrounds.

Table 3.5: Summary of systematic uncertainties in the multijet background prediction, together with their typical size ranges across the search bins. (\dagger) The parameter σ_T^{soft} is the width of the assumed underlying irreducible \cancel{E}_T distribution for a multijet event, used as an input to the Rebalancing step, and so-called because such \cancel{E}_T is dominated by low energy objects. The final output is largely independent of the precise choice for this parameter and the shape of the underlying distribution, but a systematic is assessed to cover any observed variations. Taken from [11].

| Source | Range [%] |
|---|-----------|
| Jet energy resolution | 10–20 |
| Tails of jet response in templates | 17–25 |
| σ_T^{soft} modeling \dagger | 1–25 |
| N_{jet} modeling | 1–19 |
| $N_{\text{b-tag}}$ modeling | 1–16 |

Expanded Binning

Although the baseline selection defines the class of events in which a signal of interest may be found, most signals will produce a significant number of events in only a subset of the phase space. For instance, a signal producing 4 top quarks in the final state (see Figure 3.1 second row, right) will almost always produce events with $N_{\text{jet}} \geq 7$, $N_{\text{b-tag}} > 0$, and $H_T \sim 1 \text{ TeV}$. If the entire set of selected events is considered together, the entire background is combined, potentially hiding a signal. If instead the phase space is divided into many separate regions, the background in each region is smaller, so that the background count is as small as possible in the small subset of regions that any given signal actually populates. This sensitivity enhancement motivates very fine binning of the signal region. The classic search uses M_{T2} , H_T , N_{jet} , and $N_{\text{b-tag}}$ as binning variables.

In the 2016 version of the classic search [27], the most extreme N_{jet} bin was $N_{\text{jet}} \geq 7$, and the most extreme $N_{\text{b-tag}}$ bin was $N_{\text{b-tag}} \geq 3$. This limitation was imposed by limited statistics. The background estimate for each bin is performed separately, and the observed counts are subject to Poisson statistical fluctuations. Binning too finely causes any potential sensitivity gains from better isolating signal to be lost to greater uncertainty in the expected background. The analysis binning was updated for the latest edition of the classic search [11], anticipating the improved

statistical power. The full set of new bins, along with the predicted background counts and observed event counts in data, is available in Appendix A.

The new binning extends the old binning in three ways. First, new $N_{\text{jet}} \geq 10$ bins were added to the $H_T > 1200$ GeV regions. This allows for enhanced sensitivity to signals with extremely high jet multiplicity, such as the 4 top signal previously mentioned. Similarly, new $N_{b-\text{tag}} \geq 4$ bins were added to these same regions, targeting the same signal. Sensitivity to some mass points of this signal model roughly doubled due to the addition of these bins, which have negligible background but appreciable signal counts. Finally, M_{T2} binning was generally made narrower and the last bin moved to larger M_{T2} values, for all signal regions, until the expected background in the last bin was on the order of 1 event. In all, there are 282 classic search bins, enhancing sensitivity to a broad array of potential signal models.

3.1.6 Signal Contamination

The data driven estimate procedures for the neutrino backgrounds used control regions defined using leptons. Signals capable of producing leptons can contaminate these control regions, increasing the observed control region counts above the actual Standard Model production rate. This leads directly to an overprediction of background. For instance, consider stop pair production followed by decay to tops, as shown in Figure 3.1 (row 3, right). Both tops will decay to a W and a bottom quark. If one of the W bosons decays leptonically and the lepton is reconstructed, the event will likely enter the single lepton control region. The lost lepton background, described in Section 3.1.3, will be overpredicted,

$$N_{LL}^{\text{Data;Est}} = R_{MC}^{0\ell/1\ell} N_{SL}^{\text{Data}} = R_{MC}^{0\ell/1\ell} \left(N_{SL}^{\text{Data;SM}} + N_{SL}^{\text{Data;BSM}} \right). \quad (3.3)$$

The background overprediction is $\Delta N = R_{MC}^{0\ell/1\ell} N_{SL}^{\text{Data;BSM}}$. For analysis purposes, the overprediction is modeled in simulation and treated as a reduction of the expected signal counts in every

affected bin.

$$N_{SR}^{\text{BSM;Adjusted}} = N_{SR}^{\text{BSM;Raw}} - \Delta N \quad (3.4)$$

This adjustment has the nice property that all terms are linear in the signal strength, so that it does not need to be recalculated for every signal strength considered when performing statistical analysis of the results. The same fraction of the signal is lost at every signal strength.

As a result of this loss of sensitivity due to signal contamination, the classic analysis is less effective, all things equal, when used to search for signals that sometimes produce leptons than the naive expectation based on the leptonic versus hadronic branching ratios. This is an inevitable consequence of performing an all-hadronic search.

3.1.7 Results

The full set of results for every classic search signal region, including every background prediction and the observed count, are available in Appendix A. The full set of results integrating over the M_{T2} binning are display in Figure 3.10 (upper), and the results including the M_{T2} binning for the $575 < H_T < 1200$ GeV bins, the largest set, are shown in Figure 3.10 (lower). The observed counts are consistent with the background-only hypothesis, and the results are used to set exclusion limits at 95% confidence level on the signals discussed in Section 3.1.2.

3.1.8 Limits

The CL_S statistical analysis procedure is decribed in [25]. It begins with maximum likelihood fits to the background-only and signal-plus-background hypotheses, for each signal model, considering each mass point separately. The likelihoods are products of Poisson probabilities for each signal region bin, with log-normal constraint terms for each systematic affecting the predicted counts. Correlation between uncertainties affecting different bins are fully accounted for. As stated in the previous section, the background-only hypothesis is consistent with the




Figure 3.10: Comparison of predicted background and observed data events in the classic search (upper) integrated over M_{T2} and (lower) for each of the medium H_T search regions. Taken from [11].

observation. So, the parameter of interest for each signal model is the maximum signal cross section that can be excluded with 95% confidence level. That is, the maximum possible production

rate that the signal model could have, without requiring a combination of background and signal fluctuations more improbable than 1 in 20 to be consistent with the data. If the production cross section excluded at 95% CL is less than the theoretical cross section for a given signal model, that signal model is said to be excluded at 95% CL.

The simplified supersymmetric extensions to the Standard Model shown in Figure 3.1 have only two free parameters, the masses of the pair-produced superpartner, and the mass of the lightest supersymmetric particle, the dark matter candidate $\tilde{\chi}_1^0$. Plotting the gluino or squark mass on the horizontal axis and the mass of $\tilde{\chi}_1^0$ on the vertical axis, then marking the mass points excluded at exactly 95% CL, produces the exclusion curves shown in Figures 3.11–3.14. Points to the lower left of these curves are excluded, while points above and to the right are not, as they require fluctuations no more improbable than 1 in 20 to be consistent with present data.

The CMS dataset was collected only once, and this dataset is subject to fluctuations. The multiple exclusion curves shown on each plot, some in red and some in black, provide an indication of how unusual the limits generated by this dataset were. Suppose that the background’s expected event count in a given region, summing across this entire dataset, is B . It is clearly possible that the actual number of background events that occur in the dataset in this bin could be around $B + \sqrt{B}$, or $B - \sqrt{B}$. In the first case, the analysis will draw unexpectedly strong limits on signals that populate this bin. In the second case, the analysis will draw unexpectedly weak limits on signals.

The curves shown in red are the median, 1 standard deviation, and 2 standard deviation expected exclusion curves, in the background-only hypothesis. Together, these indicate the typical range of exclusion curves across many imaginary CMS datasets in which there is no signal to find. The curves in black are those that were observed in the CMS dataset as actually recorded. When the black observed curves extend out beyond the red expected curves, the analysis likely benefitted from a downward fluctuation of background. In cases where the observed curves swing inward compared to the expected curves, the analysis may have experienced an analogous unlucky

upward fluctuation of background that mimicked a small signal, or alternatively there may be a genuine small signal lurking in the data that is resisting exclusion!

The general shape of the curves is set by a combination of the falling cross section with increasing gluino or squark mass shown in Figure 1.1, and a loss of signal efficiency and signal versus background discriminatory power when the mass splitting between the gluino or squark and $\tilde{\chi}_1^0$ is small. On the lower right hand side of the plots lie signals that produce spectacular, energetic events, but at a very low rate due to the large mass of the gluino and squark, and accordingly small pair-production cross section. Moving up along the $\tilde{\chi}_1^0$ mass axis towards the upper right, the events remain spectacular, so the exclusion curves remain roughly vertical, limited only by the production cross section. Eventually, the mass splitting becomes small enough that the loss of signal efficiency and signal versus background discriminatory power becomes significant, and the exclusion curve turns to the left, towards lower mass squarks or gluinos and higher production rates. Decreasing the gluino or squark mass at fixed $\tilde{\chi}_1^0$ mass reduces the mass splitting, so the curve generally must also drop down somewhat along the $\tilde{\chi}_1^0$ mass axis to maintain a reasonable splitting. Eventually, the curve intercepts the $M_{\tilde{\chi}_1^0} = M_{\tilde{g}}$ or $M_{\tilde{\chi}_1^0} = M_{\tilde{q}}$ line. At this mass and below, all points are excluded even in the limit of zero mass splitting.

Figure 3.11 shows the exclusion curves for gluino pair production and decay to light quarks (upper), light quarks and the Z boson (lower left), and light quarks and the W boson (lower right). Figure 3.12 shows the exclusion curves for gluino pair production and decay to bottom (left) and top (right) quarks.

Figure 3.13 shows the exclusion curves for light-flavor squark (upper left), bottom squark (upper right), and (lower) top squark pair production in which the top squark decays to a top quark. The light-flavor figure contains two curves, one which assumes that there is only a single low mass light flavor squark, and another that assumes that there are eight light flavor squarks of (approximately) degenerate mass, which implies a production cross section eight times larger. The other top squark decay modes, in which the top decays to (upper left) a bottom quark and $\tilde{\chi}^\pm$,




Figure 3.11: Exclusion limits at 95% CL_S for gluino pair production and decay to a pair of light quark jets and (upper) $\tilde{\chi}_1^0$, (lower left) a democratic split between $\tilde{\chi}_1^0, \tilde{\chi}_2^0$, which then decays to a Z boson and $\tilde{\chi}_1^0$, and $\tilde{\chi}^\pm$, which decays to a W boson and $\tilde{\chi}_1^0$, and (lower right) $\tilde{\chi}^\pm$ then W and $\tilde{\chi}_1^0$ with 100% branching fraction. Taken from [11].




Figure 3.12: Exclusion limits at 95% CL_S for gluino pair production and decay to (left) bottom quarks and (right) top quarks. Taken from [11].

which subsequently decays to a W boson and $\tilde{\chi}_1^0$, (lower) a charm and $\tilde{\chi}_1^0$, and (upper right) either $\tilde{\chi}^\pm$ and a bottom quark or $\tilde{\chi}_1^0$ and a top quark, are shown in Figure 3.14. The charm decay channel is only shown for signal models with small mass splittings, as the top squark would strictly prefer to decay to a top quark and $\tilde{\chi}_1^0$ than to a charm quark and $\tilde{\chi}_1^0$ if kinematically allowed.

Figure 3.15 shows limits placed on the mono- ϕ model. Here, the horizontal axis is the mass of the singly-produced scalar ϕ , and the vertical axis the mass of the invisible fermion ψ . A star indicates the mass point proposed by the original authors in [1] as most phenomenologically interesting, which is not yet excluded. It is worth emphasizing that the background model is nevertheless consistent with data; this signal is simply very difficult to exclude with the M_{T2} analysis methodology. To save computing resources, the mono- ϕ model is only simulated in the phenomenologically interesting subset of the mass plane, similarly to the top squark to charm model in Figure 3.14 (lower), hence the large white space beneath the considered range of masses.

Figure 3.16 shows limits for leptoquarks decaying to (upper left) a light flavor quark and neutrino, (upper right) a bottom quark and neutrino, and (lower) a top quark and neutrino. As the mass of the neutrinos, in contrast to the mass of $\tilde{\chi}_1^0$, are known to be approximately zero, the




Figure 3.13: Exclusion limit at 95% CL_S for (upper left) light-flavor squark pair production, (upper right) bottom squark pair production, and (lower) top squark pair production. Taken from [11].




Figure 3.14: Exclusion limits at 95% CLs for top squark pair production and decay to (upper left) a bottom quark and $\tilde{\chi}_1^\pm$, which subsequently decays to a W boson and $\tilde{\chi}_1^0$, (upper right) either a bottom quark and $\tilde{\chi}_1^\pm$ or top quark and $\tilde{\chi}_1^0$, or (lower) a charm quark and $\tilde{\chi}_1^0$. Taken from [11].




Figure 3.15: Exclusion limit at 95% CL_S for the mono- ϕ model. Only the portion of the mass plane of phenomenological interest was simulated. The star indicates the original authors’ proposed best fit mass point, which remains unexcluded. Taken from [11].

leptoquark limits are one dimensional, in the leptoquark masses.

The limits produced by this edition of the classic search improve upon the limits set by the previous edition [27] by hundreds of GeV, and in most cases are the strongest constraints on their respective signal models yet produced by any experiment.

3.1.9 Future of the Classic M_{T2} Search

The current limits produced by the classic M_{T2} search are impressive, and are unlikely to improve much in the near future. The pair production cross section for squarks and gluinos drops rapidly with mass, as shown in Figure 1.1. Generically, a factor of 10 improvement in sensitivity is necessary to push the exclusion limits outward by around 500 GeV along the horizontal axis. At this mature statistical stage, a factor of 10 improvement in sensitivity requires a factor of 100 increase in the integrated luminosity, infeasible in the near future. Improvement along the $\tilde{\chi}_1^0$ axis is similarly difficult due to large backgrounds and low signal efficiency for models with small mass splittings. Therefore, attention in the near future will turn to other new techniques, one of




Figure 3.16: Upper limits at 95% CL_S on the leptoquark production cross sections as a function of leptoquark mass. Unlike the limits on supersymmetric models, in which the mass of $\tilde{\chi}_1^0$ is a free parameter, the limits on leptoquarks are 1-dimensional in the leptoquark mass since the neutrino masses are known to be approximately zero. Taken from [11].

which is discussed in the next section.

3.2 Disappearing Tracks Search

3.2.1 General Description

As searches like that discussed in the previous section approach the practical limits of their sensitivities at the LHC, interest has grown in targeting plausible signals with peculiar features that more standard searches do not exploit. Among these features are those produced by relatively long-lived particles (LLPs) that do not decay promptly at the collision point, but instead travel a macroscopic distance into the detector before decaying. An overview of some possibilities in the context of supersymmetry is provided in [23].

One of these signatures is a disappearing track, produced when a charged particle, such as $\tilde{\chi}^\pm$, travels into the tracker, then decays to an invisible particle, such as $\tilde{\chi}_1^0$, with a mass so nearly equal to that of the decaying charged particle that the visible products of the decay are too low energy to be reconstructed. The supersymmetric parameter space is vast and there is no way to know which if any version of supersymmetry is realized in nature, but models including LLPs that would produce disappearing tracks at CMS have been argued to be especially plausible on theoretical grounds [17, 24, 6]. The M_{T2} analysis is already well-optimized for all-hadronic supersymmetry, so it is natural to search for all-hadronic decays of these models by extending it with a disappearing track search. This extension begins with exactly the same set of selected events as the classic search, and then further selects the subset of events possessing a disappearing track.

3.2.2 Challenges

The disappearing tracks search must confront a pair of significant challenges.

First, the background (see Section 3.2.4) entirely consists of events in which reconstruction failed, analogous to the classic search’s detector mismeasurement background that was discussed in Section 3.1.3, and is difficult to understand and estimate for many of the same reasons.

Simulation cannot be trusted to account for all of the detector details that lead to extremely rare reconstruction errors. While the classic search can mitigate the impact of this issue with selections that make the mismeasurement background subdominant compared to the better understood neutrino backgrounds, this is not possible for the disappearing track search since no such genuine background exists. Moreover, since the background is dependent on details of the detector, it is strongly affected when these details change. Most prominently, the CMS pixel detector was entirely replaced between the 2016 and 2017 data taking periods [26], and this profoundly affected CMS track reconstruction, to the extent that the disappearing track search must treat the 2016 data entirely separately from the 2017 and 2018 data. The detector’s state also evolves more subtly over the data taking period, due in large part to inevitable radiation damage, and the analysis must track, study, and account for the effect of this evolution on the occurrence of disappearing tracks.

Second, disappearing tracks are extremely rare. This is of course one of the motivations for the disappearing tracks extension—they are a powerful tool to discriminate between background and likely signal events—but it also makes them difficult to study.

Nevertheless, the disappearing tracks search produced a data-driven background estimation procedure, described in Section 3.2.5, that was successfully validated in data, and achieved the best sensitivity to disappearing tracks in supersymmetric models to date.

3.2.3 Signal Models

For the class of supersymmetric models discussed in [24] & [6], the relative mass splitting of $\tilde{\chi}^\pm$ and $\tilde{\chi}_1^0$ is proportional to $(M_W/\mu)^4$, where μ is a parameter that roughly represents the energy scale of supersymmetry. Non-observation of superpartners thus far at the LHC suggests that μ is probably large, on the order of TeV, implying $(M_W/\mu)^4 < 10^{-4}$. Such a tiny mass splitting leads to a remarkably long $\tilde{\chi}^\pm$ lifetime, shown in Figure 3.17.

Motivated by these models, the disappearing track search considers signals that include an intermediate $\tilde{\chi}^\pm$ in the superpartner decay chain, similar to the models in Figure 3.1 (upper




Figure 3.17: The rest frame lifetime of $\tilde{\chi}^\pm$ as a function of the $\tilde{\chi}^\pm$ - $\tilde{\chi}_1^0$ mass splitting. The lifetime is macroscopic for mass splittings less than 1 GeV in a broad class of supersymmetric models, a regime that is realized when the superpartner mass scale is much larger than the Standard Model mass scale. This mass splitting is so small that the charged daughter of $\tilde{\chi}^\pm$ decay is too low energy for track reconstruction. Thus, $\tilde{\chi}^\pm$ lives long enough to produce a short track in the CMS tracker, then disappears when it decays to $\tilde{\chi}_1^0$ and a lost charged daughter. For mass splittings less than $M_\pi \approx 140$ MeV, indicated on the plot, the lifetime of $\tilde{\chi}^\pm$ is so long that it typically does not decay inside the tracker. As a result, the sweet spot for the disappearing tracks search is lifetimes on the order of hundreds of MeV. The discontinuity at a mass splitting of 1.4 GeV is not physical. Taken from [6].




Figure 3.18: Diagrams for (left) gluino, (center) light-flavor squark, and (right) top squark pair production, in which the gluinos and squarks can decay via a long-lived $\tilde{\chi}^\pm$. In this analysis, the gluino is taken to decay with branching fraction 1/3 each to the $\tilde{\chi}_1^0$, $\tilde{\chi}^\pm$ plus, and $\tilde{\chi}^\pm$ minus. Squarks decay with branching fraction 1/2 each to the $\tilde{\chi}_1^0$ and the $\tilde{\chi}^\pm$ allowed by charge conservation. The $\tilde{\chi}^\pm$ mass is greater than the $\tilde{\chi}_1^0$ mass by hundreds of MeV, so that the charged product of the $\tilde{\chi}^\pm$ decay is too soft to be detected. Taken from [11].

right, lower left, and lower center) considered by the inclusive search. The $\tilde{\chi}^\pm$ is taken to be long-lived, only hundreds of MeV more massive than $\tilde{\chi}_1^0$. Specifically, the rest frame $\tilde{\chi}^\pm$ lifetime, τ_0 , is varied from $c\tau_0 = 1$ cm to $c\tau_0 = 2000$ cm. This range of lifetimes is selected because the CMS tracker extends from $r \approx 10$ cm to $r \approx 100$ cm, and a significant fraction of $\tilde{\chi}^\pm$ decays must occur inside the tracker for the disappearing track search to have meaningful sensitivity. If the decay occurs outside the tracker, the track does not disappear. The full set of models considered is shown in Figure 3.18.

In the model at left, gluinos are pair-produced and decay with equal probability either to $\tilde{\chi}_1^+, \tilde{\chi}_1^-,$ or $\tilde{\chi}_1^0$, and light-flavor quarks.

In the center second, light flavor squarks are pair-produced and decay with equal probability either to the allowed charge of $\tilde{\chi}^\pm$, or to $\tilde{\chi}_1^0$, and a light-flavor quark.

In the model at right, top squarks are pair-produced and decay with equal probability either to the allowed charge of $\tilde{\chi}^\pm$, or to $\tilde{\chi}_1^0$, and a bottom quark.

The disappearing tracks produced by $\tilde{\chi}^\pm$ decays tend to be very high quality and highly isolated (shown in comparison with background in Figure 3.19), with negligible energy deposits in any of the calorimeter cells along the track's extrapolated path. The quality metrics in this context include the track's impact parameter with respect to the primary vertex, the size of the uncertainty in the track's assigned p_T , and the number of tracker layers in which the track failed to leave a hit when one would be expected. Alongside isolation, shown in Figure ??, these quality metrics allow for signal-like disappearing tracks to be distinguished from disappearing tracks that are very likely to be due to a reconstruction error. This forms the basis of the background estimate described in Section 3.2.5.

3.2.4 Backgrounds

In contrast to the classic search, there are no irreducible disappearing track backgrounds produced by the Standard Model. However, there are a number of detector reconstruction issues




Figure 3.19: Disappearing tracks produced by $\tilde{\chi}^\pm$ decays (black triangles) tend to be much more isolated than disappearing tracks in background events (red squares). All selections listed in Table 3.6 are applied except the isolation selections. The last bin is an overflow bin, containing every track failing the relative isolation selection. The events containing those tracks will either populate control regions, or be rejected entirely, as described in Section 3.2.5. Additional material from [11].

that can lead to apparent disappearing tracks, many of which can be greatly or even entirely rejected by various cleaning selections.

Of the near-exclusively reducible backgrounds, the largest before cleaning is that produced by electrons undergoing unusually strong bremsstrahlung inside the tracker, effectively converting the electron into a photon. While electrons are charged and therefore produce tracks, photons are neutral and do not. So, these tracks disappear, but can normally be rejected due to large photon energy deposits in the electromagnetic calorimeter cells to which the track extrapolates. However, some ECAL cells do not perform well for a variety of reasons and may fail to register the photon energy deposits, and so fail to veto the disappearing track. Fortunately, the majority of

the ECAL performs well enough to make this background significant only for tracks extrapolating to defective cells.

Defective cells are mapped and vetoed using a tag-and-probe strategy, in which a PF electron “tag” identifies potential $Z \rightarrow e^+e^-$ events, and a disappearing track of opposite charge serves as the “probe.” If an event contains exactly one PF electron, the event is searched for a disappearing track of opposite charge, under the hypothesis that the event was in fact $Z \rightarrow e^+e^-$ and the second electron was not successfully reconstructed due to the conversion process described above. If such a track is found and the invariant mass of the electron-disappearing track pair is consistent with the mass of the Z boson, the region of the ECAL to which the disappearing track extrapolated is deemed faulty. $Z \rightarrow e^+e^-$ events are sufficiently common that the map of faulty ECAL cells can be assembled as a function of time. As expected, the performance of the ECAL steadily degrades over time, so that the fraction of the ECAL that must be vetoed increases from 2016 to 2017 to 2018. This veto essentially entirely rejects the electron conversion background, but at a price. It is the single largest cause of signal inefficiency, as 15-20% of signal disappearing tracks point, by chance, into vetoed regions of the ECAL.

Another, much smaller reducible background is produced by the decay of strange baryons, including the Ξ^\pm , Σ^\pm , and Ω^\pm . These baryons have lifetimes in the proper range to produce tracks 10s of centimeters long, before decaying with a small mass splitting to a neutral baryon and a charged daughter that is sometimes low enough p_T to escape reconstruction. While these tracks disappear from the perspective of the tracker and the ECAL, the neutral baryon is still detectable in the hadronic calorimeter via its nuclear interactions, which allows for efficient rejection of this background. Additionally, strange baryons are almost always found inside jets, so the vast majority of their tracks can be vetoed on the basis of isolation. Even so, a region of the HCAL with known performance issues in 2018 showed significantly increased disappearing track counts that required it to be vetoed, likely in part because of this background.

The final two backgrounds are fake tracks and mesons, mostly pions, that undergo nuclear

interactions in the tracker, neither of which can be entirely rejected.

Fake tracks are tracks comprised of hits that are not all associated to the same genuine particle. The probability that track reconstruction produces a fake track is small, and can be suppressed by requiring that tracks pass a sophisticated selection called high purity, as shown in Figure 2.5 for the 2016 tracker [13], which improved after the pixel tracker upgrade between 2016 and 2017 [35]. Furthermore, the probability that more than 3 or 4 unassociated hits will be strung together into a track is negligible, limiting the fake track background to very short tracks. It is also rare for unassociated hits to lie on a tight, relatively straight line; instead, they tend to be somewhat scattered. Therefore, fake tracks also tend to have low assigned p_T , since low energy tracks have greater curvature in the detector's magnetic field. The disappearing tracks analysis considers only tracks with $p_T > 15$ GeV in part to avoid the majority of the fake track background.

While the fake track background is limited to shorter tracks, the meson background is not and so constitutes the dominant longer disappearing track background. This background is generated when a meson undergoes a nuclear interaction inside the tracker, and showers in such a way that no single charged daughter is high enough energy to be reconstructed as a track, and any neutral products are sufficiently low energy or widely scattered that calorimeter deposits are insufficient to reject the track. While it is merely unusual for a meson to undergo a nuclear interaction in the tracker as shown in Figure 2.9, a shower leaving no detectable products is extraordinarily rare, occurring only a handful of times across three years of data taking. Mesons tend not to be isolated, allowing most of this background to be distinguished from signal-like tracks, but with one major exception. The τ lepton, almost always isolated when produced in the hard interaction, undergoes the decay $\tau^\pm \rightarrow h^\pm v$ for some meson h^\pm , almost always a π^\pm , with probability approximately 11.5% [36]. In this decay, the π^\pm effectively inherits the isolation of the τ . If this π^\pm undergoes the rare shower described above, it produces an isolated disappearing track of potentially significant length, scarcely distinguishable from a signal track. For this reason,

observation of even a relatively long, isolated, maximally signal-like disappearing track is not itself sufficient to discover a signal. An estimate of the background’s rate is necessary, to allow a comparison to the observed frequency of disappearing tracks.

3.2.5 The Short Track Selection and Data-Driven Background Estimate

The two surviving disappearing track backgrounds, fake tracks and lost pions, are both produced by extremely rare failure modes of the detector. A data-driven background estimate is mandatory, as replicating such pathological edge cases is beyond the capabilities of any feasible simulation. This requires definition of signal-depleted control regions that can be used to measure the rate at which known physics processes produce disappearing tracks. As discussed with respect to the classic search’s mismeasurement background in Sections 3.1.3 & 3.1.5, events with low M_{T2} are extremely background dominated. Low M_{T2} events compose one such control region. This dependence on M_{T2} to define a signal-depleted control region restricts the disappearing track search to multijet events, as no comparably powerful discriminant is known for monojet events. In high M_{T2} events, properties of the disappearing tracks themselves are used to divide events with signal-like disappearing tracks, called Short Tracks (ST), and background-like disappearing tracks, called Short Track Candidates (STCs), into ST signal regions and STC control regions. The ST and STC selections are summarized in Table 3.6. Signal disappearing tracks pass this selection in simulation with efficiency between 50% and 65%, while only around 1 in 1000 to 1 in 10,000 background events passing the baseline selection of the M_{T2} analysis possess a track passing the ST selection. The STC selection is not maximally background-like; rather, it selects tracks that nearly pass the ST selection, but are not STs. Instead, these tracks pass only a relaxed version of the ST selection. This choice ensures that STCs are as closely linked to STs produced by background as possible, without including a significant amount of true signal tracks. Only a few per cent of signal tracks are selected as STCs, but they are a few times more common in background events than STs.

The selections relaxed for STCs with respect to the ST definition can be grouped into two categories, those concerned with track quality, which are loosened by a factor of 3, and with isolation, which are loosened by a factor of 6. The specific track quality selections affected are the impact parameter, both along the beam axis and in the transverse plane, and the p_T error $\sigma(p_T)$. All isolation selections are loosened.

Table 3.6: Selection requirements for STs and STCs. Some requirements differ for tracks of different lengths; the length categories are described in Section 3.2.5. For the subset of medium (M) length tracks that have just four tracking layers with a measurement, the minimum required number of layers of the pixel tracking detector with a measurement is three (\dagger). The selected tracks are required not to overlap with any identified electrons or muons, of any quality. The selected tracks are as well required to not be identified as PF candidates, and not to overlap with other tracks with $p_T > 15 \text{ GeV}$, even if those tracks are not associated to PF candidates. The table also reports the factor by which a selection is relaxed to define STCs, when applicable. If no factor is reported, the requirement is not relaxed for the selection of short track candidates.

| Observable | Selection | Track length | STC factor |
|---|---------------------------------------|---------------------------|------------|
| p_T [GeV] | > 15 | All | |
| $ \eta $ | < 2.4 and not $1.38 < \eta < 1.6$ | All | |
| $\sigma(p_T) / p_T^2$ [GeV $^{-1}$] | $< 0.2; < 0.02; < 0.005$ | P; M; L | $\times 3$ |
| d_{xy} (from primary vertex) [cm] | < 0.02 (< 0.01) | P (M, L) | $\times 3$ |
| d_z (from primary vertex) [cm] | < 0.05 | All | $\times 3$ |
| Neutral isolation ($\Delta R < 0.05$) [GeV] | < 10 | All | $\times 6$ |
| Neutral isolation / p_T | < 0.1 | All | $\times 6$ |
| Isolation ($\Delta R < 0.3$) [GeV] | < 10 | All | $\times 6$ |
| Isolation / p_T | < 0.2 | All | $\times 6$ |
| Number of pixel layers | ≥ 3 (≥ 2) | P, M ‡ (M, L) | |
| Number of tracker layers | $\geq 3; < 7; \geq 7$ | P; M; L | |
| Number of lost inner hits | $= 0$ | All | |
| Number of lost outer hits | ≥ 2 | M, L | |
| Is a PF candidate? | No | All | |
| PF lepton veto ($\Delta R < 0.1$) | Yes | All | |
| Lepton veto ($\Delta R < 0.2$) | Yes | All | |
| Track veto ($\Delta R < 0.1$) | Yes | All | |
| Bad calorimeter module veto | Yes | All | |
| M_T (track, \vec{E}_T) [GeV] | > 100 , if $p_T < 150 \text{ GeV}$ | L | |

The efficiency for signal tracks to pass this selection is high, 50–65%, as shown in Figure 3.20.




Figure 3.20: The signal efficiency with respect to the full short track selection after selecting events passing the baseline kinematic signal region selection. Additional material from [11].

Binning

The background for shorter disappearing tracks is dominated by fakes, while the background for longer disappearing tracks is dominated by lost pions. Additionally, signals with different LLP lifetimes produce tracks of different lengths. Thus, binning the signal region by track length allows backgrounds of different origins to be segregated in different bins, and maximizes sensitivity to a variety of signal lifetimes. The disappearing track search adopts 3 categories of tracks length.

- Pixel-only (P) tracks have hits only inside the pixel tracker. Fakes dominate the background.
In 2017 and 2018 data, these tracks are further subdivided into pixel tracks with hits in 3 distinct layers (P3) and pixel tracks with hits in 4 distinct layers (P4).
- Medium length (M) tracks have hits in fewer than 7 distinct layers, but have at least one hit

outside the pixel detector. Lost mesons dominate the background.

- Relatively long (L) tracks have hits in 7 or more distinct layers, but still disappear, having missing expected hits in at least the two outermost layers of the tracker. Lost mesons dominate the background, and the better opportunity to measure the track allows for an especially tight selection, allowing for extremely strong background suppression.

Additionally, P and M tracks are divided into tracks with $15 < p_T < 50$ GeV (“lo”) and tracks with $p_T \geq 50$ GeV (“hi”). Most background falls into the low p_T bins, and signals generically populate the high p_T bins, making the high p_T bins the drivers of sensitivity to most signals. This division is not adopted for L tracks, as statistics are too low to allow further division of that population of tracks.

Finally, events are binned based on the global event kinematics, using N_{jet} and H_T , again partly to segregate backgrounds produced by different signals and partly to enhance sensitivity to a variety of signal models. The N_{jet} bins are $2 < N_{\text{jet}} \leq 3$ (denoted by a trailing L, targeted at squark production), and $N_{\text{jet}} > 4$ (H, targeted at gluino production). The H_T bins are $250 < H_T < 575$ GeV (denoted by a leading “L”), $575 \leq H_T < 1200$ GeV (“M”), and $H_T \geq 1200$ GeV (“H”). For events with L length tracks, the L and M H_T bins are merged into “LM” to preserve statistics.

For example, the “L HLM” region is populated by events with L length tracks, with $N_{\text{jet}} \geq 4$ and $H_T \geq 1200$ GeV. The “M LM lo” region is populated by events with M length tracks of $15 < p_T < 50$ GeV, with $2N_{\text{jet}} \leq 3$ and $250 < H_T < 575$ GeV.

The STC control region binning mirrors the ST signal region, with each STC bin mapping to a corresponding ST bin.

Short Track Candidates and f_{short}

The background estimation procedure makes use of the ratio of the counts of STs and STCs, called f_{short} ,

$$f_{\text{short}} = \frac{N_{\text{ST}}}{N_{\text{STC}}}. \quad (3.5)$$

This ratio is measured in events with $60 < M_{\text{T2}} < 100$ GeV, as these events are so background-dominated that even STs are assuredly background-dominated. A separate f_{short} is measured for every bin, except that the measurement is inclusive in H_T , exploiting an empirical invariance of f_{short} with respect to H_T to improve the statistical precision of the measurement. The ratio is then applied at high M_{T2} to estimate the number of signal-like STs produced by background, N_{ST}^{B} , using the number of STC events observed in the corresponding control region, $N_{\text{STC}}^{\text{Obs}}$,

$$N_{\text{ST}}^{\text{B}} = f_{\text{short}} N_{\text{STC}}^{\text{Obs}}. \quad (3.6)$$

This estimate is correct up to statistical fluctuations, subject to the assumption that f_{short} is invariant with respect to M_{T2} , an assumption which is validated in data in the next subsection. Even without this explicit validation, there is no reason to expect that a ratio based on track-level observables, namely the impact parameter, p_T error, and isolation, ought to be sensitive to a global event kinematic variable like M_{T2} . By tethering the background estimate to the observed count of STCs in data, the analysis is able to remain agnostic of the details of background disappearing track production.

Validation

The f_{short} -based estimate of background ST production relies on the invariance of f_{short} with respect to M_{T2} . The f_{short} ratio is measured at $60 < M_{\text{T2}} < 100$ GeV, a background-dominated region, and applied in the $M_{\text{T2}} > 200$ GeV signal region. In Figure 3.21, the invariance of f_{short} with respect to M_{T2} is validated in the intermediate region, $100 < M_{\text{T2}} < 200$ GeV, in both 2016




Figure 3.21: Validation of the disappearing tracks background estimate in (upper) 2016 and (lower) 2017–2018 data. There are no discrepancies inconsistent with statistical fluctuations. To be conservative, a systematic uncertainty (gray) is assessed region-by-region to account for any discrepancy greater than 1 statistical standard deviation (blue). Dotted vertical lines group bins that use the same measured value of f_{short} , because they differ only in H_T . Taken from [11].

and 2017–2018 data, by carrying out the background estimation procedure as used for the signal region in this M_{T2} sideband. There are no discrepancies inconsistent with statistical fluctuations,

so the estimate is considered validated. To be conservative, a systematic uncertainty is assessed bin-by-bin that is no less than the statistical uncertainty, to prevent any cases of accidental closure, and as large as is necessary to cover any observed discrepancy larger than 1 standard deviation of statistical uncertainty.

3.2.6 Systematics

Table 3.7: Summary of systematic uncertainties in the disappearing track background prediction, together with their typical size ranges across the search bins. The systematic uncertainties arising from the assumption of kinematic invariance of f_{short} and from the validation of the background prediction are always taken to be at least as large as the statistical uncertainties on the measured values of f_{short} and on the background prediction in the validation region, respectively. Taken from [11].

| Source | Range [%] |
|--|-----------|
| Limited size of data control samples (high M_{T2} STC counts) | 1–100 |
| Limited size of data f_{short} measurement samples (low M_{T2} disappearing track counts) | 5–45 |
| Kinematic invariance of f_{short} | 10–80 |
| Validation of background prediction | 25–75 |

The uncertainties uniquely affecting the disappearing tracks the background prediction, in addition to those obtained from the validation region, are summarized in Table 3.7 together with their typical size ranges across the search bins.

The first two uncertainties are purely statistical and dominate the total uncertainty.

The kinematic invariance of f_{short} is checked by varying the H_T and \cancel{E}_T definitions of the f_{short} measurement region. Any uncertainty greater than 1 statistical standard deviation is covered by applying a systematic, just as for the validation region systematic. Although the relative uncertainties thus assessed can be large, this is an artifact of extremely low statistics. Discrepancies easily consistent within 2 statistical standard deviations can nonetheless lead to assessment of a large relative systematic uncertainty. Still, these uncertainties are applied to ensure that the disappearing tracks search is conservative.

As for the classic search, the disappearing tracks signal simulation is subject to extra

uncertainties, all of which are shared except the uncertainties assessed to cover the reconstruction of signal disappearing tracks, produced by decaying $\tilde{\chi}^\pm$. There are three such uncertainties.

First, fast simulation is used for signals simulated in 2017 and 2018 conditions. The selection efficiency for signal tracks in fast simulation differs slightly from full simulation, and so a systematic is assessed to cover the difference.

A 10% uncertainty is assessed to cover any potential mismodeling of how efficiently the detector would reconstruct tracks from a long-lived $\tilde{\chi}^\pm$. This is half the difference between unity and the observed efficiency in simulation, for $\tilde{\chi}^\pm$ decays in the regions of the detector that ought to be able to reconstruct the tracks successfully.

Finally, an uncertainty of 1.5% is assessed to cover observed differences between full simulation and fast simulation in how often a $\tilde{\chi}^\pm$ that reaches the muon chamber is falsely reconstructed as a muon, triggering the lepton veto.

3.2.7 Results

Figure 3.22 shows the observed counts compared with the background estimate for both 2016 and 2017–2018 data. It is worth noting how small the expected background is, especially in the high p_T M and L bins, in which only a few events are observed across the entire 13 TeV dataset recorded from 2016 to 2018, at an event rate on the order of 1 GHz. The background-only hypothesis is consistent with the data, and so these results are interpreted as exclusion limits on the signal models shown in Figure 3.18 with a variety of lifetimes. Since counts in the M and L bins are so small, signals populating the M and L bins need only to have a few expected events to be excluded at 95% CL by these results.

Signal Contamination

As with the contamination of lepton control regions in the classic search (Section 3.1.6), the disappearing tracks search must adjust its expected signal counts for contamination of its con-




Figure 3.22: Taken from [11]

trol regions. In this case, there is some additional complexity because the signal can contaminate the background estimate by multiple routes. Any signal tracks in the f_{short} measurement region at low M_{T2} can bias the measurement of f_{short} , and signal STCs in the high M_{T2} STC control regions can bias the application of f_{short} . We can separate these two factors to express the effect

of contamination bias schematically as,

$$N_{\text{ST}}^{\text{Est}} + N_{\text{Signal}}^{\text{Bias}} = (f_{\text{short}} + \Delta f_{\text{short}})(N_{\text{STC}} + \Delta N_{\text{STC}}) = f_{\text{short}}N_{\text{STC}} + f_{\text{short}}\Delta N_{\text{STC}} + \Delta f_{\text{short}}N_{\text{STC}} + \Delta f_{\text{short}}\Delta N_{\text{STC}} \quad (3.7)$$

The bias is not linear in the signal strength, due both to the manifestly nonlinear final term, $\Delta f_{\text{short}}\Delta N_{\text{STC}}$, and to the hidden nonlinearity in Δf_{short} ,

$$f_{\text{short}} + \Delta f_{\text{short}} = \frac{N_{\text{ST}}^{\text{B}} + N_{\text{ST}}^{\text{S}}}{N_{\text{STC}}^{\text{B}} + N_{\text{STC}}^{\text{S}}} \quad (3.8)$$

(Recall that these quantities are measured in the $60 < M_{\text{T2}} < 100$ GeV sideband for the f_{short} measurement.)

If the overprediction of background due to contamination, $N_{\text{Signal}}^{\text{Bias}}$, is nonlinear, then the contamination factor C is a function of the signal strength μ , which means that the adjusted signal count must be recomputed for every trial signal strength during the statistical analysis,

$$N_{\text{Signal}}^{\text{Adjusted}} = \mu \left(N_{\text{Signal}}^{\text{Raw}} - N_{\text{Signal}}^{\text{Bias}} \right) = \mu \left(1 - \frac{N_{\text{Signal}}^{\text{Bias}}}{N_{\text{Signal}}^{\text{Raw}}} \right) N_{\text{Signal}}^{\text{Raw}} = \mu C(\mu) N_{\text{Signal}}^{\text{Raw}}. \quad (3.9)$$

This is computationally utterly infeasible. Fortunately, $N_{\text{Signal}}^{\text{Bias}}$ is *approximately* linear in μ , and the approximation error is strictly conservative, so that adopting the approximation will never lead to a false discovery or false exclusion.

First, one may approximate Δf_{short} thus,

$$f_{\text{short}} + \Delta f_{\text{short}} = \frac{N_{\text{ST}}^{\text{B}} + N_{\text{ST}}^{\text{S}}}{N_{\text{STC}}^{\text{B}}}. \quad (3.10)$$

The term $N_{\text{STC}}^{\text{S}}$ is doubly small. First, signal is rare relative to background at low M_{T2} , and second, only on the order of 1% of signal tracks fall into the STC category, so that $N_{\text{STC}}^{\text{S}} \ll N_{\text{STC}}^{\text{B}}$ in the f_{short} measurement region. This approximation linearizes Δf_{short} , and strictly makes Δf_{short}

larger by reducing the denominator, tending to overestimate the bias and, therefore, produce conservative signal counts.

Next, adjusting the applied f_{short} to $f'_{\text{short}} = f_{\text{short}} - \Delta f_{\text{short}}$, Equation 3.7 becomes

$$N_{\text{ST}}^{\text{Est}} + N_{\text{Signal}}^{\text{Bias}} = f'_{\text{short}}(N_{\text{STC}} + \Delta N_{\text{STC}}) = f'_{\text{short}}N_{\text{STC}} + f'_{\text{short}}\Delta N_{\text{STC}} \quad (3.11)$$

The second component of $f'_{\text{short}}\Delta N_{\text{STC}} = f_{\text{short}}\Delta N_{\text{STC}} - \Delta f_{\text{short}}\Delta N_{\text{STC}}$ is nonlinear in the signal strength, but is again doubly small, for the same reasons as the first neglected term. Neglecting it again produces a strictly larger estimate of the contamination, making the approximation conservative once again.

Ultimately, then, there are two adjustments to the expected signal counts. First, the analysis subtracts the effect of signal contamination of low M_{T2} STs, which bias the f_{short} measurement. Then, the analysis subtracts the effect of signal contamination of high M_{T2} STCs, which bias the application of f_{short} . Both of these effects are small. Any terms that consider the combined effect of low M_{T2} signal and signal STCs are dropped, as signal is rare in both cases, and so doubly rare in the combined case. This procedure outputs an adjustment to expected signal counts that is linear in the signal strength, and conservative, allowing for a straightforward and safe statistical interpretation.

3.2.8 Limits

The statistical procedure for extracting limits in the disappearing tracks search is exactly identical to that used in the classic search, described in Section 3.1.8, with the addition of the LLP lifetime as a parameter of interest. Figures 3.23-3.25 show limits in the mass plane for gluinos decaying to light squarks, bottom squarks, and top squarks, respectively, with the $\tilde{\chi}^\pm c\tau$ set to 10 cm (upper left), 50 cm (upper right), and 200 cm (bottom). These limits can be compared directly to those obtained by the classic search, shown respectively in Figures 3.11 (upper), 3.13




Figure 3.23: Exclusion limits at 95% CLs for direct gluino pair production where the gluinos decay to light-flavor quarks, with $c\tau_0(\tilde{\chi}^\pm) =$ (upper left) 10 cm, (upper right) 50 cm, and (lower) 200 cm. Taken from [11].




Figure 3.24: Exclusion limits at 95% CL_S for light squark pair production with $c\tau_0(\tilde{\chi}^\pm) =$ (upper left) 10 cm, (upper right) 50 cm, and (lower) 200 cm. Taken from [11].




Figure 3.25: Exclusion limits at 95% CL_S for top squark pair production with $c\tau_0(\tilde{\chi}_1^\pm) =$ (upper left) 10cm (upper right) 50cm, and (lower) 200cm. Taken from [11].




Figure 3.26: Exclusion limits at 95% CL_S on the $\tilde{\chi}_1^\pm$ mass, with $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0} + \mathcal{O}(100 \text{ MeV})$, as a function of the $\tilde{\chi}_1^\pm$ proper decay length, for (upper) direct gluino and (lower) direct light-flavor squark pair production, for representative gluino and squark masses. Taken from [11].




Figure 3.27: Exclusion limits at 95% CL_S on the $\tilde{\chi}_1^0$ mass, with $m_{\tilde{\chi}^\pm} = m_{\tilde{\chi}_1^0} + \mathcal{O}(100 \text{ MeV})$, as a function of the $\tilde{\chi}^\pm$ proper decay length, for direct top squark pair production, as obtained for a representative top squark mass. Taken from [11].

(upper left), and 3.14 (upper right). In each case, the limits improve by hundreds of GeV along each axis, due entirely to the strong background suppression of the ST requirement.

An interesting feature of these exclusion curves is evident when comparing longer to shorter $\tilde{\chi}^\pm$ lifetimes. For $c\tau = 200$ cm, the limit curve abruptly turns inward at the bottom right of each plot, where the gluino or squark is much more massive than $\tilde{\chi}^\pm$ and $\tilde{\chi}_1^0$. At $c\tau = 50$ cm, this feature is reduced in prominence, and it is inverted when $c\tau = 10$ cm. This is caused by the large Lorentz boost and correspondingly extended lab frame lifetime of $\tilde{\chi}^\pm$ in these mass scenarios. At longer lifetimes, $\tilde{\chi}^\pm$ often lives long enough to escape the tracker, or at least make it near enough to the edge not to leave a disappearing track, defined as a track with at least two tracker layers missing expected hits. With no disappearing track, the event fails the ST selection, so the signal efficiency for long $\tilde{\chi}^\pm$ lifetime is reduced in the large mass splitting scenario, and limits are weakened. In contrast, shorter lifetimes benefit from the boost, as tracks more often extend into the more sensitive M and L bins. Masses of $\tilde{\chi}^\pm$ below 91.9 GeV are excluded by searches at CERN’s Large Electron Positron collider [21], while previously occupied the LHC




Figure 3.28: Exclusion limits at 95% CLs on $\sigma/\sigma_{\text{theory}}$ as a function of the $\tilde{\chi}_1^\pm$ decay length, for (upper) gluino pair production where the gluinos decay to light-flavor quarks, (lower left) light-flavor squark pair production, and (lower right) top squark pair production, as obtained from the search for disappearing tracks. Taken from [11].

tunnel, mitigating the impact of this effect.

Figures 3.26 and 3.27 show limits at 95% CL on the $\tilde{\chi}_1^0$ mass as a function of the lifetime of $\tilde{\chi}^\pm$, at representative fixed masses of the considered gluino or squark, with the 68% and 95% uncertainty bands shown in green and yellow, respectively. Figure 3.26 (upper) considers gluino pair production at gluino mass 1900 GeV. Figure 3.26 (lower left) considers light squark pair production in the case of a single light squark of mass 900 GeV. Figure 3.26 (lower right) considers light squark pair production in the case of eight degenerate light squarks of mass 1500 GeV. Figure 3.27 considers top squark pair production and at top squark mass 1000 GeV.

There is a discontinuity at low $\tilde{\chi}^\pm$ lifetime in the exclusion curves of all four of these figures. At this discontinuity, the limits obtained for the classic search, applicable to any $\tilde{\chi}^\pm$ of lifetime short enough not to reach the calorimeters, become superior to those obtained by the disappearing track search. At these very short lifetimes, many of the charginos decay before reaching the tracker or soon after, and so do not produce tracks. As the disappearing track search only selects events with a disappearing track, it suffers from large selection inefficiencies at these very short lifetimes. The classic search does not require a track, and so avoids this selection inefficiency, at the cost of a larger background. For short enough $\tilde{\chi}^\pm$ lifetimes, to the left of the discontinuity in each figure, this tradeoff becomes advantageous as measured by the expected sensitivity.

When the exclusion curves approach the kinematic limit, at which the mass of the decaying particle and the mass of the $\tilde{\chi}^\pm$ are nearly equal, the errors tend to become greatly compressed.

Finally, Figure 3.28 show the exclusion limits at 95% CL on the production cross section of gluinos and squarks as a function of $\tilde{\chi}^\pm$ lifetime for fixed gluino or squark mass and $\tilde{\chi}_1^0$ mass, for gluinos of mass 1600 GeV decaying to light quarks and $\tilde{\chi}_1^0$ of mass 1575 GeV (upper), light squarks of mass 2000 GeV and $\tilde{\chi}_1^0$ mass 1000 GeV (lower left), and top squarks of mass 1100 GeV and $\tilde{\chi}_1^0$ mass 1000 GeV (lower right). The limits are strongest for intermediate $\tilde{\chi}^\pm$ lifetimes for which the fraction of $\tilde{\chi}^\pm$ decays producing disappearing tracks tens of centimeters

long is maximized. Sensitivity is lesser for longer and shorter lifetimes due to reduced efficiency to produce disappearing tracks, and consequently reduced signal selection efficiency.

All of these limits are the strongest produced to date by any analysis, and constitute stringent constraints on the entire class of signal models discussed in Section 3.2.3.

3.2.9 Future

The subject of LLPs in supersymmetric extensions of the Standard Model, and the potential sensitivity gains that LLPs can provide for new physics searches in general, will continue to inspire interest for the foreseeable future of the LHC’s physics program. The M_{T2} disappearing track extension is no exception. Furthermore, unlike the classic search, which has non-negligible backgrounds that will grow larger with increased integrated luminosity luminosity, the most sensitive bins of the disappearing tracks search are effectively entirely background-depleted. In fact, while the background should not meaningfully increase in the foreseeable future, the statistical precision of the background estimate *will* improve with the addition of more data. As a result, the sensitivity of the disappearing track search projects to improve roughly linearly with increasing integrated luminosity rather than as its square root. With continuing interest and better scaling of the sensitivity, this search is a potentially worthwhile candidate for continued updates as the CMS dataset expands.

Additionally, the disappearing tracks search in its current form targets only tracks that disappear inside the silicon tracker, and does not consider the case of tracks disappearing inside the muon system, as mentioned in Section 2.2.5. Indeed, it explicitly vetoes tracks potentially associated with hits in the muon system. However, sufficiently long-lived charginos would leave hits in the muon system, and so there exists a possibility to extend the lifetime sensitivity range of the disappearing tracks search by expanding the definition to include disappearing tracks in the muon system, or even non-disappearing tracks in the muon system that appear to be produced by a particle too heavy to be consistent with a muon.

In any case, the disappearing tracks signature certainly has not been fully exploited, and will remain an interesting search strategy in future LHC datasets.

Chapter 4

Conclusions

SM is robust at low energy physics but has some issues that may have consequences at the weak scale. LHC and CMS are powerful machines that can investigate the weak scale. Performed a pair of searches that are sensitive to pair-produced particles decaying to dark matter, one of these issues, that can constrain SUSY, a solution to the hierarchy problem. One of these searches targets long-lived particles producing disappearing, a challenging final state to study. Produced strongest constraints to date.

Appendix A

Full Classic Binning and Results

Table A.1: Predictions and observations for the 12 search regions with $N_{\text{jet}} = 1$. For each of the background predictions, the first uncertainty listed is statistical (from the limited size of data control samples and Monte Carlo samples), and the second is systematic. Reprinted from [11].

| $N_{\text{jet}} = 1$ | | | | | | |
|------------------------------------|---------------------------|------------------------------|-------------------------------|--------------------------|------------------------------------|---------------|
| $N_{\text{jet}}, N_{\text{b-tag}}$ | $p_T^{\text{jet}1}$ [GeV] | Lost lepton | $Z \rightarrow \nu\nu$ | Multijet | Total background | Data |
| 1j, 0b | 250-350 | $70700 \pm 400 \pm 4100$ | $167000 \pm 1000 \pm 11000$ | $530 \pm 20 \pm 160$ | 238000 $\pm 1000 \pm 14000$ | 251941 |
| | 350-450 | $13440 \pm 130 \pm 790$ | $40100 \pm 500 \pm 3100$ | $55 \pm 5 \pm 16$ | 53600 $\pm 500 \pm 3700$ | 54870 |
| | 450-575 | $3050 \pm 50 \pm 180$ | $10850^{+230}_{-220} \pm 690$ | $5.6 \pm 1.1 \pm 1.6$ | 13910 $\pm 230 \pm 840$ | 14473 |
| | 575-700 | $603^{+20}_{-19} \pm 38$ | $2590^{+110}_{-100} \pm 160$ | $0.38 \pm 0.06 \pm 0.11$ | 3200 $\pm 110 \pm 190$ | 3432 |
| | 700-1000 | $220 \pm 13 \pm 16$ | $1076^{+70}_{-66} \pm 66$ | $0.12 \pm 0.03 \pm 0.03$ | $1295^{+71}_{-67} \pm 79$ | 1304 |
| | 1000-1200 | $11.7^{+4.1}_{-3.2} \pm 0.9$ | $86^{+23}_{-19} \pm 6$ | < 0.01 | $98^{+24}_{-19} \pm 7$ | 98 |
| | ≥ 1200 | $2.8^{+2.7}_{-1.5} \pm 0.6$ | $23^{+12}_{-8} \pm 2$ | < 0.01 | $26^{+13}_{-9} \pm 2$ | 30 |
| 1j, $\geq 1b$ | 250-350 | $4210 \pm 110 \pm 260$ | $9030 \pm 230 \pm 630$ | $58 \pm 10 \pm 17$ | $13310^{+260}_{-250} \pm 820$ | 13549 |
| | 350-450 | $878 \pm 38 \pm 56$ | $2180^{+110}_{-100} \pm 170$ | $4.6 \pm 0.4 \pm 1.3$ | 3060 $\pm 110 \pm 220$ | 3078 |
| | 450-575 | $211^{+16}_{-15} \pm 13$ | $651^{+57}_{-53} \pm 44$ | $0.63 \pm 0.18 \pm 0.18$ | $863^{+59}_{-55} \pm 53$ | 810 |
| | 575-700 | $40.3^{+6.0}_{-5.5} \pm 2.5$ | $164^{+30}_{-26} \pm 11$ | $0.04 \pm 0.02 \pm 0.02$ | $205^{+31}_{-26} \pm 13$ | 184 |
| | ≥ 700 | $19.2^{+5.7}_{-4.6} \pm 1.3$ | $74^{+21}_{-16} \pm 7$ | < 0.01 | $94^{+21}_{-17} \pm 7$ | 83 |

Table A.2: Predictions and observations for the 30 search regions with $250 \leq H_T < 450$ GeV. For each of the background predictions, the first uncertainty listed is statistical (from the limited size of data control samples and Monte Carlo samples), and the second is systematic. Reprinted from [11].

| 250 $\leq H_T < 450$ GeV | | | | | | |
|------------------------------------|------------------------------|---------------------------------|---------------------------------|--------------------------|---------------------------------|---------------|
| $N_{\text{jet}}, N_{\text{b-tag}}$ | $M_{\text{T2}} [\text{GeV}]$ | Lost lepton | $Z \rightarrow \nu\nu$ | Multijet | Total background | Data |
| 2-3j, 0b | 200-300 | $73700 \pm 500 \pm 5000$ | $156000 \pm 1000 \pm 12000$ | $580 \pm 20 \pm 140$ | $231000 \pm 1000 \pm 16000$ | 240867 |
| | 300-400 | $12030 \pm 200 \pm 820$ | $31300 \pm 200 \pm 2500$ | $50 \pm 5 \pm 10$ | $43400 \pm 300 \pm 3200$ | 44074 |
| | ≥ 400 | $417^{+51}_{-47} \pm 28$ | $1450 \pm 10 \pm 140$ | $0.44 \pm 0.09 \pm 0.09$ | $1870 \pm 50 \pm 160$ | 2022 |
| 2-3j, 1b | 200-300 | $12450 \pm 170 \pm 820$ | $18700 \pm 300 \pm 1500$ | $90 \pm 8 \pm 21$ | $31300 \pm 300 \pm 2200$ | 32120 |
| | 300-400 | $2380 \pm 80 \pm 160$ | $3750 \pm 60 \pm 310$ | $6.9 \pm 1.0 \pm 1.5$ | $6130 \pm 100 \pm 430$ | 6258 |
| | ≥ 400 | $97 \pm 8 \pm 39$ | $174 \pm 3 \pm 17$ | $0.01 \pm 0.01 \pm 0.00$ | $271^{+9}_{-8} \pm 45$ | 275 |
| 2-3j, 2b | 200-300 | $2240 \pm 70 \pm 150$ | $2340^{+110}_{-100} \pm 200$ | $9.7 \pm 1.1 \pm 2.3$ | $4600^{+130}_{-120} \pm 320$ | 4709 |
| | 300-400 | $398^{+34}_{-32} \pm 27$ | $469^{+21}_{-20} \pm 39$ | $0.68 \pm 0.17 \pm 0.15$ | $868^{+40}_{-38} \pm 61$ | 984 |
| | ≥ 400 | $13.3 \pm 2.3 \pm 5.4$ | $21.7^{+1.0}_{-0.9} \pm 2.2$ | < 0.01 | $35.0 \pm 2.5 \pm 6.0$ | 30 |
| 2-6j, $\geq 3b$ | 200-300 | $507^{+32}_{-31} \pm 38$ | $179^{+35}_{-30} \pm 27$ | $1.77 \pm 0.46 \pm 0.46$ | $688^{+47}_{-43} \pm 54$ | 699 |
| | 300-400 | $69 \pm 6 \pm 15$ | $40.0^{+7.8}_{-6.6} \pm 6.0$ | $0.16 \pm 0.12 \pm 0.04$ | $109^{+10}_{-9} \pm 16$ | 102 |
| | ≥ 400 | $1.50 \pm 0.80 \pm 0.61$ | $1.43^{+0.28}_{-0.24} \pm 0.25$ | < 0.01 | $2.92^{+0.85}_{-0.83} \pm 0.67$ | 0 |
| 4-6j, 0b | 200-300 | $12500 \pm 180 \pm 800$ | $21600 \pm 300 \pm 1800$ | $250 \pm 17 \pm 58$ | $34400 \pm 400 \pm 2400$ | 35187 |
| | 300-400 | $2070 \pm 80 \pm 130$ | $4660 \pm 70 \pm 410$ | $18.2 \pm 3.6 \pm 3.8$ | $6750 \pm 110 \pm 510$ | 6725 |
| | ≥ 400 | $42 \pm 5 \pm 17$ | $155 \pm 2 \pm 64$ | $0.06 \pm 0.03 \pm 0.01$ | $197 \pm 5 \pm 67$ | 170 |
| 4-6j, 1b | 200-300 | $5750 \pm 100 \pm 380$ | $4300 \pm 150 \pm 360$ | $61 \pm 7 \pm 15$ | $10120 \pm 180 \pm 680$ | 10564 |
| | 300-400 | $784^{+43}_{-42} \pm 52$ | $928^{+32}_{-31} \pm 84$ | $2.07 \pm 0.29 \pm 0.45$ | $1710 \pm 50 \pm 120$ | 1769 |
| | ≥ 400 | $14.0 \pm 2.5 \pm 5.7$ | $31 \pm 1 \pm 13$ | $0.04 \pm 0.02 \pm 0.01$ | $45 \pm 3 \pm 14$ | 40 |
| 4-6j, 2b | 200-300 | $2550^{+70}_{-60} \pm 170$ | $921^{+68}_{-63} \pm 87$ | $10.0 \pm 1.5 \pm 2.2$ | $3480 \pm 90 \pm 230$ | 3621 |
| | 300-400 | $220^{+23}_{-21} \pm 15$ | $198^{+15}_{-14} \pm 20$ | $0.47 \pm 0.15 \pm 0.11$ | $419^{+27}_{-25} \pm 31$ | 496 |
| | ≥ 400 | $3.2 \pm 0.8 \pm 1.3$ | $6.6 \pm 0.5 \pm 2.7$ | < 0.01 | $9.8 \pm 0.9 \pm 3.1$ | 14 |
| $\geq 7j, 0b$ | 200-300 | $55^{+15}_{-13} \pm 4$ | $61^{+23}_{-17} \pm 26$ | $2.64 \pm 0.39 \pm 0.57$ | $119^{+28}_{-22} \pm 27$ | 108 |
| | 300-500 | $3.8^{+2.1}_{-2.0} \pm 0.8$ | $8.1^{+3.1}_{-2.3} \pm 4.3$ | $0.08 \pm 0.04 \pm 0.02$ | $12.0^{+3.7}_{-3.1} \pm 4.4$ | 30 |
| | ≥ 500 | $0.0^{+3.2}_{-0.0} \pm 0.0$ | $0.0^{+1.2}_{-0.0} \pm 0.0$ | < 0.01 | $0.0^{+3.4}_{-0.0} \pm 0.0$ | 0 |
| $\geq 7j, 1b$ | 200-300 | $48.0^{+9.1}_{-8.2} \pm 3.5$ | $19^{+19}_{-11} \pm 10$ | $0.33 \pm 0.14 \pm 0.09$ | $68^{+21}_{-13} \pm 11$ | 95 |
| | ≥ 300 | $3.0 \pm 1.4 \pm 1.2$ | $2.5^{+2.4}_{-1.3} \pm 1.7$ | $0.03 \pm 0.02 \pm 0.01$ | $5.6^{+2.8}_{-1.9} \pm 2.1$ | 12 |
| $\geq 7j, 2b$ | 200-300 | $41.3^{+7.7}_{-7.0} \pm 3.1$ | $6.0^{+5.8}_{-3.2} \pm 3.7$ | $0.29 \pm 0.14 \pm 0.06$ | $47.6^{+9.7}_{-7.7} \pm 5.0$ | 30 |
| | ≥ 300 | $2.15^{+0.78}_{-0.76} \pm 0.87$ | $0.74^{+0.72}_{-0.40} \pm 0.57$ | < 0.01 | $2.9^{+1.1}_{-0.9} \pm 1.1$ | 1 |
| $\geq 7j, \geq 3b$ | 200-300 | $7.3^{+1.7}_{-1.5} \pm 0.9$ | $1.0^{+1.0}_{-0.6} \pm 1.1$ | $0.04 \pm 0.04 \pm 0.01$ | $8.4^{+1.9}_{-1.6} \pm 1.5$ | 17 |
| | ≥ 300 | $0.47 \pm 0.35 \pm 0.20$ | $0.12^{+0.11}_{-0.06} \pm 0.14$ | < 0.01 | $0.59^{+0.37}_{-0.35} \pm 0.24$ | 0 |

Table A.3: Predictions and observations for the 40 search regions with $450 \leq H_T < 575$ GeV. For each of the background predictions, the first uncertainty listed is statistical (from the limited size of data control samples and Monte Carlo samples), and the second is systematic. Reprinted from [11].

| $450 \leq H_T < 575$ GeV | | | | | | |
|------------------------------------|----------------|---------------------------------|---------------------------------|--------------------------|--|--------------|
| $N_{\text{jet}}, N_{\text{b-tag}}$ | M_{T2} [GeV] | Lost lepton | $Z \rightarrow vv$ | Multijet | Total background | Data |
| 2-3j, 0b | 200-300 | $8860 \pm 110 \pm 640$ | $20100 \pm 200 \pm 1300$ | $69 \pm 13 \pm 16$ | 29100 $\pm 300 \pm 1900$ | 28956 |
| | 300-400 | $4230 \pm 80 \pm 300$ | $11770 \pm 140 \pm 790$ | $10.6 \pm 0.8 \pm 2.4$ | 16000 $\pm 200 \pm 1000$ | 15876 |
| | 400-500 | $1510 \pm 60 \pm 110$ | $5020 \pm 60 \pm 360$ | $2.86 \pm 0.62 \pm 0.60$ | 6540 $\pm 80 \pm 440$ | 6527 |
| | ≥ 500 | $121^{+24}_{-21} \pm 9$ | $580 \pm 7 \pm 63$ | $0.07 \pm 0.03 \pm 0.02$ | 701^{+25}_{-22} ± 68 | 740 |
| 2-3j, 1b | 200-300 | $1326 \pm 43 \pm 88$ | $2500 \pm 80 \pm 170$ | $17.0 \pm 8.4 \pm 3.8$ | 3840^{+100}_{-90} ± 240 | 3859 |
| | 300-400 | $737 \pm 35 \pm 49$ | $1464^{+49}_{-48} \pm 99$ | $1.62 \pm 0.20 \pm 0.43$ | 2200 $\pm 60 \pm 140$ | 2065 |
| | 400-500 | $259^{+25}_{-23} \pm 19$ | $626^{+21}_{-20} \pm 45$ | $0.49 \pm 0.10 \pm 0.12$ | 885^{+32}_{-31} ± 58 | 907 |
| | ≥ 500 | $19.1^{+2.8}_{-2.7} \pm 7.8$ | $72.4 \pm 2.4 \pm 7.9$ | $0.04 \pm 0.02 \pm 0.02$ | 92 $\pm 4 \pm 11$ | 79 |
| 2-3j, 2b | 200-300 | $201 \pm 15 \pm 13$ | $322^{+31}_{-28} \pm 25$ | $1.34 \pm 0.62 \pm 0.47$ | 524^{+35}_{-32} ± 35 | 463 |
| | 300-400 | $83.8^{+9.6}_{-9.1} \pm 9.1$ | $188^{+18}_{-17} \pm 15$ | $0.26 \pm 0.07 \pm 0.07$ | 272^{+21}_{-19} ± 20 | 304 |
| | 400-500 | $31.8^{+4.1}_{-4.0} \pm 6.7$ | $80.4^{+7.7}_{-7.1} \pm 6.6$ | $0.02 \pm 0.01 \pm 0.01$ | 112^{+9}_{-8} ± 10 | 120 |
| | ≥ 500 | $2.16^{+0.67}_{-0.66} \pm 0.88$ | $9.3^{+0.9}_{-0.8} \pm 1.1$ | < 0.01 | 11.4 $\pm 1.1 \pm 1.4$ | 15 |
| 2-6j, $\geq 3b$ | 200-300 | $232^{+17}_{-16} \pm 15$ | $57^{+17}_{-13} \pm 7$ | $2.20 \pm 0.70 \pm 0.80$ | 291^{+24}_{-21} ± 19 | 297 |
| | 300-400 | $81^{+12}_{-11} \pm 6$ | $33.6^{+9.9}_{-7.8} \pm 4.3$ | $0.26 \pm 0.08 \pm 0.08$ | 115^{+16}_{-14} ± 8 | 76 |
| | 400-500 | $10.7^{+2.1}_{-2.0} \pm 2.3$ | $11.4^{+3.4}_{-2.7} \pm 1.5$ | < 0.01 | 22.1^{+4.0}_{-3.4} ± 2.8 | 24 |
| | ≥ 500 | $1.08 \pm 0.58 \pm 0.44$ | $1.03^{+0.30}_{-0.24} \pm 0.17$ | < 0.01 | 2.11^{+0.65}_{-0.62} ± 0.48 | 0 |
| 4-6j, 0b | 200-300 | $5660 \pm 90 \pm 370$ | $8560 \pm 170 \pm 600$ | $143 \pm 7 \pm 35$ | 14360 $\pm 190 \pm 890$ | 15047 |
| | 300-400 | $2250 \pm 60 \pm 150$ | $4790^{+100}_{-90} \pm 350$ | $24.3 \pm 2.6 \pm 6.2$ | 7060 $\pm 110 \pm 460$ | 6939 |
| | 400-500 | $428^{+32}_{-30} \pm 28$ | $1220 \pm 20 \pm 110$ | $1.42 \pm 0.21 \pm 0.52$ | 1650 $\pm 40 \pm 130$ | 1817 |
| | ≥ 500 | $14.8 \pm 2.2 \pm 6.0$ | $86 \pm 2 \pm 35$ | $0.04 \pm 0.02 \pm 0.01$ | 101 $\pm 3 \pm 36$ | 104 |
| 4-6j, 1b | 200-300 | $2810 \pm 60 \pm 190$ | $1880 \pm 80 \pm 130$ | $63 \pm 15 \pm 19$ | 4750 $\pm 100 \pm 300$ | 4736 |
| | 300-400 | $937 \pm 36 \pm 63$ | $1054^{+45}_{-43} \pm 78$ | $5.4 \pm 0.4 \pm 1.4$ | 2000 $\pm 60 \pm 130$ | 2039 |
| | 400-500 | $138^{+17}_{-16} \pm 10$ | $269 \pm 11 \pm 25$ | $0.36 \pm 0.10 \pm 0.10$ | 407^{+20}_{-19} ± 31 | 403 |
| | ≥ 500 | $7.5 \pm 2.2 \pm 3.0$ | $19.1 \pm 0.8 \pm 7.9$ | $0.01 \pm 0.01 \pm 0.00$ | 26.5 $\pm 2.3 \pm 8.5$ | 27 |
| 4-6j, 2b | 200-300 | $1343^{+38}_{-37} \pm 89$ | $414^{+39}_{-35} \pm 33$ | $11.5 \pm 1.0 \pm 3.3$ | 1770 $\pm 50 \pm 110$ | 1767 |
| | 300-400 | $418^{+24}_{-23} \pm 29$ | $232^{+22}_{-20} \pm 19$ | $1.35 \pm 0.35 \pm 0.39$ | 651^{+32}_{-31} ± 43 | 636 |
| | 400-500 | $45.6^{+3.9}_{-3.8} \pm 9.6$ | $59.1^{+5.5}_{-5.1} \pm 5.9$ | $0.03 \pm 0.02 \pm 0.01$ | 105^{+7}_{-6} ± 12 | 120 |
| | ≥ 500 | $1.59 \pm 0.89 \pm 0.65$ | $4.2 \pm 0.4 \pm 1.7$ | < 0.01 | 5.8 $\pm 1.0 \pm 1.9$ | 7 |
| $\geq 7j, 0b$ | 200-300 | $149^{+17}_{-16} \pm 13$ | $169^{+31}_{-27} \pm 34$ | $11.5 \pm 0.8 \pm 3.0$ | 329^{+36}_{-31} ± 38 | 354 |
| | 300-400 | $38.9^{+5.8}_{-5.6} \pm 8.2$ | $64^{+12}_{-10} \pm 17$ | $1.24 \pm 0.42 \pm 0.32$ | 104^{+13}_{-12} ± 20 | 110 |
| | ≥ 400 | $1.28 \pm 0.82 \pm 0.52$ | $8.8^{+1.6}_{-1.4} \pm 3.8$ | $0.03 \pm 0.02 \pm 0.01$ | 10.1^{+1.8}_{-1.6} ± 3.8 | 10 |
| $\geq 7j, 1b$ | 200-300 | $191^{+13}_{-12} \pm 15$ | $67^{+19}_{-15} \pm 15$ | $4.4 \pm 0.5 \pm 1.2$ | 262^{+23}_{-19} ± 23 | 268 |
| | 300-400 | $37.8^{+3.4}_{-3.3} \pm 8.0$ | $25.3^{+7.2}_{-5.7} \pm 7.3$ | $0.30 \pm 0.07 \pm 0.08$ | 63^{+8}_{-7} ± 11 | 65 |
| | ≥ 400 | $2.31 \pm 0.69 \pm 0.94$ | $3.5^{+1.0}_{-0.8} \pm 1.5$ | $0.01 \pm 0.01 \pm 0.00$ | 5.8^{+1.2}_{-1.0} ± 1.8 | 3 |
| $\geq 7j, 2b$ | 200-300 | $173^{+12}_{-11} \pm 13$ | $19.9^{+5.7}_{-4.5} \pm 5.2$ | $1.24 \pm 0.18 \pm 0.33$ | 194^{+13}_{-12} ± 15 | 197 |
| | 300-400 | $26.8 \pm 2.6 \pm 5.7$ | $7.6^{+2.2}_{-1.7} \pm 2.4$ | $0.09 \pm 0.04 \pm 0.03$ | 34.6^{+3.4}_{-3.1} ± 6.3 | 44 |
| | ≥ 400 | $1.40 \pm 0.44 \pm 0.57$ | $1.02^{+0.29}_{-0.23} \pm 0.46$ | < 0.01 | 2.42^{+0.53}_{-0.49} ± 0.73 | 3 |
| $\geq 7j, \geq 3b$ | 200-300 | $55.4^{+4.8}_{-4.7} \pm 7.3$ | $2.3^{+0.7}_{-0.6} \pm 1.1$ | $0.15 \pm 0.06 \pm 0.06$ | 57.8^{+4.8}_{-4.7} ± 7.4 | 37 |
| | 300-400 | $6.4 \pm 1.2 \pm 1.5$ | $0.86^{+0.25}_{-0.20} \pm 0.46$ | $0.01 \pm 0.01 \pm 0.00$ | 7.3 $\pm 1.2 \pm 1.6$ | 9 |
| | ≥ 400 | $0.06 \pm 0.01 \pm 0.03$ | $0.12 \pm 0.03 \pm 0.06$ | < 0.01 | 0.18^{+0.04}_{-0.03} ± 0.07 | 0 |

Table A.4: Predictions and observations for the 47 search regions with $575 \leq H_T < 1200$ GeV, $N_{\text{jet}} < 7$. For each of the background predictions, the first uncertainty listed is statistical (from the limited size of data control samples and Monte Carlo samples), and the second is systematic. Reprinted from [11].

| $575 \leq H_T < 1200$ GeV, $N_{\text{jet}} < 7$ | | | | | | |
|---|------------------------------|---------------------------------|---------------------------------|--------------------------|---|--------------|
| $N_{\text{jet}}, N_{\text{b-tag}}$ | $M_{\text{T2}} [\text{GeV}]$ | Lost lepton | $Z \rightarrow vv$ | Multijet | Total background | Data |
| 2-3j, 0b | 200-300 | $5270 \pm 60 \pm 370$ | $11550 \pm 160 \pm 790$ | $93 \pm 20 \pm 30$ | 16900 $\pm 200 \pm 1100$ | 17256 |
| | 300-400 | $2560 \pm 50 \pm 180$ | $7770^{+110}_{-100} \pm 540$ | $11.9 \pm 1.3 \pm 4.4$ | 10340 $^{+120}_{-110} \pm 680$ | 10145 |
| | 400-500 | $1101^{+32}_{-31} \pm 77$ | $3900 \pm 50 \pm 280$ | $1.33 \pm 0.24 \pm 0.41$ | 5000 $\pm 60 \pm 340$ | 5021 |
| | 500-600 | $502^{+24}_{-23} \pm 35$ | $2250 \pm 30 \pm 170$ | $0.37 \pm 0.07 \pm 0.12$ | 2760 $\pm 40 \pm 200$ | 2706 |
| | 600-700 | $180^{+16}_{-15} \pm 13$ | $746 \pm 10 \pm 73$ | $0.09 \pm 0.03 \pm 0.03$ | 926 $^{+19}_{-18} \pm 80$ | 1066 |
| | 700-800 | $52.1^{+7.3}_{-6.5} \pm 5.5$ | $256 \pm 3 \pm 36$ | $0.01 \pm 0.01 \pm 0.00$ | 308 $^{+8}_{-7} \pm 38$ | 347 |
| | 800-900 | $17.7^{+2.6}_{-2.3} \pm 2.2$ | $107 \pm 1 \pm 20$ | < 0.01 | 125 $\pm 3 \pm 21$ | 111 |
| | 900-1000 | $6.0 \pm 0.9 \pm 1.3$ | $39.4 \pm 0.5 \pm 8.5$ | $0.01 \pm 0.01 \pm 0.00$ | 45.4 $^{+1.1}_{-1.0} \pm 8.7$ | 39 |
| | 1000-1100 | $3.3^{+1.1}_{-1.0} \pm 1.0$ | $13.3 \pm 0.2 \pm 3.9$ | < 0.01 | 16.6 $\pm 1.1 \pm 4.1$ | 11 |
| | ≥ 1100 | $0.31^{+0.09}_{-0.08} \pm 0.12$ | $2.5 \pm 0.0 \pm 1.1$ | < 0.01 | 2.8 $\pm 0.1 \pm 1.1$ | 2 |
| 2-3j, 1b | 200-300 | $826^{+27}_{-26} \pm 54$ | $1480^{+60}_{-50} \pm 100$ | $38 \pm 15 \pm 12$ | 2340 $\pm 60 \pm 140$ | 2499 |
| | 300-400 | $426^{+21}_{-20} \pm 28$ | $994^{+38}_{-37} \pm 69$ | $2.33 \pm 0.26 \pm 0.84$ | 1422 $^{+43}_{-42} \pm 90$ | 1366 |
| | 400-600 | $282^{+18}_{-17} \pm 20$ | $788^{+30}_{-29} \pm 55$ | $0.27 \pm 0.06 \pm 0.10$ | 1071 $^{+35}_{-34} \pm 69$ | 1057 |
| | 600-800 | $43.5^{+3.2}_{-3.1} \pm 6.5$ | $129 \pm 5 \pm 12$ | < 0.01 | 172 $\pm 6 \pm 15$ | 225 |
| | 800-1000 | $4.6 \pm 0.7 \pm 1.3$ | $18.8 \pm 0.7 \pm 3.3$ | < 0.01 | 23.4 $\pm 1.0 \pm 3.6$ | 22 |
| | ≥ 1000 | $0.34 \pm 0.08 \pm 0.14$ | $2.05 \pm 0.08 \pm 0.90$ | < 0.01 | 2.38 $\pm 0.11 \pm 0.91$ | 1 |
| 2-3j, 2b | 200-300 | $105.1^{+9.2}_{-8.7} \pm 7.6$ | $181^{+20}_{-18} \pm 15$ | $3.8 \pm 0.5 \pm 1.3$ | 290 $^{+22}_{-20} \pm 20$ | 316 |
| | 300-400 | $55.0^{+6.7}_{-6.3} \pm 7.5$ | $122^{+14}_{-12} \pm 10$ | $0.27 \pm 0.06 \pm 0.10$ | 177 $^{+15}_{-14} \pm 14$ | 159 |
| | 400-600 | $36.5^{+4.6}_{-4.3} \pm 5.5$ | $97^{+11}_{-10} \pm 8$ | $0.08 \pm 0.03 \pm 0.03$ | 133 $^{+12}_{-11} \pm 11$ | 107 |
| | 600-800 | $4.7 \pm 0.8 \pm 1.3$ | $15.8^{+1.8}_{-1.6} \pm 1.6$ | < 0.01 | 20.6 $^{+1.9}_{-1.8} \pm 2.2$ | 21 |
| | ≥ 800 | $0.59 \pm 0.19 \pm 0.24$ | $2.56^{+0.29}_{-0.26} \pm 0.45$ | < 0.01 | 3.14 $^{+0.35}_{-0.32} \pm 0.52$ | 1 |
| 2-6j, $\geq 3b$ | 200-300 | $299^{+17}_{-16} \pm 22$ | $73^{+15}_{-13} \pm 10$ | $6.2 \pm 0.4 \pm 2.1$ | 379 $^{+22}_{-21} \pm 28$ | 345 |
| | 300-400 | $100 \pm 10 \pm 7$ | $43.5^{+8.8}_{-7.4} \pm 6.2$ | $0.68 \pm 0.09 \pm 0.24$ | 144 $^{+14}_{-12} \pm 11$ | 132 |
| | 400-600 | $32.5^{+6.3}_{-5.6} \pm 2.5$ | $31.2^{+6.3}_{-5.3} \pm 4.4$ | $0.08 \pm 0.03 \pm 0.03$ | 63.8 $^{+8.9}_{-7.7} \pm 5.8$ | 48 |
| | 600-800 | $3.16^{+0.95}_{-0.90} \pm 0.68$ | $5.4^{+1.1}_{-0.9} \pm 0.8$ | < 0.01 | 8.6 $^{+1.4}_{-1.3} \pm 1.1$ | 4 |
| | ≥ 800 | $0.10 \pm 0.03 \pm 0.04$ | $0.71^{+0.14}_{-0.12} \pm 0.15$ | < 0.01 | 0.81 $^{+0.15}_{-0.12} \pm 0.16$ | 0 |
| 4-6j, 0b | 200-300 | $6280 \pm 70 \pm 420$ | $9470 \pm 160 \pm 650$ | $360 \pm 20 \pm 110$ | 16100 $\pm 180 \pm 1000$ | 16292 |
| | 300-400 | $2700 \pm 50 \pm 180$ | $5410 \pm 90 \pm 380$ | $53 \pm 1 \pm 17$ | 8160 $\pm 100 \pm 520$ | 8330 |
| | 400-500 | $927^{+28}_{-27} \pm 62$ | $2420 \pm 40 \pm 180$ | $7.7 \pm 0.4 \pm 2.4$ | 3350 $\pm 50 \pm 230$ | 3576 |
| | 500-600 | $324^{+17}_{-16} \pm 22$ | $1171^{+20}_{-19} \pm 100$ | $1.46 \pm 0.12 \pm 0.46$ | 1500 $\pm 30 \pm 110$ | 1516 |
| | 600-700 | $95.4^{+9.4}_{-8.7} \pm 6.4$ | $413 \pm 7 \pm 47$ | $0.33 \pm 0.06 \pm 0.10$ | 509 $^{+12}_{-11} \pm 50$ | 543 |
| | 700-800 | $35.6^{+5.0}_{-4.5} \pm 3.6$ | $171 \pm 3 \pm 27$ | $0.03 \pm 0.02 \pm 0.01$ | 206 $^{+6}_{-5} \pm 27$ | 178 |
| | 800-900 | $13.4^{+2.0}_{-1.8} \pm 1.6$ | $64 \pm 1 \pm 11$ | $0.02 \pm 0.01 \pm 0.01$ | 77 $\pm 2 \pm 11$ | 62 |
| | 900-1000 | $4.39^{+0.78}_{-0.73} \pm 0.93$ | $23.6 \pm 0.4 \pm 5.3$ | < 0.01 | 28.0 $^{+0.9}_{-0.8} \pm 5.4$ | 20 |
| | 1000-1100 | $0.64 \pm 0.16 \pm 0.20$ | $6.3 \pm 0.1 \pm 2.0$ | < 0.01 | 6.9 $\pm 0.2 \pm 2.0$ | 3 |
| | ≥ 1100 | $0.78 \pm 0.58 \pm 0.32$ | $0.89^{+0.02}_{-0.01} \pm 0.40$ | < 0.01 | 1.68 $\pm 0.58 \pm 0.52$ | 1 |
| 4-6j, 1b | 200-300 | $2900 \pm 50 \pm 200$ | $2220^{+80}_{-70} \pm 150$ | $154 \pm 16 \pm 50$ | 5270 $\pm 90 \pm 330$ | 5335 |
| | 300-400 | $1066 \pm 29 \pm 74$ | $1267^{+44}_{-42} \pm 89$ | $19.2 \pm 0.9 \pm 6.2$ | 2350 $\pm 50 \pm 150$ | 2547 |
| | 400-600 | $504^{+22}_{-21} \pm 35$ | $840^{+29}_{-28} \pm 61$ | $2.98 \pm 0.21 \pm 0.93$ | 1347 $^{+36}_{-35} \pm 88$ | 1284 |
| | 600-800 | $35.3^{+5.9}_{-5.2} \pm 2.6$ | $138 \pm 5 \pm 14$ | $0.09 \pm 0.03 \pm 0.03$ | 174 $^{+8}_{-7} \pm 16$ | 151 |
| | 800-1000 | $3.89^{+0.83}_{-0.77} \pm 0.82$ | $19.3^{+0.7}_{-0.6} \pm 4.3$ | $0.01 \pm 0.01 \pm 0.00$ | 23.2 $^{+1.1}_{-1.0} \pm 4.5$ | 18 |
| | ≥ 1000 | $0.18 \pm 0.07 \pm 0.07$ | $1.57 \pm 0.05 \pm 0.65$ | < 0.01 | 1.75 $\pm 0.09 \pm 0.65$ | 1 |
| 4-6j, 2b | 200-300 | $1500 \pm 30 \pm 100$ | $473^{+36}_{-33} \pm 36$ | $42 \pm 2 \pm 13$ | 2020 $\pm 50 \pm 130$ | 1968 |
| | 300-400 | $508 \pm 20 \pm 35$ | $270^{+20}_{-19} \pm 21$ | $4.9 \pm 0.3 \pm 1.6$ | 783 $^{+29}_{-28} \pm 50$ | 788 |
| | 400-600 | $167 \pm 12 \pm 12$ | $179^{+14}_{-13} \pm 14$ | $0.57 \pm 0.08 \pm 0.18$ | 346 $^{+18}_{-17} \pm 23$ | 354 |
| | 600-800 | $11.9^{+1.3}_{-1.2} \pm 2.5$ | $29.5^{+2.2}_{-2.1} \pm 3.5$ | $0.02 \pm 0.01 \pm 0.01$ | 41.4 $^{+2.6}_{-2.4} \pm 4.6$ | 37 |
| | ≥ 800 | $0.91 \pm 0.23 \pm 0.37$ | $4.4 \pm 0.95 \pm 1.8$ | < 0.01 | 5.4 $\pm 0.4 \pm 1.9$ | 7 |

Table A.5: Predictions and observations for the 34 search regions with $575 \leq H_T < 1200$ GeV, $N_{\text{jet}} \geq 7$. For each of the background predictions, the first uncertainty listed is statistical (from the limited size of data control samples and Monte Carlo samples), and the second is systematic. Reprinted from [11].

| $575 \leq H_T < 1200$ GeV, $N_{\text{jet}} \geq 7$ | | | | | | |
|--|------------------------------|---------------------------------|---------------------------------|--------------------------|---------------------------------|-------------|
| $N_{\text{jet}}, N_{\text{b-tag}}$ | $M_{\text{T2}} [\text{GeV}]$ | Lost lepton | $Z \rightarrow vv$ | Multijet | Total background | Data |
| 7-9j, 0b | 200-300 | $589^{+27}_{-26} \pm 39$ | $573^{+47}_{-43} \pm 64$ | $90 \pm 10 \pm 28$ | $1252^{+55}_{-52} \pm 93$ | 1340 |
| | 300-400 | $265^{+19}_{-18} \pm 18$ | $279^{+23}_{-21} \pm 42$ | $14.9 \pm 0.5 \pm 4.7$ | $559^{+29}_{-28} \pm 51$ | 581 |
| | 400-600 | $92^{+10}_{-9} \pm 6$ | $159^{+13}_{-12} \pm 28$ | $2.72 \pm 0.18 \pm 0.85$ | $253^{+16}_{-15} \pm 30$ | 243 |
| | 600-800 | $8.6 \pm 1.2 \pm 1.8$ | $22.8^{+1.9}_{-1.7} \pm 6.4$ | $0.10 \pm 0.03 \pm 0.03$ | $31.6^{+2.2}_{-2.1} \pm 6.8$ | 32 |
| | ≥ 800 | $0.51 \pm 0.16 \pm 0.21$ | $3.0 \pm 0.2 \pm 1.3$ | < 0.01 | $3.5 \pm 0.3 \pm 1.3$ | 2 |
| 7-9j, 1b | 200-300 | $733 \pm 21 \pm 52$ | $278^{+28}_{-25} \pm 33$ | $48 \pm 3 \pm 16$ | $1059^{+35}_{-33} \pm 73$ | 1052 |
| | 300-400 | $252^{+13}_{-12} \pm 18$ | $135^{+14}_{-12} \pm 21$ | $7.7 \pm 0.4 \pm 2.5$ | $395^{+19}_{-17} \pm 32$ | 387 |
| | 400-600 | $71.3^{+6.9}_{-5.5} \pm 5.2$ | $77^{+8}_{-7} \pm 14$ | $1.36 \pm 0.13 \pm 0.45$ | $150 \pm 10 \pm 16$ | 131 |
| | 600-800 | $4.26^{+0.73}_{-0.71} \pm 0.90$ | $11.0^{+1.1}_{-1.0} \pm 3.1$ | $0.03 \pm 0.02 \pm 0.01$ | $15.3^{+1.3}_{-1.2} \pm 3.3$ | 20 |
| | ≥ 800 | $0.11 \pm 0.04 \pm 0.05$ | $1.48^{+0.15}_{-0.13} \pm 0.63$ | < 0.01 | $1.60^{+0.15}_{-0.14} \pm 0.63$ | 1 |
| 7-9j, 2b | 200-300 | $675 \pm 20 \pm 51$ | $82^{+8}_{-7} \pm 10$ | $20.9 \pm 3.0 \pm 6.7$ | $777^{+22}_{-21} \pm 56$ | 750 |
| | 300-400 | $211 \pm 11 \pm 16$ | $39.8^{+4.0}_{-3.6} \pm 6.4$ | $2.42 \pm 0.19 \pm 0.79$ | $253^{+12}_{-11} \pm 19$ | 259 |
| | 400-600 | $55.4^{+5.5}_{-5.2} \pm 4.2$ | $22.7^{+2.3}_{-2.1} \pm 4.2$ | $0.50 \pm 0.07 \pm 0.16$ | $78.6^{+5.9}_{-5.6} \pm 6.6$ | 72 |
| | 600-800 | $3.00^{+0.63}_{-0.62} \pm 0.64$ | $3.25^{+0.32}_{-0.30} \pm 0.93$ | $0.01 \pm 0.01 \pm 0.01$ | $6.3 \pm 0.7 \pm 1.2$ | 7 |
| | ≥ 800 | $0.27 \pm 0.20 \pm 0.11$ | $0.44 \pm 0.04 \pm 0.19$ | < 0.01 | $0.71 \pm 0.20 \pm 0.22$ | 1 |
| 7-9j, 3b | 200-300 | $185 \pm 8 \pm 18$ | $11.3^{+1.1}_{-1.0} \pm 1.9$ | $3.6 \pm 0.2 \pm 1.2$ | $200 \pm 8 \pm 18$ | 184 |
| | 300-400 | $52.0 \pm 3.8 \pm 5.0$ | $5.5 \pm 0.5 \pm 1.2$ | $0.72 \pm 0.12 \pm 0.26$ | $58.3^{+3.9}_{-3.8} \pm 5.3$ | 59 |
| | 400-600 | $13.6 \pm 1.8 \pm 1.3$ | $3.13^{+0.31}_{-0.29} \pm 0.82$ | $0.05 \pm 0.02 \pm 0.02$ | $16.8 \pm 1.8 \pm 1.6$ | 14 |
| | ≥ 600 | $0.49 \pm 0.21 \pm 0.20$ | $0.51 \pm 0.05 \pm 0.21$ | < 0.01 | $1.00 \pm 0.21 \pm 0.29$ | 2 |
| 7-9j, $\geq 4b$ | 200-300 | $38.8 \pm 3.1 \pm 7.4$ | $2.01^{+0.20}_{-0.18} \pm 0.71$ | $0.55 \pm 0.08 \pm 0.19$ | $41.3^{+3.2}_{-3.1} \pm 7.4$ | 38 |
| | 300-400 | $14.5^{+2.0}_{-1.9} \pm 2.8$ | $0.98^{+0.10}_{-0.09} \pm 0.43$ | $0.06 \pm 0.02 \pm 0.02$ | $15.6^{+2.0}_{-1.9} \pm 2.8$ | 16 |
| | ≥ 400 | $3.75^{+0.98}_{-0.97} \pm 0.70$ | $0.65 \pm 0.06 \pm 0.35$ | < 0.01 | $4.40^{+0.98}_{-0.97} \pm 0.79$ | 3 |
| $\geq 10j$, 0b | 200-300 | $11.5 \pm 1.6 \pm 1.0$ | $4.4^{+0.4}_{-0.3} \pm 2.3$ | $3.1 \pm 0.8 \pm 1.1$ | $19.0 \pm 1.8 \pm 2.8$ | 27 |
| | 300-500 | $5.6 \pm 1.0 \pm 0.5$ | $3.0 \pm 0.2 \pm 1.7$ | $0.55 \pm 0.08 \pm 0.20$ | $9.1 \pm 1.0 \pm 1.8$ | 4 |
| | ≥ 500 | $0.30 \pm 0.11 \pm 0.12$ | $0.44^{+0.04}_{-0.03} \pm 0.24$ | $0.02 \pm 0.01 \pm 0.01$ | $0.76 \pm 0.11 \pm 0.27$ | 3 |
| $\geq 10j$, 1b | 200-300 | $21.0 \pm 1.8 \pm 1.6$ | $3.5 \pm 0.3 \pm 1.9$ | $1.92 \pm 0.18 \pm 0.72$ | $26.4 \pm 1.8 \pm 2.7$ | 32 |
| | 300-500 | $7.7 \pm 1.0 \pm 0.6$ | $2.4 \pm 0.2 \pm 1.4$ | $0.45 \pm 0.07 \pm 0.17$ | $10.5 \pm 1.1 \pm 1.6$ | 15 |
| | ≥ 500 | $0.83^{+0.42}_{-0.41} \pm 0.07$ | $0.36^{+0.04}_{-0.03} \pm 0.20$ | $0.02 \pm 0.01 \pm 0.01$ | $1.20^{+0.42}_{-0.41} \pm 0.22$ | 0 |
| $\geq 10j$, 2b | 200-300 | $21.8 \pm 1.8 \pm 1.6$ | $1.05 \pm 0.10 \pm 0.66$ | $0.64 \pm 0.08 \pm 0.24$ | $23.5 \pm 1.8 \pm 1.8$ | 26 |
| | 300-500 | $8.8 \pm 1.2 \pm 0.6$ | $0.69^{+0.07}_{-0.06} \pm 0.45$ | $0.16 \pm 0.04 \pm 0.06$ | $9.6^{+1.3}_{-1.2} \pm 0.8$ | 9 |
| | ≥ 500 | $0.22 \pm 0.13 \pm 0.02$ | $0.10 \pm 0.01 \pm 0.06$ | < 0.01 | $0.32 \pm 0.13 \pm 0.07$ | 0 |
| $\geq 10j$, 3b | 200-300 | $9.9 \pm 1.3 \pm 1.2$ | $0.25 \pm 0.02 \pm 0.20$ | $0.29 \pm 0.05 \pm 0.12$ | $10.4 \pm 1.3 \pm 1.2$ | 14 |
| | ≥ 300 | $1.59 \pm 0.50 \pm 0.18$ | $0.19 \pm 0.02 \pm 0.16$ | $0.02 \pm 0.01 \pm 0.01$ | $1.80 \pm 0.50 \pm 0.25$ | 2 |
| $\geq 10j$, $\geq 4b$ | ≥ 200 | $3.9 \pm 1.2 \pm 0.8$ | $0.00^{+0.17}_{-0.00} \pm 0.00$ | $0.05 \pm 0.02 \pm 0.02$ | $4.0 \pm 1.2 \pm 0.8$ | 6 |

Table A.6: Predictions and observations for the 37 search regions with $1200 \leq H_T < 1500$ GeV, $N_{\text{jet}} < 7$. For each of the background predictions, the first uncertainty listed is statistical (from the limited size of data control samples and Monte Carlo samples), and the second is systematic. Reprinted from [11].

| $1200 \leq H_T < 1500$ GeV, $N_{\text{jet}} < 7$ | | | | | | |
|--|------------------------------|---------------------------------|---------------------------------|--------------------------|---------------------------------|-------------|
| $N_{\text{jet}}, N_{\text{b-tag}}$ | $M_{\text{T2}} [\text{GeV}]$ | Lost lepton | $Z \rightarrow vv$ | Multijet | Total background | Data |
| 2-3j, 0b | 200-400 | $315 \pm 15 \pm 21$ | $656^{+51}_{-47} \pm 73$ | $39 \pm 16 \pm 12$ | $1009^{+55}_{-52} \pm 85$ | 1128 |
| | 400-600 | $43.0^{+5.2}_{-4.7} \pm 4.9$ | $185^{+14}_{-13} \pm 30$ | $0.03 \pm 0.02 \pm 0.01$ | $228^{+15}_{-14} \pm 31$ | 207 |
| | 600-800 | $14.1^{+2.1}_{-2.0} \pm 1.7$ | $64 \pm 5 \pm 17$ | < 0.01 | $78 \pm 5 \pm 17$ | 83 |
| | 800-1000 | $6.4^{+1.1}_{-1.0} \pm 1.3$ | $32.5^{+2.5}_{-2.3} \pm 7.6$ | < 0.01 | $38.9^{+2.7}_{-2.5} \pm 7.8$ | 36 |
| | 1000-1200 | $3.23^{+0.61}_{-0.59} \pm 0.99$ | $17.5 \pm 1.3 \pm 5.2$ | < 0.01 | $20.7^{+1.5}_{-1.4} \pm 5.3$ | 19 |
| | ≥ 1200 | $0.87^{+0.14}_{-0.13} \pm 0.35$ | $6.0^{+0.5}_{-0.4} \pm 2.6$ | < 0.01 | $6.9 \pm 0.5 \pm 2.6$ | 4 |
| 2-3j, 1b | 200-400 | $61.5^{+7.2}_{-6.5} \pm 4.2$ | $78^{+19}_{-16} \pm 10$ | $9.7 \pm 0.7 \pm 3.0$ | $149^{+21}_{-17} \pm 12$ | 157 |
| | 400-600 | $10.1 \pm 1.4 \pm 1.0$ | $21.9^{+5.4}_{-4.4} \pm 3.8$ | $0.03 \pm 0.02 \pm 0.01$ | $32.0^{+5.6}_{-4.6} \pm 4.1$ | 27 |
| | 600-800 | $2.36^{+0.36}_{-0.35} \pm 0.41$ | $7.5^{+1.9}_{-1.5} \pm 2.0$ | < 0.01 | $9.8^{+1.9}_{-1.6} \pm 2.1$ | 9 |
| | 800-1000 | $0.78^{+0.16}_{-0.15} \pm 0.19$ | $3.84^{+0.95}_{-0.78} \pm 0.93$ | < 0.01 | $4.62^{+0.97}_{-0.79} \pm 0.96$ | 6 |
| | 1000-1200 | $0.43 \pm 0.08 \pm 0.14$ | $2.13^{+0.53}_{-0.43} \pm 0.64$ | < 0.01 | $2.56^{+0.54}_{-0.44} \pm 0.66$ | 2 |
| | ≥ 1200 | $0.14^{+0.05}_{-0.04} \pm 0.06$ | $0.71^{+0.18}_{-0.14} \pm 0.31$ | < 0.01 | $0.86^{+0.18}_{-0.15} \pm 0.31$ | 0 |
| 2-3j, 2b | 200-400 | $4.8^{+2.0}_{-1.6} \pm 0.3$ | $11^{+11}_{-6} \pm 2$ | $1.38 \pm 0.13 \pm 0.43$ | $18^{+11}_{-6} \pm 2$ | 18 |
| | 400-600 | $0.61^{+0.30}_{-0.25} \pm 0.07$ | $3.2^{+3.1}_{-2.7} \pm 0.7$ | < 0.01 | $3.8^{+3.1}_{-1.8} \pm 0.7$ | 5 |
| | 600-800 | $0.21^{+0.11}_{-0.09} \pm 0.04$ | $1.1^{+1.1}_{-0.6} \pm 0.4$ | < 0.01 | $1.3^{+1.1}_{-0.6} \pm 0.4$ | 2 |
| | 800-1000 | $0.07^{+0.04}_{-0.03} \pm 0.02$ | $0.56^{+0.55}_{-0.31} \pm 0.18$ | < 0.01 | $0.63^{+0.55}_{-0.31} \pm 0.18$ | 1 |
| | ≥ 1000 | $0.03 \pm 0.02 \pm 0.01$ | $0.42^{+0.41}_{-0.23} \pm 0.18$ | < 0.01 | $0.46^{+0.41}_{-0.23} \pm 0.18$ | 1 |
| 2-6j, $\geq 3b$ | 200-400 | $22.6^{+4.7}_{-4.2} \pm 1.8$ | $0.0^{+6.6}_{-0.0} \pm 0.0$ | $4.4 \pm 0.2 \pm 1.5$ | $27.0^{+8.1}_{-4.2} \pm 2.4$ | 25 |
| | 400-600 | $1.58^{+0.51}_{-0.48} \pm 0.34$ | $0.0^{+1.6}_{-0.0} \pm 0.0$ | $0.02 \pm 0.01 \pm 0.01$ | $1.6^{+1.7}_{-0.5} \pm 0.3$ | 3 |
| | ≥ 600 | $0.47^{+0.27}_{-0.26} \pm 0.19$ | $0.00^{+0.94}_{-0.00} \pm 0.00$ | < 0.01 | $0.47^{+0.98}_{-0.26} \pm 0.19$ | 4 |
| 4-6j, 0b | 200-400 | $606^{+21}_{-20} \pm 41$ | $909^{+63}_{-59} \pm 90$ | $208 \pm 12 \pm 64$ | $1720^{+70}_{-60} \pm 130$ | 1768 |
| | 400-600 | $84.3^{+7.4}_{-6.9} \pm 5.8$ | $234^{+16}_{-15} \pm 34$ | $0.88 \pm 0.09 \pm 0.27$ | $319^{+18}_{-17} \pm 36$ | 301 |
| | 600-800 | $21.1^{+3.2}_{-2.9} \pm 2.3$ | $75 \pm 5 \pm 17$ | $0.06 \pm 0.02 \pm 0.02$ | $96 \pm 6 \pm 17$ | 99 |
| | 800-1000 | $7.6^{+1.2}_{-1.1} \pm 1.1$ | $35.2^{+2.4}_{-2.3} \pm 8.0$ | $0.01 \pm 0.01 \pm 0.00$ | $42.7^{+2.7}_{-2.5} \pm 8.2$ | 41 |
| | 1000-1200 | $2.23^{+0.36}_{-0.33} \pm 0.61$ | $14.1^{+1.0}_{-0.9} \pm 4.2$ | < 0.01 | $16.3 \pm 1.0 \pm 4.2$ | 15 |
| | ≥ 1200 | $0.47^{+0.10}_{-0.09} \pm 0.19$ | $3.0 \pm 0.2 \pm 1.3$ | < 0.01 | $3.5 \pm 0.2 \pm 1.3$ | 5 |
| 4-6j, 1b | 200-400 | $278^{+15}_{-14} \pm 20$ | $254^{+33}_{-30} \pm 28$ | $97 \pm 2 \pm 30$ | $629^{+36}_{-33} \pm 50$ | 579 |
| | 400-600 | $30.3^{+4.0}_{-3.7} \pm 2.7$ | $65^{+9}_{-8} \pm 10$ | $0.33 \pm 0.06 \pm 0.10$ | $96^{+9}_{-8} \pm 11$ | 79 |
| | 600-800 | $8.2^{+1.4}_{-1.3} \pm 1.0$ | $21.0^{+2.8}_{-2.5} \pm 4.8$ | $0.02 \pm 0.01 \pm 0.01$ | $29.2^{+3.1}_{-2.8} \pm 5.0$ | 16 |
| | 800-1000 | $2.36^{+0.56}_{-0.54} \pm 0.50$ | $9.8^{+1.3}_{-1.1} \pm 2.3$ | $0.01 \pm 0.01 \pm 0.00$ | $12.2^{+1.4}_{-1.3} \pm 2.4$ | 9 |
| | 1000-1200 | $1.00 \pm 0.24 \pm 0.31$ | $4.0 \pm 0.5 \pm 1.2$ | < 0.01 | $5.0^{+0.6}_{-0.5} \pm 1.2$ | 6 |
| | ≥ 1200 | $0.07 \pm 0.02 \pm 0.03$ | $0.86^{+0.11}_{-0.10} \pm 0.37$ | < 0.01 | $0.92^{+0.11}_{-0.10} \pm 0.37$ | 1 |
| 4-6j, 2b | 200-400 | $120.4^{+9.1}_{-8.7} \pm 9.8$ | $45^{+18}_{-13} \pm 5$ | $26.0 \pm 0.6 \pm 8.1$ | $191^{+20}_{-16} \pm 15$ | 194 |
| | 400-600 | $11.9 \pm 1.4 \pm 1.5$ | $11.5^{+4.6}_{-3.4} \pm 1.8$ | $0.11 \pm 0.03 \pm 0.04$ | $23.4^{+4.8}_{-3.7} \pm 2.6$ | 27 |
| | 600-800 | $3.49 \pm 0.83 \pm 0.75$ | $3.7^{+1.5}_{-1.1} \pm 1.0$ | < 0.01 | $7.2^{+1.7}_{-1.4} \pm 1.3$ | 7 |
| | 800-1000 | $0.66 \pm 0.16 \pm 0.20$ | $1.73^{+0.69}_{-0.51} \pm 0.48$ | < 0.01 | $2.38^{+0.71}_{-0.54} \pm 0.53$ | 3 |
| | ≥ 1000 | $0.15 \pm 0.04 \pm 0.06$ | $0.84^{+0.34}_{-0.25} \pm 0.36$ | < 0.01 | $1.00^{+0.34}_{-0.25} \pm 0.36$ | 0 |

Table A.7: Predictions and observations for the 31 search regions with $1200 \leq H_T < 1500$ GeV, $N_{\text{jet}} \geq 7$. For each of the background predictions, the first uncertainty listed is statistical (from the limited size of data control samples and Monte Carlo samples), and the second is systematic. Reprinted from [11].

| $1200 \leq H_T < 1500$ GeV, $N_{\text{jet}} \geq 7$ | | | | | | |
|---|----------------|---------------------------------|---------------------------------|--------------------------|--|------------|
| $N_{\text{jet}}, N_{\text{b-tag}}$ | M_{T2} [GeV] | Lost lepton | $Z \rightarrow vv$ | Multijet | Total background | Data |
| 7-9j, 0b | 200-400 | $120.4^{+9.8}_{-9.2} \pm 9.0$ | $108^{+26}_{-21} \pm 21$ | $91 \pm 3 \pm 29$ | $\mathbf{319}^{+28}_{-24} \pm 38$ | 379 |
| | 400-600 | $16.5^{+1.9}_{-1.8} \pm 2.0$ | $25.8^{+6.3}_{-5.1} \pm 5.7$ | $0.80 \pm 0.09 \pm 0.25$ | $\mathbf{43.1}^{+6.5}_{-5.4} \pm 6.3$ | 45 |
| | 600-800 | $2.94 \pm 0.42 \pm 0.63$ | $8.6^{+2.1}_{-1.7} \pm 2.1$ | $0.06 \pm 0.02 \pm 0.02$ | $\mathbf{11.6}^{+2.1}_{-1.8} \pm 2.2$ | 17 |
| | 800-1000 | $0.77^{+0.14}_{-0.13} \pm 0.24$ | $2.90^{+0.70}_{-0.58} \pm 1.00$ | $0.01 \pm 0.01 \pm 0.00$ | $\mathbf{3.7}^{+0.7}_{-0.6} \pm 1.0$ | 3 |
| | ≥ 1000 | $0.11 \pm 0.03 \pm 0.05$ | $1.09^{+0.26}_{-0.22} \pm 0.50$ | < 0.01 | $\mathbf{1.21}^{+0.27}_{-0.22} \pm 0.50$ | 0 |
| 7-9j, 1b | 200-400 | $133.8^{+8.0}_{-7.7} \pm 9.8$ | $36^{+13}_{-10} \pm 8$ | $58 \pm 2 \pm 18$ | $\mathbf{228}^{+15}_{-13} \pm 23$ | 247 |
| | 400-600 | $16.6^{+2.9}_{-2.7} \pm 1.3$ | $8.7^{+3.2}_{-2.4} \pm 2.1$ | $0.46 \pm 0.07 \pm 0.14$ | $\mathbf{25.8}^{+4.3}_{-3.6} \pm 2.7$ | 23 |
| | 600-800 | $1.83^{+0.43}_{-0.41} \pm 0.28$ | $2.9^{+1.1}_{-0.8} \pm 0.8$ | $0.03 \pm 0.02 \pm 0.01$ | $\mathbf{4.8}^{+1.1}_{-0.9} \pm 0.8$ | 7 |
| | 800-1000 | $0.65^{+0.24}_{-0.23} \pm 0.18$ | $0.95^{+0.34}_{-0.26} \pm 0.34$ | $0.02 \pm 0.01 \pm 0.01$ | $\mathbf{1.62}^{+0.42}_{-0.35} \pm 0.39$ | 2 |
| | ≥ 1000 | $0.22 \pm 0.19 \pm 0.09$ | $0.36^{+0.13}_{-0.10} \pm 0.17$ | < 0.01 | $\mathbf{0.58}^{+0.23}_{-0.21} \pm 0.19$ | 0 |
| 7-9j, 2b | 200-400 | $124.0^{+7.6}_{-7.4} \pm 9.1$ | $9.9^{+3.6}_{-2.7} \pm 2.5$ | $21.4 \pm 0.5 \pm 6.9$ | $\mathbf{155} \pm 8 \pm 12$ | 162 |
| | 400-600 | $15.0^{+2.8}_{-2.6} \pm 1.3$ | $2.41^{+0.87}_{-0.66} \pm 0.67$ | $0.12 \pm 0.03 \pm 0.04$ | $\mathbf{17.5}^{+3.0}_{-2.7} \pm 1.5$ | 18 |
| | 600-800 | $2.47^{+0.78}_{-0.76} \pm 0.53$ | $0.81^{+0.29}_{-0.22} \pm 0.26$ | $0.01 \pm 0.01 \pm 0.00$ | $\mathbf{3.29}^{+0.83}_{-0.79} \pm 0.60$ | 1 |
| | ≥ 800 | $0.24 \pm 0.11 \pm 0.10$ | $0.36^{+0.13}_{-0.10} \pm 0.16$ | < 0.01 | $\mathbf{0.60}^{+0.17}_{-0.15} \pm 0.19$ | 1 |
| 7-9j, 3b | 200-400 | $30.0 \pm 2.6 \pm 3.2$ | $1.89^{+0.68}_{-0.52} \pm 0.64$ | $5.0 \pm 0.3 \pm 1.8$ | $\mathbf{36.9}^{+2.7}_{-2.6} \pm 3.8$ | 46 |
| | 400-600 | $4.1^{+1.1}_{-1.0} \pm 0.6$ | $0.45^{+0.16}_{-0.12} \pm 0.18$ | $0.02 \pm 0.01 \pm 0.01$ | $\mathbf{4.6}^{+1.1}_{-1.0} \pm 0.6$ | 2 |
| | ≥ 600 | $0.92^{+0.50}_{-0.49} \pm 0.38$ | $0.23^{+0.08}_{-0.06} \pm 0.11$ | < 0.01 | $\mathbf{1.15} \pm 0.50 \pm 0.40$ | 1 |
| 7-9j, $\geq 4b$ | 200-400 | $9.1 \pm 1.6 \pm 1.8$ | $0.26^{+0.10}_{-0.07} \pm 0.23$ | $0.88 \pm 0.10 \pm 0.32$ | $\mathbf{10.3} \pm 1.6 \pm 1.9$ | 9 |
| | ≥ 400 | $0.44^{+0.24}_{-0.23} \pm 0.08$ | $0.10^{+0.04}_{-0.03} \pm 0.09$ | < 0.01 | $\mathbf{0.53} \pm 0.24 \pm 0.12$ | 0 |
| $\geq 10j$, 0b | 200-400 | $7.7^{+1.2}_{-1.1} \pm 0.8$ | $2.7^{+0.6}_{-0.5} \pm 2.8$ | $8.3 \pm 0.9 \pm 3.0$ | $\mathbf{18.7}^{+1.6}_{-1.5} \pm 4.1$ | 17 |
| | 400-600 | $1.00 \pm 0.32 \pm 0.22$ | $0.56^{+0.13}_{-0.11} \pm 0.62$ | $0.11 \pm 0.03 \pm 0.04$ | $\mathbf{1.66}^{+0.35}_{-0.34} \pm 0.66$ | 1 |
| | ≥ 600 | $0.10^{+0.35}_{-0.04} \pm 0.04$ | $0.14^{+0.08}_{-0.03} \pm 0.14$ | $0.01 \pm 0.01 \pm 0.00$ | $\mathbf{0.24}^{+0.36}_{-0.05} \pm 0.15$ | 0 |
| $\geq 10j$, 1b | 200-400 | $15.2 \pm 1.8 \pm 1.4$ | $1.1^{+0.4}_{-0.3} \pm 1.2$ | $5.3 \pm 0.2 \pm 1.9$ | $\mathbf{21.6}^{+1.9}_{-1.8} \pm 2.7$ | 22 |
| | 400-600 | $1.27^{+0.38}_{-0.36} \pm 0.11$ | $0.22^{+0.08}_{-0.06} \pm 0.26$ | $0.05 \pm 0.02 \pm 0.02$ | $\mathbf{1.55}^{+0.39}_{-0.37} \pm 0.29$ | 6 |
| | ≥ 600 | $0.03 \pm 0.02 \pm 0.01$ | $0.05^{+0.10}_{-0.01} \pm 0.05$ | < 0.01 | $\mathbf{0.07}^{+0.11}_{-0.02} \pm 0.05$ | 0 |
| $\geq 10j$, 2b | 200-400 | $16.9 \pm 1.8 \pm 1.5$ | $0.44^{+0.16}_{-0.12} \pm 0.50$ | $2.7 \pm 0.2 \pm 1.0$ | $\mathbf{20.1} \pm 1.8 \pm 1.9$ | 16 |
| | 400-600 | $2.62^{+0.71}_{-0.68} \pm 0.30$ | $0.09 \pm 0.03 \pm 0.11$ | $0.01 \pm 0.01 \pm 0.00$ | $\mathbf{2.73}^{+0.71}_{-0.68} \pm 0.32$ | 2 |
| | ≥ 600 | $0.23 \pm 0.15 \pm 0.10$ | $0.02^{+0.08}_{-0.01} \pm 0.02$ | < 0.01 | $\mathbf{0.25}^{+0.17}_{-0.15} \pm 0.10$ | 0 |
| $\geq 10j$, 3b | 200-400 | $5.58^{+0.86}_{-0.85} \pm 0.61$ | $0.12^{+0.11}_{-0.03} \pm 0.16$ | $1.04 \pm 0.10 \pm 0.42$ | $\mathbf{6.74}^{+0.87}_{-0.86} \pm 0.76$ | 6 |
| | ≥ 400 | $0.51 \pm 0.22 \pm 0.06$ | $0.03^{+0.11}_{-0.01} \pm 0.04$ | < 0.01 | $\mathbf{0.54}^{+0.25}_{-0.22} \pm 0.08$ | 0 |
| $\geq 10j, \geq 4b$ | ≥ 200 | $2.59 \pm 0.82 \pm 0.62$ | $0.10^{+0.13}_{-0.03} \pm 0.13$ | $0.31 \pm 0.06 \pm 0.13$ | $\mathbf{3.00}^{+0.83}_{-0.82} \pm 0.65$ | 7 |

Table A.8: Predictions and observations for the 30 search regions with $H_T \geq 1500$ GeV, $N_{\text{jet}} < 7$. For each of the background predictions, the first uncertainty listed is statistical (from the limited size of data control samples and Monte Carlo samples), and the second is systematic. Reprinted from [11].

| $H_T \geq 1500$ GeV, $N_{\text{jet}} < 7$ | | | | | | |
|---|------------------------------|---------------------------------|---------------------------------|--------------------------|---------------------------------|------------|
| $N_{\text{jet}}, N_{\text{b-tag}}$ | $M_{\text{T2}} [\text{GeV}]$ | Lost lepton | $Z \rightarrow vv$ | Multijet | Total background | Data |
| 2-3j, 0b | 400-600 | $27.2^{+4.4}_{-3.9} \pm 2.5$ | $150^{+14}_{-13} \pm 19$ | $0.16 \pm 0.04 \pm 0.05$ | $177^{+15}_{-13} \pm 20$ | 125 |
| | 600-800 | $7.8^{+1.4}_{-1.2} \pm 0.8$ | $38.7^{+3.6}_{-3.3} \pm 8.4$ | < 0.01 | $46.5^{+3.9}_{-3.6} \pm 8.6$ | 37 |
| | 800-1000 | $2.29^{+0.39}_{-0.34} \pm 0.35$ | $17.2^{+1.6}_{-1.5} \pm 3.4$ | < 0.01 | $19.5^{+1.7}_{-1.5} \pm 3.4$ | 19 |
| | 1000-1200 | $1.20^{+0.21}_{-0.19} \pm 0.26$ | $9.0 \pm 0.8 \pm 1.8$ | < 0.01 | $10.2^{+0.9}_{-0.8} \pm 1.9$ | 14 |
| | 1200-1400 | $0.80^{+0.16}_{-0.14} \pm 0.22$ | $4.9^{+0.5}_{-0.4} \pm 1.3$ | < 0.01 | $5.7^{+0.5}_{-0.4} \pm 1.4$ | 4 |
| | 1400-1800 | $0.43^{+0.09}_{-0.08} \pm 0.15$ | $2.80^{+0.26}_{-0.24} \pm 0.98$ | < 0.01 | $3.23^{+0.28}_{-0.26} \pm 0.99$ | 3 |
| | ≥ 1800 | $0.05 \pm 0.02 \pm 0.02$ | $0.41^{+0.04}_{-0.03} \pm 0.19$ | < 0.01 | $0.46 \pm 0.04 \pm 0.19$ | 0 |
| 2-3j, 1b | 400-600 | $5.2^{+1.1}_{-1.0} \pm 0.6$ | $13.4^{+4.9}_{-3.7} \pm 1.9$ | $0.09 \pm 0.03 \pm 0.03$ | $18.7^{+5.0}_{-3.8} \pm 2.1$ | 23 |
| | 600-800 | $1.52^{+0.43}_{-0.41} \pm 0.27$ | $3.5^{+1.3}_{-1.0} \pm 1.0$ | < 0.01 | $5.0^{+1.3}_{-1.0} \pm 1.0$ | 3 |
| | 800-1000 | $0.38 \pm 0.09 \pm 0.10$ | $1.53^{+0.55}_{-0.42} \pm 0.35$ | < 0.01 | $1.90^{+0.56}_{-0.43} \pm 0.37$ | 3 |
| | 1000-1200 | $0.10 \pm 0.03 \pm 0.03$ | $0.81^{+0.29}_{-0.22} \pm 0.24$ | < 0.01 | $0.91^{+0.29}_{-0.22} \pm 0.24$ | 4 |
| | ≥ 1200 | $0.19 \pm 0.06 \pm 0.08$ | $0.73^{+0.26}_{-0.20} \pm 0.31$ | < 0.01 | $0.92^{+0.27}_{-0.21} \pm 0.32$ | 0 |
| 2-3j, 2b | ≥ 400 | $0.63^{+0.49}_{-0.36} \pm 0.26$ | $0.0^{+3.0}_{-0.0} \pm 0.0$ | < 0.01 | $0.6^{+3.0}_{-0.4} \pm 0.3$ | 2 |
| 2-6j, $\geq 3b$ | 400-600 | $1.72^{+0.73}_{-0.68} \pm 0.42$ | $1.1^{+2.4}_{-0.9} \pm 0.3$ | $0.03 \pm 0.02 \pm 0.01$ | $2.8^{+2.5}_{-1.1} \pm 0.6$ | 1 |
| | ≥ 600 | $0.37^{+0.19}_{-0.18} \pm 0.16$ | $0.5^{+1.2}_{-0.4} \pm 0.2$ | < 0.01 | $0.9^{+1.2}_{-0.5} \pm 0.2$ | 0 |
| 4-6j, 0b | 400-600 | $46.4^{+5.6}_{-5.1} \pm 3.6$ | $176^{+15}_{-14} \pm 23$ | $1.62 \pm 0.13 \pm 0.46$ | $224^{+16}_{-15} \pm 24$ | 207 |
| | 600-800 | $10.6^{+2.3}_{-1.9} \pm 1.2$ | $45.5^{+4.0}_{-3.7} \pm 9.9$ | $0.07 \pm 0.03 \pm 0.02$ | $56^{+5}_{-4} \pm 10$ | 62 |
| | 800-1000 | $4.5^{+1.1}_{-1.0} \pm 0.5$ | $20.3^{+1.8}_{-1.6} \pm 3.9$ | < 0.01 | $24.8^{+2.1}_{-1.9} \pm 4.1$ | 31 |
| | 1000-1200 | $1.35^{+0.30}_{-0.26} \pm 0.24$ | $10.6 \pm 0.9 \pm 2.1$ | < 0.01 | $11.9^{+1.0}_{-0.9} \pm 2.2$ | 12 |
| | 1200-1400 | $0.89^{+0.27}_{-0.25} \pm 0.23$ | $5.7 \pm 0.5 \pm 1.5$ | < 0.01 | $6.6^{+0.6}_{-0.5} \pm 1.6$ | 9 |
| | 1400-1600 | $0.20 \pm 0.05 \pm 0.07$ | $2.64^{+0.23}_{-0.21} \pm 0.92$ | < 0.01 | $2.84^{+0.24}_{-0.22} \pm 0.92$ | 3 |
| | ≥ 1600 | $0.09 \pm 0.03 \pm 0.04$ | $1.18 \pm 0.10 \pm 0.51$ | < 0.01 | $1.27^{+0.11}_{-0.10} \pm 0.51$ | 2 |
| 4-6j, 1b | 400-600 | $21.0^{+3.7}_{-3.3} \pm 2.0$ | $32.6^{+7.0}_{-5.8} \pm 5.5$ | $0.81 \pm 0.09 \pm 0.23$ | $54.5^{+7.9}_{-6.7} \pm 6.3$ | 72 |
| | 600-800 | $4.79^{+0.91}_{-0.83} \pm 0.62$ | $8.4^{+1.8}_{-1.5} \pm 2.3$ | $0.02 \pm 0.01 \pm 0.01$ | $13.2^{+2.0}_{-1.7} \pm 2.5$ | 20 |
| | 800-1000 | $1.27^{+0.26}_{-0.24} \pm 0.27$ | $3.71^{+0.79}_{-0.66} \pm 0.92$ | $0.03 \pm 0.02 \pm 0.01$ | $5.01^{+0.84}_{-0.71} \pm 0.97$ | 8 |
| | 1000-1400 | $0.89^{+0.21}_{-0.20} \pm 0.28$ | $3.00^{+0.64}_{-0.54} \pm 0.93$ | < 0.01 | $3.89^{+0.68}_{-0.57} \pm 0.98$ | 6 |
| | ≥ 1400 | $0.40^{+0.34}_{-0.33} \pm 0.16$ | $0.72^{+0.15}_{-0.13} \pm 0.31$ | < 0.01 | $1.12^{+0.37}_{-0.36} \pm 0.36$ | 3 |
| 4-6j, 2b | 400-600 | $7.2^{+1.2}_{-1.1} \pm 1.1$ | $4.3^{+2.9}_{-1.9} \pm 1.4$ | $0.17 \pm 0.04 \pm 0.05$ | $11.7^{+3.2}_{-2.2} \pm 1.9$ | 11 |
| | 600-800 | $1.66^{+0.41}_{-0.40} \pm 0.46$ | $1.12^{+0.76}_{-0.48} \pm 0.55$ | $0.01 \pm 0.01 \pm 0.00$ | $2.79^{+0.86}_{-0.63} \pm 0.73$ | 3 |
| | ≥ 800 | $0.32 \pm 0.13 \pm 0.13$ | $0.99^{+0.67}_{-0.43} \pm 0.52$ | < 0.01 | $1.31^{+0.68}_{-0.45} \pm 0.54$ | 4 |

Table A.9: Predictions and observations for the 21 search regions with $H_T \geq 1500$ GeV, $N_{\text{jet}} \geq 7$. For each of the background predictions, the first uncertainty listed is statistical (from the limited size of data control samples and Monte Carlo samples), and the second is systematic. Reprinted from [11].

| $H_T \geq 1500$ GeV, $N_{\text{jet}} \geq 7$ | | | | | | |
|--|----------------|---------------------------------|---------------------------------|--------------------------|--|-----------|
| $N_{\text{jet}}, N_{\text{b-tag}}$ | M_{T2} [GeV] | Lost lepton | $Z \rightarrow vv$ | Multijet | Total background | Data |
| 7-9j, 0b | 400-600 | $14.3^{+1.8}_{-1.7} \pm 1.7$ | $32.3^{+7.5}_{-6.2} \pm 4.3$ | $1.50 \pm 0.13 \pm 0.44$ | $\mathbf{48.1}^{+7.7}_{-6.4} \pm 5.0$ | 36 |
| | 600-800 | $3.77^{+0.56}_{-0.55} \pm 0.69$ | $8.3^{+1.9}_{-1.6} \pm 2.2$ | $0.18 \pm 0.04 \pm 0.05$ | $\mathbf{12.3}^{+2.0}_{-1.7} \pm 2.3$ | 9 |
| | 800-1000 | $1.16^{+0.18}_{-0.17} \pm 0.30$ | $3.70^{+0.86}_{-0.71} \pm 0.83$ | $0.01 \pm 0.01 \pm 0.00$ | $\mathbf{4.86}^{+0.88}_{-0.73} \pm 0.90$ | 6 |
| | 1000-1400 | $0.58 \pm 0.11 \pm 0.19$ | $2.96^{+0.69}_{-0.57} \pm 0.86$ | $0.01 \pm 0.01 \pm 0.00$ | $\mathbf{3.55}^{+0.69}_{-0.58} \pm 0.89$ | 4 |
| | ≥ 1400 | $0.05 \pm 0.01 \pm 0.02$ | $0.71^{+0.17}_{-0.14} \pm 0.30$ | < 0.01 | $\mathbf{0.76}^{+0.17}_{-0.14} \pm 0.30$ | 2 |
| 7-9j, 1b | 400-600 | $12.8^{+2.5}_{-2.3} \pm 1.6$ | $9.2^{+4.2}_{-3.0} \pm 1.4$ | $0.82 \pm 0.09 \pm 0.24$ | $\mathbf{22.9}^{+4.9}_{-3.8} \pm 2.3$ | 25 |
| | 600-800 | $3.49^{+0.94}_{-0.89} \pm 0.76$ | $2.4^{+1.1}_{-0.8} \pm 1.0$ | $0.06 \pm 0.02 \pm 0.02$ | $\mathbf{5.9}^{+1.4}_{-1.2} \pm 1.2$ | 7 |
| | ≥ 800 | $1.09^{+0.34}_{-0.32} \pm 0.45$ | $2.10^{+0.96}_{-0.69} \pm 0.93$ | < 0.01 | $\mathbf{3.2}^{+1.0}_{-0.8} \pm 1.0$ | 2 |
| 7-9j, 2b | 400-600 | $8.1^{+1.8}_{-1.6} \pm 1.0$ | $2.4^{+1.1}_{-0.8} \pm 0.4$ | $0.35 \pm 0.06 \pm 0.10$ | $\mathbf{10.9}^{+2.1}_{-1.8} \pm 1.2$ | 10 |
| | 600-800 | $1.78^{+0.54}_{-0.52} \pm 0.40$ | $0.62^{+0.28}_{-0.20} \pm 0.25$ | $0.02 \pm 0.01 \pm 0.01$ | $\mathbf{2.41}^{+0.61}_{-0.56} \pm 0.49$ | 5 |
| | ≥ 800 | $0.40^{+0.19}_{-0.18} \pm 0.17$ | $0.55^{+0.25}_{-0.18} \pm 0.25$ | $0.01 \pm 0.01 \pm 0.00$ | $\mathbf{0.96}^{+0.31}_{-0.26} \pm 0.30$ | 0 |
| 7-9j, 3b | 400-800 | $2.40^{+0.74}_{-0.72} \pm 0.29$ | $0.32^{+0.15}_{-0.10} \pm 0.12$ | $0.10 \pm 0.03 \pm 0.03$ | $\mathbf{2.82}^{+0.76}_{-0.72} \pm 0.32$ | 2 |
| | ≥ 800 | $0.16 \pm 0.09 \pm 0.07$ | $0.08^{+0.04}_{-0.03} \pm 0.04$ | < 0.01 | $\mathbf{0.24} \pm 0.09 \pm 0.08$ | 0 |
| 7-9j, $\geq 4b$ | ≥ 400 | $0.52^{+0.23}_{-0.22} \pm 0.08$ | $0.07^{+0.03}_{-0.02} \pm 0.06$ | $0.02 \pm 0.01 \pm 0.01$ | $\mathbf{0.61}^{+0.23}_{-0.22} \pm 0.10$ | 1 |
| $\geq 10j$, 0b | 400-800 | $1.41 \pm 0.38 \pm 0.33$ | $1.52^{+0.35}_{-0.29} \pm 0.34$ | $0.23 \pm 0.05 \pm 0.08$ | $\mathbf{3.17}^{+0.52}_{-0.48} \pm 0.49$ | 11 |
| | ≥ 800 | $0.05 \pm 0.02 \pm 0.02$ | $0.37^{+0.09}_{-0.07} \pm 0.17$ | $0.01 \pm 0.01 \pm 0.00$ | $\mathbf{0.43}^{+0.09}_{-0.08} \pm 0.17$ | 0 |
| $\geq 10j$, 1b | 400-800 | $2.16^{+0.71}_{-0.69} \pm 0.25$ | $0.56^{+0.25}_{-0.18} \pm 0.16$ | $0.14 \pm 0.04 \pm 0.05$ | $\mathbf{2.85}^{+0.76}_{-0.71} \pm 0.31$ | 3 |
| | ≥ 800 | $0.55 \pm 0.30 \pm 0.22$ | $0.13^{+0.06}_{-0.04} \pm 0.07$ | < 0.01 | $\mathbf{0.68}^{+0.31}_{-0.30} \pm 0.23$ | 0 |
| $\geq 10j$, 2b | ≥ 400 | $1.98^{+0.69}_{-0.67} \pm 0.24$ | $0.30^{+0.14}_{-0.10} \pm 0.12$ | $0.05 \pm 0.02 \pm 0.02$ | $\mathbf{2.33}^{+0.70}_{-0.68} \pm 0.28$ | 0 |
| $\geq 10j$, 3b | ≥ 400 | $0.77 \pm 0.35 \pm 0.09$ | $0.00^{+0.45}_{-0.00} \pm 0.00$ | $0.05 \pm 0.03 \pm 0.02$ | $\mathbf{0.82}^{+0.57}_{-0.35} \pm 0.09$ | 1 |
| $\geq 10j$, $\geq 4b$ | ≥ 400 | $0.09 \pm 0.05 \pm 0.01$ | $0.00^{+0.45}_{-0.00} \pm 0.00$ | < 0.01 | $\mathbf{0.09}^{+0.45}_{-0.05} \pm 0.01$ | 0 |

Bibliography

- [1]
- [2] Cms luminosity public results. <https://twiki.cern.ch/twiki/bin/view/CMSPublic/LumiPublicResults>. Accessed: 2019-11-08.
- [3] Morad Aaboud et al. Search for long-lived charginos based on a disappearing-track signature in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *JHEP*, 06:022, 2018.
- [4] Georges Aad et al. Search for charginos nearly mass degenerate with the lightest neutralino based on a disappearing-track signature in pp collisions at $\sqrt(s)=8$ TeV with the ATLAS detector. *Phys. Rev.*, D88(11):112006, 2013.
- [5] W Adam, R Frhwirth, A Strandlie, and T Todorov. Reconstruction of electrons with the gaussian-sum filter in the cms tracker at the lhc. *Journal of Physics G: Nuclear and Particle Physics*, 31(9):N9N20, Jul 2005.
- [6] Alan J Barr, Christopher G Lester, M. Andy Parker, Ben C Allanach, and Peter Richardson. Discovering anomaly-mediated supersymmetry at the lhc. *Journal of High Energy Physics*, 2003(03):045045, Mar 2003.
- [7] G L Bayatian and etal. *CMS Physics: Technical Design Report Volume 1: Detector Performance and Software*. Technical Design Report CMS. CERN, Geneva, 2006. There is an error on cover due to a technical problem for some items.
- [8] Michael Benedikt, Paul Collier, V Mertens, John Poole, and Karlheinz Schindl. *LHC Design Report*. CERN Yellow Reports: Monographs. CERN, Geneva, 2004.
- [9] Christoph Borschensky, Michael Krmer, Anna Kulesza, Michelangelo Mangano, Sanjay Padhi, Tilman Plehn, and Xavier Portell. Squark and gluino production cross sections in

pp

p p collisions at

$$\sqrt{s} = 13, 14, 33$$

$s = 13, 14, 33$ and

100

100 tev. *The European Physical Journal C*, 74(12), Dec 2014.

- [10] Sergio Cittolin, Attila Rcz, and Paris Sphicas. *CMS The TriDAS Project: Technical Design Report, Volume 2: Data Acquisition and High-Level Trigger. CMS trigger and data-acquisition project*. Technical Design Report CMS. CERN, Geneva, 2002.
- [11] CMS Collaboration. Searches for physics beyond the standard model with the m_{T2} variable in hadronic final states with and without disappearing tracks in proton-proton collisions at $\sqrt{s} = 13$ tev, 2019.
- [12] The CMS Collaboration. CMS physics technical design report, volume II: Physics performance. *Journal of Physics G: Nuclear and Particle Physics*, 34(6):995–1579, apr 2007.
- [13] The CMS Collaboration. Description and performance of track and primary-vertex reconstruction with the cms tracker. *Journal of Instrumentation*, 9(10):P10009P10009, Oct 2014.
- [14] Seth I. Cooper. Phase i upgrade of the cms hadron calorimeter. *Nuclear and Particle Physics Proceedings*, 273-275:1002 – 1007, 2016. 37th International Conference on High Energy Physics (ICHEP).
- [15] Daniele del Re. Timing performance of the CMS ECAL and prospects for the future. *Journal of Physics: Conference Series*, 587:012003, feb 2015.
- [16] R Gandhi, Christopher S Quigg, M H Reno, and I Sarcevic. Neutrino Interactions at Ultra-high Energies. *Phys. Rev. D*, 58(hep-ph/9807264. FERMILAB-PUB-98-087-T. 9):93009. 32 p, Jul 1998.
- [17] Gian F Giudice, Riccardo Rattazzi, Markus A Luty, and Hitoshi Murayama. Gaugino mass without singlets. *Journal of High Energy Physics*, 1998(12):027027, Dec 1998.
- [18] V. Khachatryan et al. Search for disappearing tracks in proton-proton collisions at $\sqrt{s} = 8$ TeV. *JHEP*, 01:096, 2015.
- [19] V. Khachatryan, A. M. Sirunyan, A. Tumasyan, W. Adam, E. Asilar, T. Bergauer, J. Brandstetter, E. Brondolin, M. Dragicevic, and et al. Search for new physics with the m_{t2} variable in all-jets final states produced in pp collisions at $s = 13$

$\sqrt{s} = 13$

tev. *Journal of High Energy Physics*, 2016(10), Oct 2016.

- [20] V. Khachatryan, A.M. Sirunyan, A. Tumasyan, W. Adam, E. Asilar, T. Bergauer, J. Brandstetter, E. Brondolin, M. Dragicevic, J. Er, and et al. Jet energy scale and resolution in the

cms experiment in pp collisions at 8 tev. *Journal of Instrumentation*, 12(02):P02014P02014, Feb 2017.

- [21] LEP2 SUSY Working Group, ALEPH, DELPHI, L3 and OPAL experiments. Combined LEP chargino results, up to 208 GeV for low Δm , 2002.
- [22] C.G. Lester and D.J. Summers. Measuring masses of semiinvisibly decaying particles pair produced at hadron colliders. *Phys. Lett. B*, 463:99, 1999.
- [23] Zhen Liu and Brock Tweedie. The fate of long-lived superparticles with hadronic decays after lhc run 1. *Journal of High Energy Physics*, 2015(6), Jun 2015.
- [24] Lisa Randall and Raman Sundrum. Out of this world supersymmetry breaking. *Nuclear Physics B*, 557(1-2):79118, Sep 1999.
- [25] A L Read. Presentation of search results: the cl s technique. *Journal of Physics G: Nuclear and Particle Physics*, 28(10):2693, 2002.
- [26] Anirban Saha. Phase 1 upgrade of the CMS pixel detector. *Journal of Instrumentation*, 12(02):C02033–C02033, feb 2017.
- [27] Albert M Sirunyan et al. Search for new phenomena with the M_{T2} variable in the all-hadronic final state produced in protonproton collisions at $\sqrt{s} = 13$ TeV. *Eur. Phys. J.*, C77(10):710, 2017.
- [28] Albert M Sirunyan et al. Search for disappearing tracks as a signature of new long-lived particles in proton-proton collisions at $\sqrt{s} = 13$ TeV. *JHEP*, 08:016, 2018.
- [29] Albert M Sirunyan et al. Search for natural and split supersymmetry in proton-proton collisions at $\sqrt{s} = 13$ TeV in final states with jets and missing transverse momentum. *JHEP*, 05:025, 2018.
- [30] Albert M Sirunyan et al. Inclusive search for supersymmetry in pp collisions at $\sqrt{s} = 13$ TeV using razor variables and boosted object identification in zero and one lepton final states. *JHEP*, 03:031, 2019.
- [31] Albert M Sirunyan et al. Search for supersymmetry in proton-proton collisions at 13 TeV in final states with jets and missing transverse momentum. 2019.
- [32] A.M. Sirunyan, A. Tumasyan, W. Adam, E. Asilar, T. Bergauer, J. Brandstetter, E. Brondolin, M. Dragicevic, J. Er, M. Flechl, and et al. Particle-flow reconstruction and global event description with the cms detector. *Journal of Instrumentation*, 12(10):P10003P10003, Oct 2017.
- [33] Torbjrn Sjstrand. The lund monte carlo for e+ e- jet physics. *Computer Physics Communications*, 28(3):229 – 254, 1983.

- [34] Torbjørn Sjstrand, Stephen Mrenna, and Peter Skands. PYTHIA 6.4 physics and manual. *Journal of High Energy Physics*, 2006(05):026–026, may 2006.
- [35] Jory Sonneveld. Commissioning and first results from the cms phase 1 upgrade pixel detector, 2018.
- [36] M. Tanabashi, K. Hagiwara, K. Hikasa, K. Nakamura, Y. Sumino, F. Takahashi, J. Tanaka, K. Agashe, G. Aielli, C. Amsler, M. Antonelli, D. M. Asner, H. Baer, Sw. Banerjee, R. M. Barnett, T. Basaglia, C. W. Bauer, J. J. Beatty, V. I. Belousov, J. Beringer, S. Bethke, A. Bettini, H. Bichsel, O. Biebel, K. M. Black, E. Blucher, O. Buchmuller, V. Burkert, M. A. Bychkov, R. N. Cahn, M. Carena, A. Ceccucci, A. Cerri, D. Chakraborty, M.-C. Chen, R. S. Chivukula, G. Cowan, O. Dahl, G. D’Ambrosio, T. Damour, D. de Florian, A. de Gouvêa, T. DeGrand, P. de Jong, G. Dissertori, B. A. Dobrescu, M. D’Onofrio, M. Doser, M. Drees, H. K. Dreiner, D. A. Dwyer, P. Eerola, S. Eidelman, J. Ellis, J. Erler, V. V. Ezhela, W. Fettscher, B. D. Fields, R. Firestone, B. Foster, A. Freitas, H. Gallagher, L. Garren, H.-J. Gerber, G. Gerbier, T. Gershon, Y. Gershtein, T. Gherghetta, A. A. Godizov, M. Goodman, C. Grab, A. V. Gritsan, C. Grojean, D. E. Groom, M. Grunewald, A. Gurtu, T. Gutsche, H. E. Haber, C. Hanhart, S. Hashimoto, Y. Hayato, K. G. Hayes, A. Hebecker, S. Heinemeyer, B. Heltsley, J. J. Hernández-Rey, J. Hisano, A. Höcker, J. Holder, A. Holtkamp, T. Hyodo, K. D. Irwin, K. F. Johnson, M. Kado, M. Karliner, U. F. Katz, S. R. Klein, E. Klemp, R. V. Kowalewski, F. Krauss, M. Kreps, B. Krusche, Yu. V. Kuyanov, Y. Kwon, O. Lahav, J. Laiho, J. Lesgourgues, A. Liddle, Z. Ligeti, C.-J. Lin, C. Lippmann, T. M. Liss, L. Littenberg, K. S. Lugovsky, S. B. Lugovsky, A. Lusiani, Y. Makida, F. Maltoni, T. Mannel, A. V. Manohar, W. J. Marciano, A. D. Martin, A. Masoni, J. Matthews, U.-G. Meißner, D. Milstead, R. E. Mitchell, K. Mönig, P. Molaro, F. Moortgat, M. Moskovic, H. Murayama, M. Narain, P. Nason, S. Navas, M. Neubert, P. Nevski, Y. Nir, K. A. Olive, S. Pagan Griso, J. Parsons, C. Patrignani, J. A. Peacock, M. Pennington, S. T. Petcov, V. A. Petrov, E. Pianori, A. Piepke, A. Pomarol, A. Quadt, J. Rademacker, G. Raffelt, B. N. Ratcliff, P. Richardson, A. Ringwald, S. Roesler, S. Rolli, A. Romanouk, L. J. Rosenberg, J. L. Rosner, G. Rybka, R. A. Ryutin, C. T. Sachrajda, Y. Sakai, G. P. Salam, S. Sarkar, F. Sauli, O. Schneider, K. Scholberg, A. J. Schwartz, D. Scott, V. Sharma, S. R. Sharpe, T. Shutt, M. Silari, T. Sjöstrand, P. Skands, T. Skwarnicki, J. G. Smith, G. F. Smoot, S. Spanier, H. Spieler, C. Spiering, A. Stahl, S. L. Stone, T. Sumiyoshi, M. J. Syphers, K. Terashi, J. Terning, U. Thoma, R. S. Thorne, L. Tiator, M. Titov, N. P. Tkachenko, N. A. Törnqvist, D. R. Tovey, G. Valencia, R. Van de Water, N. Varelas, G. Venanzoni, L. Verde, M. G. Vincter, P. Vogel, A. Vogt, S. P. Wakely, W. Walkowiak, C. W. Walter, D. Wands, D. R. Ward, M. O. Wascko, G. Weiglein, D. H. Weinberg, E. J. Weinberg, M. White, L. R. Wiencke, S. Willocq, C. G. Wohl, J. Womersley, C. L. Woody, R. L. Workman, W.-M. Yao, G. P. Zeller, O. V. Zenin, R.-Y. Zhu, S.-L. Zhu, F. Zimmermann, P. A. Zyla, J. Anderson, L. Fuller, V. S. Lugovsky, and P. Schaffner. Review of particle physics. *Phys. Rev. D*, 98:030001, Aug 2018.