

# Magnetorotational instability: a review

Shi Sim, David P. Larson, and Wonjae Lee

University of California, San Diego

(Received 4 June 2013)

We review magnetorotational instability and its role in transporting angular momentum within accretion disks. Theoretical, experimental and numerical simulation research work is surveyed.

## 1. Introduction

Turbulence generating magnetorotational instability (MRI) was first discovered by [Velikhov \(1959\)](#) and [Chandrasekhar \(1960\)](#), and was later re-discovered by [Balbus & Hawley \(1998\)](#) for astrophysical applications. MRI has since then been confirmed by robust numerical simulations, but to date has not been verified experimentally or through observations. To understand the importance of MRI, we have to look into the accretion disk theory where many astrophysical phenomena take place in.

Accretion disks are disk made up of gas, dust, and plasma that rotates around and gradually collapses onto an object in the center, e.g leading to formation of a star. However, accretion can happen only with an efficient mechanism for rapidly transporting angular momentum outwards. It was suggested that turbulence drives angular momentum outwards but there was no known mechanism that would generate turbulence. These Keplerian disks satisfies the Rayleigh stability criterion (see [Rayleigh 1916](#)) against centrifugal instability. Thus, there has to be other mechanisms that generates turbulence and this is where MRI and non-linear hydrodynamic instabilities comes into the picture. There has been other mechanisms suggested but MRI seem like the most probable explanation at this point.

MRI is not only applicable to astrophysical phenomena but geophysical ones as well, e.g. the Earth's magnetic field. It is used to better understand how planetary magnetic field might have formed on Earth or other planets and how and why the field is sustained or decayed through time.

In this report, we attempt to explain and illustrate the instability in an astrophysical sense and go through what has been done experimentally and numerically in terms of MRI.

## 2. Physical Explanation

### 2.1. Perfectly conducting fluid in magnetic field

The magnetic field lines are tied to the conducting fluid in which they are embedded. We can show this by comparing a transport equation of magnetic field with a equation for a line element moving with fluid. We first combine Ohm's Law in ideal conductor and Faraday's equation to get a transport equation for magnetic field,

$$\begin{aligned}\frac{\partial \mathbf{B}}{\partial t} &= \nabla \times (\mathbf{u} \times \mathbf{B}) \\ &= \mathbf{B} \cdot \nabla \mathbf{u} - \mathbf{u} \cdot \nabla \mathbf{B} - \mathbf{B} \nabla \cdot \mathbf{u}.\end{aligned}\tag{2.1}$$

Assuming incompressibility, we get

$$\frac{D\mathbf{B}}{Dt} = (\mathbf{B} \cdot \nabla)\mathbf{u}, \quad (2.2)$$

where  $\frac{D}{Dt}$  is the convective derivative defined as

$$\frac{D}{Dt} = \frac{\partial}{\partial t} + (\mathbf{u} \cdot \nabla) \quad (2.3)$$

Considering a short line element  $d\mathbf{l}$  moving with fluid, we can express the rate of change of  $d\mathbf{l}$  as

$$\frac{D}{Dt}(d\mathbf{l}) = \mathbf{u}(\mathbf{x} + d\mathbf{l}) - \mathbf{u}(\mathbf{x}) = (d\mathbf{l} \cdot \nabla)\mathbf{u}. \quad (2.4)$$

Comparing two equations above, we can conclude that  $\mathbf{B}$  and  $d\mathbf{l}$  obey the same equation (see [Davidson 2001](#)). Therefore, the field lines are frozen into the fluid.

## 2.2. Magnetic tension

When the fluid elements are displaced from their equilibrium position, the magnetic field lines move together with fluid elements and behave like elastic bands frozen-into the fluid. Using Ampere's law, the Lorenz force ( $\mathbf{J} \times \mathbf{B}$ ) may be written as

$$\mathbf{J} \times \mathbf{B} = -\nabla \left( \frac{B^2}{2\mu} \right) + \frac{(\mathbf{B} \cdot \nabla)\mathbf{B}}{\mu} = -\nabla \left( \frac{B^2}{2\mu} \right) + \frac{\partial}{\partial s} \left[ \frac{B^2}{2\mu} \right] \hat{e}_t - \frac{B^2}{\mu R} \hat{e}_n \quad (2.5)$$

where  $R$  is radius of curvature,  $s$  is a coordinate along a magnetic field line, and  $\hat{e}_t$  and  $\hat{e}_n$  are tangential and normal unit vectors respectively. The last two terms are magnetic tension forces in tangential and normal directions. These forces can also be interpreted as tensile stress of  $B^2/2\mu$  acting on the end of the tube (see [Davidson 2001](#)).

This tensile force which is also known as Faraday or Maxwell tension is analogous to a force acting on a spring. Considering a displacement  $\boldsymbol{\xi} = \mathbf{v}\delta t$ , the Faraday's equation can be written as  $\delta\mathbf{B} = ikB\boldsymbol{\xi}$ . The magnetic tension for small displacement per unit density  $\rho$  is

$$\frac{(\mathbf{B} \cdot \nabla)\delta\mathbf{B}}{\mu\rho} = \frac{ikB\delta\mathbf{B}}{\mu\rho} = -\frac{k^2B^2}{\mu\rho}\boldsymbol{\xi} = -K\boldsymbol{\xi}, \quad (2.6)$$

where  $K$  is comparable to the spring constant (see [Wikipedia 2013](#); [Balbus & Hawley 1998](#)). The equation has same form of Hooke's Law for a spring.

## 2.3. Magnetorotational instability

We present a relatively simple physical explanation of the magnetorotational instability from [Balbus \(2011\)](#). Consider Keplerian motion of conducting fluid orbiting around a central body of mass  $M_c$ . Two adjacent fluid elements  $m_i$  and  $m_o$ , at radial position  $r_i$  and  $r_o$ , are orbiting around gravitational center with angular velocities  $\Omega_i = \sqrt{GM_c/r_i^3}$  and  $\Omega_o = \sqrt{GM_c/r_o^3}$  respectively. Therefore the angular velocity of the inner element is higher than that of the outer elements ( $d\Omega^2/dr < 0$ ), but the angular momentum of the inner element is smaller than that of the outer element ( $dr^4\Omega^2/dr > 0$ ).

If a magnetic field line is connecting the conducting fluid elements, the magnetic field will move together with the two elements. Because of the velocity shear in Keplerian motion, the magnetic field will be stretched and bent. Therefore the magnetic field will exert restoring force on the fluid element making the inner element pulled back and the outer element dragged forward. The inner




FIGURE 1. A schematic diagram from Balbus (2011) showing magnetorotational instability. Two fluid elements  $m_i$  (inner) and  $m_o$  (outer) orbit a mass ( $M_c$ ) with the magnetic tension between the two elements represented by a spring. Over time  $m_i$  loses angular momentum, moving closer to  $M_c$  while  $m_o$  gains angular momentum and moves away from  $M_c$ .

element lose angular momentum, therefore it must fall to an orbit of smaller radius. On the other hand the outer element gains angular momentum and moves to the orbit of larger radius (see Balbus 2011; Wikipedia 2013). This outward angular momentum transport makes the small initial displacement get larger. The magnetic fields are stretched even more and gives positive feedback leading the system unstable. This instability is known as a magnetorotational instability (MRI) and is illustrated in Figure 1.

One can show that the dispersion relation for the incompressible ideal flow rotating with angular velocity  $\Omega_0$  is

$$\omega^4 - [2(k^2 V_A^2) + \kappa^2] \omega^2 + (k^2 V_A^2) \left[ (k^2 V_A^2) + r \frac{d\Omega^2}{dr} \right] = 0, \quad (2.7)$$

where  $\omega$  and  $k$  are angular frequency and wave number of a perturbation  $\xi \propto e^{i(kz + \omega t)}$ ,  $V_A = B/\sqrt{\mu\rho}$  is Alfvén speed for the imposed magnetic field  $B$ , and  $\kappa^2$  is the epicyclic frequency, defined as

$$\kappa^2 = 4\Omega_0^2 + r \frac{d\Omega^2}{dr} \quad (2.8)$$

See Balbus & Hawley (1991, 1998); Balbus (2003) for more details. If  $d\Omega^2/dr < 0$ , then one of  $\omega^2$  has negative root for long wave length modes satisfying

$$k^2 < -\frac{r}{V_A^2} \frac{d\Omega^2}{dr}. \quad (2.9)$$

Therefore, the instability criterion for the magnetorotational instability (MRI) is represented as radially decreasing angular velocity,

$$\frac{d\Omega^2}{dr} < 0 \quad (\text{UNSTABLE}) \quad (2.10)$$

Note that the MRI is different from the conventional hydrodynamic instability known as Couette-Taylor centrifugal instability. The Couette-Taylor centrifugal instability criterion for axisymmetric perturbation is represented as radially decreasing angular momentum (see Charu & de Forcrand-

Millard 2011):

$$\frac{dr^4\Omega^2}{dr} < 0 \quad (\text{UNSTABLE}) \quad (2.11)$$

The Keplerian disk is a good example of flows which are stable to the hydrodynamic instability but unstable to the MRI.

### 3. Theoretical Work

Acheson & Hide (1973) and Knobloch (1992) showed the linear stability analysis of rotating magneto-fluid bounded in coaxial cylinder. For the case of gaseous astrophysical disks in unbounded geometry was shown in Balbus & Hawley (1991). The condition for stability was shown to be radially increasing angular velocity profile. We present a linear stability analysis of MRI for radially bounded case following Acheson (1972), Acheson (1973), Knobloch (1992), and Julien & Knobloch (2010).

#### 3.1. Governing equation

The wave dispersion equation of a cylindrical magneto-fluid can be obtained from the magnetohydrodynamic (MHD) equations. Assuming the fluid is inviscid and perfectly conducting, the ideal MHD equations are

$$\rho \left( \frac{\partial \mathbf{u}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{u} \right) = -\nabla p + \mathbf{J} \times \mathbf{B} \quad (3.1)$$

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{u}) = 0 \quad (3.2)$$

$$\frac{d}{dt} \left( \frac{p}{\rho^\gamma} \right) = 0 \quad (3.3)$$

$$\mathbf{E} + \mathbf{u} \times \mathbf{B} = 0 \quad (3.4)$$

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t} \quad (3.5)$$

$$\nabla \times \mathbf{B} = \mu \mathbf{J}, \quad (3.6)$$

where  $\mathbf{u}$  fluid velocity,  $\rho$  is fluid density,  $p$  is pressure.  $d/dt = \partial/\partial t + \mathbf{u} \cdot \nabla$  is convective derivative and  $\gamma$  is the ratio of specific heats.  $\mathbf{E}$ ,  $\mathbf{B}$  and  $\mathbf{J}$  are electric field, magnetic field and current density respectively (see Freidberg 1987).

To investigate the magnetorotational instability, one can consider homogeneous incompressible fluid rotating with angular velocity  $\Omega(r) = V(r)/r$  in externally imposed magnetic fields  $\mathbf{B}_0 = [0, B_\phi(r), B_z(r)]$ . Combining electromagnetic equations with momentum relation, we get the appropriate MHD equations,

$$\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} = -\frac{1}{\rho} \nabla \left( P + \frac{\mathbf{B}^2}{2\mu} \right) + \frac{1}{\mu\rho} (\mathbf{B} \cdot \nabla) \mathbf{B} \quad (3.7)$$

$$\frac{\partial \mathbf{B}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{B} = (\mathbf{B} \cdot \nabla) \mathbf{u} \quad (3.8)$$

$$\nabla \cdot \mathbf{u} = 0 \quad (3.9)$$

$$\nabla \cdot \mathbf{B} = 0. \quad (3.10)$$

### 3.2. Linear perturbation equation and eigenvalue problem

We can get linearized equations by perturbing the basic state by small amount of  $\mathbf{u}_1$  and  $\mathbf{b}_1$  for velocity and magnetic fields. The linear perturbation is assumed to have the form

$$f = \Re \left[ \hat{f}(r) e^{i(m\phi + kz + \omega t)} \right]. \quad (3.11)$$

According to [Acheson \(1972\)](#), the normal mode equations are

$$\hat{b}_r = \frac{\hat{u}_r}{\omega} \left( kB_z + \frac{mB_\phi}{r} \right) \quad (3.12)$$

$$\hat{b}_\phi = -\frac{(\hat{u}_r B_\phi)'}{i\omega} + \frac{k}{\omega} (\hat{u}_\phi B_z - \hat{u}_z B_\phi) \quad (3.13)$$

$$\hat{b}_z = -\frac{(r\hat{u}_r B_z)'}{ri\omega} - \frac{m(\hat{u}_\phi B_z - \hat{u}_z B_\phi)}{r\omega} \quad (3.14)$$

$$\hat{u}_z = -\frac{(r\hat{u}_r)'}{rik} - \frac{m\hat{u}_\phi}{rk} \quad (3.15)$$

$$\begin{aligned} \left( 1 + \frac{m^2}{r^2 k^2} \right) ri\hat{u}_\phi = & -\frac{m}{rk^2} (r\hat{u}_r)' \\ & - \frac{\hat{u}_r}{\left( V_{Az} + \frac{mV_{A\phi}}{rk} \right)^2 \frac{k^2}{\omega^2} - 1} \left\{ -\frac{2\Omega r}{\omega} + \frac{2kV_{A\phi}}{\omega^2} \left( V_{Az} + \frac{mV_{A\phi}}{rk} \right) \right\} \end{aligned} \quad (3.16)$$

where  $V_{A\phi} = B_\phi(r)/\sqrt{\mu\rho}$  and  $V_{Az} = B_z(r)/\sqrt{\mu\rho}$  are Alfvén speeds for associated external magnetic field components.

Solving the normal mode equation set for radial velocity perturbation  $\hat{u}_r = u$  and considering axisymmetric perturbation ( $m = 0$ ), [Acheson \(1973\)](#) obtained following eigenvalue problem,

$$\begin{aligned} \frac{d}{dr} \left[ (\omega^2 - k^2 V_{Az}^2) \left( \frac{du}{dr} + \frac{u}{r} \right) \right] - k^2 \left[ \omega^2 - k^2 V_{Az}^2 + r \frac{d}{dr} \left( \frac{V_{A\phi}^2}{r^2} - \frac{V^2}{r^2} \right) \right] u \\ = -\frac{4k^2}{r^2} \frac{(kV_{A\phi}V_{Az} + \omega V)^2}{(\omega^2 - k^2 V_{Az}^2)} u. \end{aligned} \quad (3.17)$$

### 3.3. Stability criterion

#### 3.3.1. Standard magnetorotational instability

Consider a standard MRI of radially bounded coaxial fluid cylinder with externally imposed axial magnetic field but without axial current flowing. Therefore, we have  $V_{Az} = \text{constant} \neq 0$  and  $V_{A\phi} = 0$ . Considering boundary condition  $u(r_1) = u(r_2) = 0$ , we multiply the eigenvalue equation by complex conjugate of  $u$  and integrate over radial coordinate,

$$(\omega^2 - k^2 V_{Az}^2)^2 = \frac{k^2}{D} \int_{r_1}^{r_2} \left[ \frac{\omega^2}{r^2} \frac{d}{dr} r^2 V^2 - r^2 k^2 V_{Az}^2 \frac{d}{dr} \left( \frac{V^2}{r^2} \right) \right] |u|^2 dr \quad (3.18)$$

where

$$D \equiv \int_{r_1}^{r_2} \left( r \left| \frac{du}{dr} \right|^2 + \frac{|u|^2}{r} + k^2 r |u|^2 \right) dr > 0 \quad (3.19)$$

According to [Chandrasekhar \(1960\)](#),  $\omega^2$  must be real. We get stable modes with  $\omega^2 > 0$  and

unstable modes with  $\omega^2 < 0$ . If the angular velocity increases radially outward,  $\frac{d}{dr} \left( \frac{V^2}{r^2} \right) > 0$ , the system is stable because  $\omega^2$  is bounded from below by positive number,

$$\omega^2 > \frac{r^2 k^2 V_{Az}^2 \frac{d}{dr} \left( \frac{V^2}{r^2} \right)}{4 \frac{V^2}{r} + r^2 \frac{d}{dr} \left( \frac{V^2}{r^2} \right)} > 0. \quad (3.20)$$

If we have radially decreasing angular velocity profile,  $\frac{d}{dr} \left( \frac{V^2}{r^2} \right) < 0$ , somewhere  $r_1 < r < r_2$ , then  $\omega^2$  may have negative solution which makes the system unstable.

### 3.3.2. Helical magnetorotational instability

When the external nonzero magnetic fields in axial and azimuthal directions are considered, it was found that the eigenvalue equation can be written as

$$\frac{d}{dr} r \frac{du}{dr} - \frac{u}{r} - k^2 r u = \frac{k^2}{(\omega^2 - k^2 V_{Az}^2)^2} \left[ r^2 \frac{d}{dr} \left( \frac{V_{A\phi}^2 - V^2}{r^2} \right) (\omega^2 - k^2 V_{Az}^2) - \frac{4}{r} (k V_{A\phi} V_{Az} - \omega V)^2 \right] u \quad (3.21)$$

According to [Knobloch \(1992\)](#) and [Julien & Knobloch \(2010\)](#), the exponentially growing mode  $\omega = -i\lambda$ ,  $\lambda > 0$  is possible when the eigenvalue relation has following form

$$(\lambda^2 + k^2 V_{Az}^2)^2 = \frac{k^2}{D} \int_{r_1}^{r_2} \left[ r^2 \frac{d}{dr} \left( \frac{V_{A\phi}^2 - V^2}{r^2} \right) (\lambda^2 + k^2 V_{Az}^2) + \frac{4}{r} (k V_{A\phi} V_{Az} - i\lambda V)^2 \right] |u|^2 dr. \quad (3.22)$$

Considering the imaginary part of the equation, we have

$$\int_{r_1}^{r_2} \frac{1}{r} V_{\phi} V |u|^2 dr = 0 \quad (3.23)$$

[Knobloch \(1992\)](#) showed that exponentially growing instability is only possible when  $V_{A\phi}$  or  $V$  changes sign somewhere in  $r_1 < r < r_2$ .

## 4. Laboratory Experiments

Attempts to generate MRI in laboratory environments did not begin until 2001, four decades after the first magnetohydrodynamic (MHD) experiments (see [Donnelley & Ozima 1960](#)). In contrast to MRI, the first MHD experiments focused on stabilizing, rather than generating, instabilities within Taylor-Couette flow using magnetic fields. Experiments aimed at generating MRI were not devised until the after published work indicating MRI's role as the transport mechanism of angular momentum within accretion disks (need to add citation of paper).

### 4.1. Standard MRI

Following [Ji et al. \(2001\)](#), experimental generation of standard MRI typically involves applying an axial magnetic field to a stable, quasi-Keplerian flow.

One outstanding technical challenge in generating standard MRI is achieving the required flow conditions without damaging the experimental setup. (add more here)

Standard MRI experiments must also be careful to prevent Ekman effects (add more here)

Also, researchers must be careful to keep the background flow laminar so as to prevent uncertainty on whether MRI or turbulence was the first instability (add more here)




FIGURE 2. Diagram from [Sisan \*et al.\* \(2004\)](#) showing the experimental setup. The fixed outer steel sphere contains liquid sodium (the conducting fluid), with the magnetic field imposed coaxially to the rotating inner copper sphere. The magnetic fields and fluid velocity are measured using Hall probes and Doppler velocimetry.

[Ji \*et al.\* \(2001\)](#), [Goodman & Ji \(2002\)](#) = first modern MRI experiments

[Sisan \*et al.\* \(2004\)](#) = inner rotating (copper) sphere, outer fixed (steel) sphere, liquid sodium; found  $\beta_c$  above which MRI occurred (NOTE: unique experiment =  $j$ , write more about it)

#### 4.2. Helical MRI

As noted in Section 4.1, a difficulty with generation of MRI is attaining sufficient speeds without damaging the experimental setup. Although researchers are actively working on solving this issue, there are also several groups researching helical MRI (HMRI), a type of MRI which requires less extreme flow conditions.

HMRI is MRI generated by applying axial and azimuthal magnetic fields to a magnetized fluid and was first proposed by [Hollerbach & Rudiger \(2005\)](#). Hollerbach's paper showed that the addition of an azimuthal magnetic field would allowed MRI to be generated at a lower Reynolds number than standard MRI requires.

[Stefani \*et al.\* \(2006, 2007\)](#) = experimentally proved helical MRI

[Stefani \*et al.\* \(2009\)](#) = improve helical MRI experiment

[Stefani \*et al.\* \(2012\)](#) = DRESYN experiment setup

#### 4.3. Azimuthal MRI

It should be noted that there are experiments aimed at azimuthal MRI (azimuthal magnetic fields only)

## 5. Numerical Work

In terms of MRI, numerical simulations on top of experiments mentioned in the previous section are also employed to try to explain the anomalous viscosity and magnetic field generation. Direct numerical simulations (DNS) are typically used with the advance of supercomputers and dramatically decreased computing time. DNS of the magnetohydrodynamics (MHD) Navier Stokes equations have had much success in the determination of the existence of MRI. Many of the numerical simulations assume incompressible flow if Alfven speed slower than sound speed as in that

regime, the MHD equations can be simplified to be incompressible since it is not fundamental to the instability Balbus & Hawley (1991).

A strong magnetic field will suppress MRI as shown by Liu (2008). From linear stability analysis of the exact nonlinear streaming solution done by Goodman & Xu (1994), we know the magnetic field perturbations grows exponentially, indicating that the solution is unstable. 3D numerical simulation will help to confirm this analysis.

With regards to 2D vs 3D simulations, a good comparison was done by Hawley *et al.* (1995) where they introduced the now well-known local shearing box model which incorporated coriolis and tidal forcing but neglects background gradients in pressure and density (not important unless radial oscillations are significant). Periodic boundary conditions were used with an added shearing component to the radial direction. "Method of characteristics-constrained transport" algorithm as explained in Hawley *et al.* (1995) was implemented. This model with an initial input magnetic field with zero- volume average initiates MRI but may obliterate the magnetic field and thus suppress the turbulence.

The comparison of the 3D simulations results from HGB with the 2D results from Balbus & Hawley (1994) shows significant difference. The 2D results have a general channel/streaming flow solution whereas the 3D results may go through the channel flow but eventually evolves further into a turbulent flow with some not going through the channel flow at all. Note: vertical component of gravity, hence buoyancy, was ignored and since their simulations showed significant density contrast, the shearing box model is not self-consistent. Purely hydrodynamic turbulence did not give significant angular momentum transport needed even when the initial conditions had strong turbulence structures. Their 3D results suggest that accretion disks are well described by Euler's, ignoring the viscous terms from Navier Stokes.

Recent numerical simulations using similar models have shown that the resulting simulations are highly dependent on the numerical method used and resolution of the grid unless explicit diffusion coefficients and the appropriate dissipative scales are resolved (Fromang & Papaloizou 2007; Fromang *et al.* 2007). This issue has been considered in Lesur & Longaretti (2007) for cases with nonzero net vertical flux, where they found that both the viscosity and resistivity affect the amount of angular momentum transported by magnetohydrodynamic turbulence.

The use of the shearing box model has its limitations as pointed out by Hawley *et al.* (1995) as they do not permit any dynamics involving the background shear, which is taken as imposed and constant in space and time (Regev & Umurhan 2008). With regards to resolution which is tied in with the model used, the efficiency of angular momentum transport appears to decrease with improved resolution as shown by Fromang *et al.* (2007), suggesting that transport rates estimated on basis of ideal MHD are affected by grid-scale dissipation and overestimate the efficiency of angular momentum extraction by MRI. Kapyla & Korpi (2011) also worked on resolution dependence of  $\alpha$ , the Shakura-Sunyaev number and the possible decline of the angular momentum transport rate with decreasing Pm. Even though the shearing box model is restrictive in resolving the behavior of the accretion disk, it is nonetheless useful in understanding the dynamo process that sustains the magnetic field Lesur & Ogilvie (2008).

Squire's theorem says that for each unstable 3D disturbances, there are corresponding 2D disturbance that are more unstable. How does that apply here?

Non-axisymmetric numerical simulations done by Fleming *et al.* (2000) suggest saturation does occur even when Elsasser number  $\Delta \geq 1$  if non-axisymmetric disturbances are allowed to evolve.

Liu (2008) (also Princeton MRI experiments) simulated nonlinear development of MRI in a non-ideal magnetohydrodynamic Taylor-Couette flow (mimicking an on-going experiment) using



ZEUS-MP 2.0 code (Hayes *et al.* (2006)), which is a time-explicit, compressible, astrophysical ideal MHD parallel 3D code with added viscosity and resistivity for axisymmetric flows in cylindrical coordinates. He shows that the saturation of MRI causes a inflowing 'jet' feature which is opposite to the usual Ekman circulation and enhances angular momentum transport radially outward, which is what is needed, agreeing with HGB. Note: Stable MRI regime ( $Re \leq 1600$ ) enhances vertical angular momentum transport while in unstable MRI regime ( $Re \geq 3200$ ), MRI kicks in, resulting in more radial angular momentum transport as compared with vertical.

Kirillov *et al.* (2012) presents a unifying description of the helical and azimuthal versions of MRI, and they also identify the universal character of the 'Liu' limit  $2(1 - 2) \approx -0.8284$  for the critical Rossby number. From this universal characteristics, they are led to the prediction that the instability will be governed by a mode with an azimuthal wavenumber that is proportional to the ratio of axial to azimuthal applied magnetic field, when this ratio becomes large and the Rossby number is close to the Liu limit)

Miller & Stone (1999) worked on vertically stratified disks using a shearing box model and found that turbulent magnetised disk can produce a magnetised corona in laminar flow through MRI. Stone *et al.* (1996) presented a 3D MHD simulation of the nonlinear evolution of MRI in a local shearing box method with a global vertical direction and vertically stratified disks. They found that the instability generated and maintained MHD turbulence.

From Julien & Knobloch (2010): Reduced equations- Case A ( $\delta = \epsilon, \Lambda = O(1)$  )

- a) Single Mode theory
- b) Stress-free boundary conditions
- c) Random initial conditions

Reduced equations- Case B ( $\epsilon = O(\delta), \Lambda \gg 1$  )

- a) Single Mode Solutions
- b) Multi-mode Solutions
- c) Random initial Perturbations
- d) Dissipation and Saturation

## 6. Conclusion

Concluding remarks

## REFERENCES

- ACHESON, D.J. 1972 On the hydromagnetic stability of a rotating fluid annulus. *Journal of Fluid Mechanics* **52**, 529–541.
- ACHESON, D.J. 1973 Hydromagnetic wavelike instabilities in a rapidly rotating stratified fluid. *Journal of Fluid Mechanics* **61**, 609–624.
- ACHESON, D.J. & HIDE, R. 1973 Hydromagnetics of rotating fluids. *Reports on Progress in Physics* **36** (2), 159.
- BALBUS, S.A., GAMMIE-C.F. & HAWLEY, J.F. 1994 Fluctuations, dissipation and turbulence in accretion discs. *Monthly Notices of the Royal Astronomical Society* **271**, 197–201.
- BALBUS, S.A. & HAWLEY, J.F. 1991 A powerful local shear instability in weakly magnetized disks. i - linear analysis. ii - nonlinear evolution. *Astrophysical Journal* **376**, 214–233.
- BALBUS, S. A. 2003 Enhanced angular momentum transport in accretion disks. *Ann. Rev. Astron. Astrophys* **41**, 555–597.

- BALBUS, S. A. 2011 *Magnetohydrodynamics of Protostellar Disks*, pp. 237–282.
- BALBUS, STEVEN A. & HAWLEY, JOHN F. 1998 Instability, turbulence, and enhanced transport in accretion disks. *Rev. Mod. Phys.* **70**, 1–53.
- CHANDRASEKHAR, S. 1960 The stability of non-dissipative couette flow in hydromagnetics. *Proceedings of the National Academy of Science* **46**, 253–257.
- CHARRU, FRANCOIS & DE FORCRAND-MILLARD, PATRICIA 2011 *Hydrodynamic instabilities*, , vol. 37. Cambridge University Press.
- DAVIDSON, PETER ALAN 2001 *An introduction to magnetohydrodynamics*, , vol. 25. Cambridge university press.
- DONNELLEY, R.J. & OZIMA, M. 1960 Hydromagnetic stability of flow between rotating cylinders. *Phys. Rev. Lett.* **4**, 497–498.
- FLEMING, T.P., STONE, J.I.M. & HAWLEY, J.F. 2000 The effect of resistivity on the nonlinear stage of the magnetorotational instability in accretion disks. *Astrophysical Journal* **530**, 464–477.
- FREIDBERG, JEFFREY P. 1987 *Ideal magnetohydrodynamics*. Plenum Press, New York, NY.
- FROMANG, S. & PAPALOIZOU, J. 2007 Mhd simulations of the magnetorotational instability in a shearing box with zero net flux. *Astronomy & Astrophysics* **476** (3), 1113–1122.
- FROMANG, S., PAPALOIZOU, J., LESUR, G. & HEINEMANN, T. 2007 Mhd simulations of the magnetorotational instability in a shearing box with zero net flux - the effect of transport coefficients. *Astronomy & Astrophysics* **476** (3), 1123–1132.
- GOODMAN, J. & JI, H.T. 2002 Magnetorotational instability of dissipative couette flow. *Journal of Fluid Mechanics* **462**, 365–382.
- GOODMAN, J. & XU, G.H. 1994 Parasitic instabilities in magnetized, differentially rotating-disks. *Astrophysical Journal* **432**, 213–223.
- HAWLEY, J.F., GAMMIE, C.F. & BALBUS, S.A. 1995 Local 3-dimensional magnetohydrodynamic simulations of accretion disks. *Astrophysical Journal* **440**, 742–763.
- HAYES, JOHN C., NORMAN, MICHAEL L., FIEDLER, ROBERT A., BORDNER, JAMES O., LI, PAK SHING, CLARK, STEPHEN E., UD DOULA, ASIF & MAC LOW, MORDECAI-MARK 2006 Simulating radiating and magnetized flows in multiple dimensions with zeus-mp. *Astrophysical Journal Supplement Series* **165** (1), 188–228.
- HOLLERBACH, RAINER & RUDIGER, GUNTHER 2005 New type of magnetorotational instability in cylindrical taylor-couette flow. *Physical Review Letters* **95**.
- Ji, H.T., GOODMAN, J. & KAGEYAMA, A. 2001 Magnetorotational instability in a rotating liquid metal annulus. *Monthly Notices of the Royal Astronomical Society* **325** (2), L1–L5.
- JULIEN, K. & KNOBLOCH, E. 2010 Magnetorotational instability: recent developments. *Philosophical Transactions of the Royal Society A* **368**, 1607–1633.
- KAPYLA, P.J. & KORPI, M.J. 2011 Magnetorotational instability driven dynamos at low magnetic prandtl numbers. *Monthly Notices of the Royal Astronomical Society* **413** (2), 901–907.
- KIRILLOV, OLEG N., STEFANI, FRANK & FUKUMOTO, YASUhide 2012 A unifying picture of helical and azimuthal magnetorotational instability, and the universal significance of the liu limit. *The Astrophysical Journal* **756** (1), 83.
- KNOBLOCH, E. 1992 On the stability of magnetized accretion discs. *Monthly Notices of the Royal Astronomical Society* **255**, 25P–28P.
- LESUR, G. & LONGARETTI, P.Y. 2007 Impact of dimensionless numbers on the efficiency of magnetorotational instability induced turbulent transport. *Monthly Notices of the Royal Astronomical Society* **378** (4), 1471–1480.
- LESUR, GEOFFROY & OGILVIE, GORDON I. 2008 Localized magnetorotational instability and its role in the accretion disc dynamo. *Monthly Notices of the Royal Astronomical Society* **391** (3), 1437–1450.
- LIU, WEI 2008 Numerical study of the magnetorotational instability in princeton mri experiment. *The Astrophysical Journal* **684** (1), 515.
- MILLER, K.A. & STONE, J.M. 1999 3d mhd simulations of the magneto-rotational instability in vertically stratified disks. *Numerical Astrophysics* .
- RAYLEIGH, LORD 1916 On convection currents in horizontal layer of fluid when the higher temperature is on the under side. *Phil. Mag.* **32**, 529–546.

- REGEV, O. & UMURHAN, O.M. 2008 On the viability of the shearing box approximation for numerical studies of mhd turbulence in accretion disks. *Astronomy & Astrophysics* **481** (1), 21–32.
- SISAN, DANIEL R., MUJICA, NICOLÁS, TILLOTSON, W. ANDREW, HUANG, YI-MIN, DORLAND, WILLIAM, HASSAM, ADIL B., ANTONSEN, THOMAS M. & LATHROP, DANIEL P. 2004 Experimental observation and characterization of the magnetorotational instability. *Phys. Rev. Lett.* **93**, 114502.
- STEFANI, F., ECKERT, S., GERBETH, G., GIESECKE, A., GUNDRUM, TH., STEGLICH, C., WEIER, T. & WUSTMANN, B. 2012 DRESDYN - a new facility for mhd experiments with liquid sodium. *Magnetohydrodynamics* **48** (1), 103–113, 8th PAMIR International Conference on Fundamental MHD and Liquid Metal Technology, Borgo Corsica, France, SEP 05-09, 2011.
- STEFANI, FRANK, GERBETH, GUNTER, GUNDRUM, THOMAS, HOLLERBACH, RAINER, PRIEDE, J. ANIS, RÜDIGER, GÜNTHER & SZKLARSKI, JACEK 2009 Helical magnetorotational instability in a taylor-couette flow with strongly reduced ekman pumping. *Phys. Rev. E* **80**.
- STEFANI, FRANK, GUNDRUM, THOMAS, GERBETH, GUNTER, RÜDIGER, GÜNTHER, SCHULTZ, MANFRED, SZKLARSKI, JACEK & HOLLERBACH, RAINER 2006 Experimental evidence for magnetorotational instability in a taylor-couette flow under the influence of a helical magnetic field. *Phys. Rev. Lett.* **97**.
- STEFANI, FRANK, GUNDRUM, THOMAS, GERBETH, GUNTER, RDIGER, GNTER, SZKLARSKI, JACEK & HOLLERBACH, RAINER 2007 Experiments on the magnetorotational instability in helical magnetic fields. *New Journal of Physics* **9** (8).
- STONE, J.M., HAWLEY, J.F., GAMMIE, C.F. & BALBUS, S.A. 1996 Three-dimensional magnetohydrodynamical simulations of vertically stratified accretion disks. *The Astrophysical Journal* **463** (2), 656–673.
- VELIKHOV, E.P. 1959 Stability of an ideally conducting liquid flowing between cylinders rotating in a magnetic field. *Soviet Physics JETP-USSR* **9** (5), 995–998.
- WIKIPEDIA 2013 Magnetorotational instability — Wikipedia, the free encyclopedia. [Online; accessed 29-May-2013].