Homework 02 (26Sep22)

Rubric for each assignment:

Context	Points
Precision of the answer Answer Markdown readability Code readability	80% 10% 10%

Guidance:

• Upload your answers in the Blackboard submission portal as:

 $last name-first name-homework-xx.pdf \ {\rm or} \ last name-first name-homework-xx.ipynb$

Table of Problems

- Problem 1 (10 pts) D-He-3 fusion.
- 1.1)(5 pts) Q value.
- 1.2)(3 pts) Compare to D-T?
- 1.3)(2 pts) He-3 production.
- Problem 2 (35 pts) UMLRR Co-60 sphere source.
- 2.1)(5 pts) Activity.
- 2.2)(2 pts) Mass.
- 2.3)(5 pts) Final mass.
- 2.4)(3 pts) Mass loss.
- 2.5)(10 pts) Alternative mass loss.
- 2.6)(10 pts) Electricity conversion.
- Problem 3 (45 pts) Production of radionuclide.
- 3.1)(5 pts) Reaction.
- 3.2)(10 pts) Mathematical model.
- 3.3)(20 pts) Maximum activity.
- 3.4)(10 pts) Maximum flux.
- Problem 4 (10 pts) Heat decay for 1 GW(t) operation in 1 year.
- 4.1)(2 pts) 1 day.
- 4.2)(2 pts) 1 month.
- 4.3)(2 pts) 1 year.
- 4.4)(4 pts) Repeat for 6-month operation.

Problem 1 (20 pts)

1.1)(5 pts) A proposed fusion reaction is as follows: ${}^{2}\text{H} + {}^{3}\text{He} \rightarrow {}^{4}\text{He} + {}^{1}_{1}\text{p}$. Calculate the Q value of the reaction and the partition of energy of the products.

Answer:

1.2)(3 pts) Is this reaction a better proposition than the D-T reaction? Explain why or why not.

Explanation:

1.3)(2 pts) How can ³He be produced?

Answer:

Problem 2 (35 pts)

The UMLRR advertised a 60 Co sphere with activity $A=100\,\mathrm{kCi}$.

2.1)(5 pts) Does the activity stay the same with time? If not, what is the value in 1 year?

Answer:

2.2)(2 pts) Is the mass of the sphere after 1 year more than the initial mass, less or the same? Why?

Answer:

2.3)(5 pts) Calculate the initial mass of the sphere and final mass after 1 year.

Answer:

2.4)(3 pts) Calculate the mass loss.

Answer:

2.5)(10 pts) Find an alternative way to calculate the mass loss and compare with 2.4).

Answer:

2.6)(10 pts) If the energy emitted in one year by the ⁶⁰Co source could be converted in electricity, how long a typical residence in Massachussets could be powered for in 2022? (assume a conversion efficiency of 20%).

Answer:

Problem 3 (45 pts)

A stable nuclide X is reacted with particle n (e.g. a neutron) to give a radionuclide Y with radioactive decay constant λ_Y . Denote c_{X0} as the concentration of X when the reaction starts, assume there is no Y when the reaction starts, and that Y does not react with the particle n. It is of practical interest to know the maximum activity of Y and at what time the maximum is reached. Please address the sub-items below.

3.1)(5 pts) Write the reaction mechanism.

Answer:

3.2)(10 pts) Setup the mathematical model and state your assumptions.

Answer:

3.3)(20 pts) Calculate the maximum activity of Y and when it occurs relative to the beginning of the reaction.

Answer:

3.4)(10 pts) Often times when designing the irradiation apparatus for the reaction in Problem 3.1), the particle n spatial flux magnitude is wanted for a given production activity maximum $A_{Y,\max}$. Derive the equation needed to find this flux value and explain how to solve it.

Answer:

Problem 4 (10 pts)

A nuclear fission reactor operates at a power of 1.5 GW(t) for 1 year. Calculate the heat decay power:

4.1)(2 pts) 1 day after shutdown.

Answer:

4.2)(2 pts) 1 month after shutdown.

Answer:

4.3)(2 pts) 1 year after shutdown.

Answer:

4.4)(4 pts) Repeat the above for 6 months of operation and compare results.

Answer