# Quantum Love: The Interplay of Science with Emotion

Daniel Prelipcean

Jacobs University Bremen d.prelipcean@jacobs-university.de

August 30, 2020



# Physics is Fun!

"Physics is like sex: sure, it may give some practical results, but that's not why we do it."

# Physics is Fun!

"Physics is like sex: sure, it may give some practical results, but that's not why we do it."

(Richard Feynman)



Figure 1: Nobel Prize Picture <sup>1</sup>



<sup>&</sup>lt;sup>1</sup>wikipedia

### Motivation and Idea




Figure 2: Male Testosterone Levels <sup>2</sup>

### Motivation and Idea




Figure 2: Male Testosterone Levels <sup>2</sup>

<sup>&</sup>lt;sup>2</sup>become-the-alpha-male.com/blog/category/testosterone/ → ← ≥ → ← ≥ → へ ○ ○

### Disclaimer

Introduction

If you feel offended, let me know such that I can skip the respective part of my presentation.

If you feel offended, but do not want to speak up, let me know afterwards, such that I can change it and improve in future presentations.

The main object in Quantum Mechanics is the wavefunction  $\psi$ .



The main object in Quantum Mechanics is the wavefunction  $\psi$ .

$$|\psi\rangle$$

The main object in Quantum Mechanics is the wavefunction  $\psi$ .

$$|\psi\rangle$$

The main object in Quantum Mechanics is the wavefunction  $\psi$ .

$$\hat{\mathbb{O}} \ket{\psi}$$

The main object in Quantum Mechanics is the wavefunction  $\psi$ .

$$\hat{\mathbb{O}} \ket{\psi} = \mathbf{o} \ket{\psi}$$

The main object in Quantum Mechanics is the wavefunction  $\psi$ .

The main object in Quantum Mechanics is the wavefunction  $\psi$ .

$$\hat{\mathbb{O}} \left| \psi \right> = \mathbf{o} \left| \psi \right>$$
 operator — igenvalue



The main object in Quantum Mechanics is the wavefunction  $\psi$ .

$$\hat{\mathbb{O}} \left| \psi \right> = \mathbf{o} \left| \psi \right>$$
 operator — eigenvalue



The main object in Quantum Mechanics is the wavefunction  $\psi$ .

Employing Dirac notation, this is written as:

$$\hat{\mathbb{O}} \left| \psi \right> = \mathbf{o} \left| \psi \right>$$
 operator — igenvalue





= nervous



# States Spaces

Superposition Principle: Male State Space

$$|m(x)\rangle = \sum_{i=0}^{n} c_i |m_i(x)\rangle$$
 (1)

$$\sum_{i} c_{j}$$
  $|m_{j}(x)\rangle$ 

(2)

$$|w(k)\rangle = \sum_{j} c_{j}$$
  $|m_{j}(x)\rangle$ 

(2)

$$|w(k)\rangle = \sum_{j} c_{j}$$
  $|m_{j}(x)\rangle e^{-ikx}$  (2)

#### where:

• the exponential = our first attempt to understand something

$$|w(k)\rangle = \sum_{j} c_{j} \frac{1}{(\sqrt{2\pi})^{d}} \int d^{d}x |m_{j}(x)\rangle e^{-ikx}$$
 (2)

- the exponential = our first attempt to understand something
- d dimensions = cannot even comprehend in how many ways a woman can think

$$|w(k)\rangle = \sum_{j} c_{j} \frac{1}{(\sqrt{2\pi})^{d}} \int d^{d}x |m_{j}(x)\rangle e^{-ikx} \delta(k_{\mu}k^{\mu} + m^{2})$$
 (2)

- the exponential = our first attempt to understand something
- d dimensions = cannot even comprehend in how many ways a woman can think
- $\bullet$  the delta function  $\delta = {\rm engage}$  in a conversation only when appropriate



$$|w(k)\rangle = \sum_{j} c_{j} \frac{1}{(\sqrt{2\pi})^{d}} \int d^{d}x |m_{j}(x)\rangle e^{-ikx} \delta(k_{\mu}k^{\mu} + m^{2})\Theta(k_{\mu}x^{\mu})$$
(2)

- the exponential = our first attempt to understand something
- d dimensions = cannot even comprehend in how many ways a woman can think
- $oldsymbol{\circ}$  the delta function  $\delta=$  engage in a conversation only when appropriate
- **4** step function  $\Theta(\mu) = \text{apologize when mistaken}$



Operators

## The Love operator

Let  $\hat{L}$  be the Love operator, with the desired property:

# The Love operator

Let  $\hat{L}$  be the Love operator, with the desired property:

$$\hat{L}|w\rangle = (\text{loves me})|w\rangle$$
 (3)

Let  $\hat{L}$  be the Love operator, with the desired property:

$$\hat{L}|w\rangle = (\text{loves me})|w\rangle$$
 (3)

Problem: Hard to find exact representation of  $\hat{L}$  for different ladies  $|w_i\rangle$ .

Let  $\hat{L}$  be the Love operator, with the desired property:

$$\hat{L}|w\rangle = (\text{loves me})|w\rangle$$
 (3)

Problem: Hard to find exact representation of  $\hat{L}$  for different ladies  $|w_j\rangle$ .

### Daniel's Conjecture:



# The Love operator

Let  $\hat{L}$  be the Love operator, with the desired property:

$$\hat{L}|w\rangle = (\text{loves me})|w\rangle$$
 (3)

Problem: Hard to find exact representation of  $\hat{L}$  for different ladies  $|w_j\rangle$ .

**Daniel's Conjecture:** One can find a suitable representation of  $\hat{L}$  only for a single woman  $|w\rangle = |g\rangle$ , that is, the girlfriend.



$$\hat{L}\ket{w}$$

$$\hat{L} |w\rangle \approx \widehat{\mathsf{Flowers}} |w\rangle$$

$$\hat{L} \ket{w} \approx \widehat{\mathsf{Flowers}} \ket{w} = \mathsf{impressed} \ket{w}$$

$$\hat{L} \ket{w} \approx \widehat{\mathsf{Flowers}} \ket{w} = \mathsf{impressed} \ket{w}$$

$$\hat{L} \ket{w} \approx \widehat{\mathsf{Flowers}} \ket{w} = \mathsf{impressed} \ket{w}$$

$$\hat{L}\ket{w} pprox \widehat{\mathsf{Flowers}}\ket{w} = \mathsf{impressed}\ket{w}$$

$$\hat{L} \ket{w} \approx \widehat{\mathsf{Flowers}} \ket{w} = \mathsf{impressed} \ket{w}$$

$$\hat{L}\ket{w}$$

$$\hat{L}|w\rangle \approx \widehat{\mathsf{Romantic Dinner}}|w\rangle$$

$$\hat{L} \ket{w} \approx \widehat{\mathsf{Romantic Dinner}} \ket{w} = \mathsf{impressed} \ket{w}$$

$$\hat{L}|w\rangle \approx \widehat{\mathsf{Romantic Dinner}}|w\rangle = \mathsf{impressed}|w\rangle$$

Pics/Nicoleta.jpg



$$\hat{L}|w\rangle \approx \widehat{\mathsf{Romantic Dinner}}|w\rangle = \mathsf{impressed}|w\rangle$$

$$\hat{L}|w\rangle \approx \widehat{\mathsf{Romantic Dinner}}|w\rangle = \mathsf{impressed}|w\rangle$$

$$\hat{L} \ket{w} \approx \widehat{\mathsf{Romantic Dinner}} \ket{w} = \mathsf{impressed} \ket{w}$$

Operators may change the state completely, e.g. the cheating operator  $\hat{C}$ .

Operators



Operators may change the state completely, e.g. the cheating operator  $\hat{\mathcal{C}}$ .

Let  $|g\rangle$  be your girlfriend, and  $|w\rangle$  be another girl. Then:

$$\hat{S} |w\rangle = \hat{C} |g\rangle = \tag{4}$$

Operators may change the state completely, e.g. the cheating operator  $\hat{\mathcal{C}}$ .

Let  $|g\rangle$  be your girlfriend, and  $|w\rangle$  be another girl. Then:

$$\hat{S} |w\rangle = \hat{C} |g\rangle = 0 |0\rangle = 0 \tag{4}$$

For eternity.



Operators may change the state completely, e.g. the cheating operator  $\hat{C}$ .

Let  $|g\rangle$  be your girlfriend, and  $|w\rangle$  be another girl. Then:

$$\hat{S} |w\rangle = \hat{C} |g\rangle = 0 |0\rangle = 0 \tag{4}$$

For eternity.

**Conclusion:** Not all operator equations are eingenvalue problems.

Let  $\hat{S}$  be the sex operator, with the desired property:

$$\hat{S}|g\rangle = \sec|g\rangle \tag{5}$$

Let  $\hat{S}$  be the sex operator, with the desired property:

$$\hat{S}|g\rangle = \sec|g\rangle \tag{5}$$

However, it can be the case that:

$$\hat{S}|g\rangle = \text{harrasment}|g\rangle$$
 (6)

So be very careful with this one!

Let  $\hat{S}$  be the sex operator, with the desired property:

$$\hat{S}|g\rangle = \sec|g\rangle \tag{5}$$

However, it can be the case that:

$$\hat{S}|g\rangle = \text{harrasment}|g\rangle$$
 (6)

So be very careful with this one!

**Hopeful Lemma:** As  $t \to \infty$ , the above case will happen with probability P = 0, that is, never again.



Practically, one may have to apply it multiple times to reach the desired result:

$$\hat{S}^n |g\rangle = \sec |g\rangle \tag{7}$$

Practically, one may have to apply it multiple times to reach the desired result:

$$\hat{S}^n |g\rangle = \sec |g\rangle \tag{7}$$

Scientists are especially interested in limit cases:



Practically, one may have to apply it multiple times to reach the desired result:

$$\hat{S}^n |g\rangle = \sec |g\rangle \tag{7}$$

Scientists are especially interested in limit cases: e.g. n = 0, i.e. you do not have to do anything and get:

$$\hat{S}^{0}|g\rangle = |g\rangle = sex|g\rangle \tag{8}$$

Practically, one may have to apply it multiple times to reach the desired result:

$$\hat{S}^n |g\rangle = \sec |g\rangle \tag{7}$$

Operators

Scientists are especially interested in limit cases: e.g. n = 0, i.e. you do not have to do anything and get:

$$\hat{S}^{0}|g\rangle = |g\rangle = \sec|g\rangle \tag{8}$$

Open question: Prove whether such states exist (or not).



A more realistic scenario is the following:

$$\hat{L}^{n}|g\rangle = \prod_{j=1}^{N} (L_{j})|g\rangle = (\text{in love})|g\rangle$$
(9)

A more realistic scenario is the following:

$$\hat{L}^{n}|g\rangle = \prod_{j=1}^{N} (L_{j})|g\rangle = (\text{in love})|g\rangle$$
(9)

That is, one has to apply it multiple times, possibly using different representations or common approximations (as before).



A more realistic scenario is the following:

$$\hat{L}^{n}|g\rangle = \prod_{j=1}^{N} (L_{j})|g\rangle = (\text{in love})|g\rangle$$
 (9)

That is, one has to apply it multiple times, possibly using different representations or common approximations (as before).

Most interesting case:  $n \to \infty$ .



A more realistic scenario is the following:

$$\hat{L}^{n}|g\rangle = \prod_{j=1}^{N} (L_{j})|g\rangle = (\text{in love})|g\rangle$$
 (9)

That is, one has to apply it multiple times, possibly using different representations or common approximations (as before).

Most interesting case:  $n \to \infty$ .

#### Problem:

How to apply an infinite amount of work?



A more realistic scenario is the following:

$$\hat{L}^{n}|g\rangle = \prod_{j=1}^{N} (L_{j})|g\rangle = (\text{in love})|g\rangle$$
(9)

That is, one has to apply it multiple times, possibly using different representations or common approximations (as before).

Most interesting case:  $n \to \infty$ .

#### Problem:

How to apply an infinite amount of work? ( $\approx 0$  in her eyes)



## The Love operator solution

Answer: Apply the Interest operator  $\hat{I}$ :

$$\hat{I}|g\rangle = (\text{interested in me})|g\rangle$$
 (10)

## The Love operator solution

Answer: Apply the Interest operator  $\hat{I}$ :

$$\hat{I}|g\rangle = (\text{interested in me})|g\rangle$$
 (10)

such that she wants to apply to Love operator on me, that is:

$$\hat{L}_{w} |b\rangle = (?) |b\rangle \tag{11}$$

Quantum Physics in a nutshell:

Quantum Physics in a nutshell:

Define your wave functions in a suitable basis.



Quantum Physics in a nutshell:

- Define your wave functions in a suitable basis.
- Investigate how operators apply on the wave functions (and pay much attention how you can apply them).

### Quantum Physics in a nutshell:

- Define your wave functions in a suitable basis.
- Investigate how operators apply on the wave functions (and pay much attention how you can apply them).
- Obtain physical results that can be experimentally achieved.

# Acknowledgements

I acknowledge



Conclusion

## Acknowledgements

I acknowledge psychological support from my girlfriend Nicoleta (JUB, Class of 2017)

☐Pics/grad.JPG

Figure 3: Success!  $(n \gg 1)$ 

## Acknowledgements

☐Pics/grad.JPG

I acknowledge psychological support from my girlfriend Nicoleta (JUB, Class of 2017) and financial support from my parents, through the Grant RO-B961217.

Figure 3: Success!  $(n \gg 1)$ 

□Pics/fam.jpg

Figure 4: Prelipcean Family

Humans



Humans = social beings, who urge to feel love.



Humans = social beings, who urge to feel love.

Tell "I love you" more often, and louder!



Conclusion

Humans = social beings, who urge to feel love.

Tell "I love you" more often, and louder!

Even the most aesthetic equation cannot compare to a sincere "I love you"

from your loved and

from your loved ones.



Humans = social beings, who urge to feel love.

Tell "I love you" more often, and louder!

Even the most aesthetic equation cannot compare to a sincere "I love you"

from your loved and

from your loved ones.

Thank You and May the Force be with You!

