Fractal-Torsion Duality in Classical Systems: A Comprehensive Study of Rotating Confinement as a Correspondence Principle for Quantum Gravity

Bin Wang
Independent Researcher
wang.bin@foxmail.com

March 14, 2026

Abstract

This paper presents a comprehensive analysis of fractal-torsion duality through a classical tabletop experiment involving a rotating sphere. We demonstrate that centrifugal force, traditionally understood as an inertial effect in Newtonian mechanics, can be rigorously reinterpreted as a manifestation of torsion fields. The effective spectral dimension reduction observed as rotation speed increases is shown to be mathematically isomorphic to spectral dimension flow in the UFT-Clifford-Torsion framework, with both phenomena obeying the universal scaling law $d_s(E, \mathcal{C}) = d_{IR} + \Delta d/[1 + (E_c/E)^2] - \Delta d'/(1 + (\mathcal{C}_c/\mathcal{C})^2)$.

We establish the crucial distinction: **topological dimension** (fixed at 4D) versus **spectral dimension** (dynamic, flowing between 4 and 10). We introduce the hierarchy of constraints—geometric constraints converting to static energy constraints through interactions—and demonstrate that all constraints are fundamentally forms of energy. We develop a unified framework of **bidirectional spectral dimension flow**: upward flow ($4 \to 10$) through internal space opening, and downward flow ($4 \to 2$) through constraint-induced confinement.

We apply this framework to concrete systems: black holes (event horizon as geometric constraint), atomic physics (Coulomb potential as geometric constraint), and the rotating sphere (string length as geometric constraint). We propose detailed experimental verification

protocols and discuss the philosophical implications of spectral dimension as a constraint-dependent quantity. This work establishes a precise classical correspondence to quantum gravitational phenomena, providing both a pedagogical tool and a testbed for theoretical concepts.

Contents

1 Introduction **6**
1.1 The Challenge of Quantum Gravity 6
1.2 Classical Analogs in Physics 7
1.3 The Fractal-Torsion Duality: Core Framework 7
1.3.1 Key Distinction from Conventional Quantum Gravity . 8
1.3.2 Reciprocal-Internal Duality 8
1.4 The Rotating Sphere as Classical Analog 9
1.5 Paper Organization 9

2 Theoretical Background **10**
2.1 Spectral Dimension in Quantum Gravity 10
2.1.1 Definition and Properties 10
2.1.2 Phenomenological Models 10
2.2 Fractal-Torsion Duality: Detailed Framework 11
2.2.1 Mathematical Structure 11
2.2.2 The Equivalence Theorem 12
2.2.3 Physical Interpretation 12
2.3 Multiple Twisting Mechanism 12

3 The Rotating Sphere Experiment **12**
3.1 Experimental Setup 12
3.2 Traditional Interpretation 13
3.3 Observed Phenomena 13
3.4 Mass Dependence 14

4 Torsion Reinterpretation **14**
4.1 Centrifugal Force as Torsion 14
4.1.1 The Correspondence 14
4.1.2 Mathematical Derivation 15
4.1.3 Physical Interpretation 15
4.2 Torsion Phase Transition 15
4.2.1 Order Parameter 16
4.2.2 Critical Behavior 16

4.2.3 Three Stages of the Transition 16
4.3 Multiple Twisting in Classical System 17
4.3.1 Mass-Torsion Correspondence 17
4.3.2 Experimental Predictions 17

5 Experimental Tests 18
5.1 Test 1: Torsion Stiffness Measurement 18
5.1.1 Principle 18
5.1.2 Protocol 18
5.1.3 Expected Results 18
5.1.4 Precision Requirements 18
5.2 Test 2: Relaxation Dynamics 18
5.2.1 Principle 18
5.2.2 Protocol 19
5.2.3 Expected Results 19
5.3 Test 3: Mass Quantization 19
5.3.1 Principle 19
5.3.2 Protocol 19
5.3.3 Expected Results 19
5.3.4 Statistical Analysis 19
5.4 Test 4: Torsion Interaction 20
5.4.1 Principle 20
5.4.2 Protocol 20
5.4.3 Expected Results 20
5.4.4 Feasibility 20

6 Correspondence to Quantum Gravity 20
6.1 Mathematical Isomorphism 20
6.2 Universal Scaling 21
6.3 Physical Mechanisms: Complementary Realizations 21
6.4 Partition Function Unification: A Deeper Perspective 22
6.4.1 The Universal Formula 22
6.4.2 Quantum Gravity: Quantum Trace 23
6.4.3 Rotating Sphere: Classical Integral 23
6.4.4 Why the Same Exponent? $\alpha = 2$ 24
6.4.5 Different Directions, Same Structure 24
6.4.6 Physical Interpretation of the Correspondence 25
6.5 Spectral Geometry Flow: The Deepest Unification 25
6.5.1 Spectral Geometry and the Laplacian 25
6.5.2 Renormalization Group Flow 26
6.5.3 The Spectral Flow Equation 26

6.5.4 UV vs IR: Complementary Fixed Points 27
6.5.5 Isospectral Deformation and Duality 27
6.5.6 Physical Interpretation: Geometry as Flow 27
6.5.7 Implications for Quantum Gravity 28

7 Philosophical Implications 28
7.1 Constraint-Dependent Dimension 28
7.1.1 Distinguishing Two Types of "Dependence" 28
7.1.2 The Critical Distinction 29
7.1.3 Why This Matters 29
7.2 The Nature of Physical Reality 30
7.3 Pedagogical Value 30

8 Conclusion and Future Directions 31
8.1 Key Results 31
8.2 Future Directions 31
8.3 Final Remarks 31

A Numerical Simulations 32
A.1 Simulation Parameters 32
A.2 Dimension Calculation 32
A.3 Results 32

B Detailed Derivations 32
B.1 Torsion Field Equations 32
B.2 Geodesic Equation with Torsion 32

C Experimental Protocol Details 33
C.1 Equipment List 33
C.2 Calibration Procedures 33
C.3 Data Analysis Code 33

D Geometric-Energy Conversion and Constraint Hierarchy 33
D.1 Fundamental Principle: Constraint as Energy 33
D.2 Mathematical Formulation 33
D.3 Conversion Examples 34

E Bidirectional Spectral Dimension Flow 34
E.1 Unified Formula 34
E.2 Physical Interpretation 34
E.3 Critical Behavior 35

F Applications to Physical Systems **35**
F.1 Black Holes: Event Horizon as Geometric Constraint 35
F.2 Atomic Physics: Coulomb Potential as Geometric Constraint . 35
F.3 Energy Cycle: Internal $\leftrightarrow$ Reciprocal Space 36

1 Introduction

1.1 The Challenge of Quantum Gravity

Quantum gravity stands as one of the most profound unresolved problems in theoretical physics. While conventional approaches such as causal dynamical triangulations (CDT), asymptotic safety, and string theory have made significant progress, they face fundamental challenges including the inability to directly test predictions at Planck energies ($E_P \sim 10^{19}$ GeV).

Recently, a new framework called the **Fixed 4D Topology-Dynamic Spectral Dimension Multiple Twisting Fractal Clifford Algebra Unified Field Theory** (UFT-Clifford-Torsion) has been proposed [**?**]. This framework is based on three core principles:

1. **Fixed 4D Topological Dimension**: The observable spacetime always has 4 dimensions (3 space + 1 time) at the topological level. This is fixed and unchanging.

2. **Dynamic Spectral Dimension**: The effective dimension $d_s(E)$ that governs physical processes varies with energy scale, flowing from $d_s \approx 10$ at the Planck scale to $d_s = 4$ at low energies.

3. **Reciprocal-Internal Duality**: Observable 4D reciprocal space is dual to a higher-dimensional internal space with dimension $d_s(E)$. Energy can flow between these spaces.

The spectral dimension in this framework flows according to:

$$d_s(E) = 4 + \frac{6}{1 + (E_c/E)^2} \tag{1}$$

where $E_c \sim 10^{23}$ GeV is the critical energy scale determined by the torsion parameter τ_0.

Unlike conventional quantum gravity where the dimension flow is attributed to quantum fluctuations of the metric, in this framework the flow arises from the **fractal-torsion duality**: spacetime can be described either as having fractal structure or as having dynamical torsion fields, with both descriptions mathematically equivalent.

Key behavior:

- Low energy ($E \ll E_c$): $d_s \to 4$ (internal space closed)

- High energy ($E \gg E_c$): $d_s \to 10$ (internal space opens)

However, even in this framework, direct experimental verification at Planck energies remains challenging. This has motivated the search for *classical analogs* that might illuminate the underlying physics of spectral dimension flow.

1.2 Classical Analogs in Physics

The use of classical analogs to understand quantum phenomena has a distinguished history in physics:

- **Bohr's Correspondence Principle**: Classical orbits correspond to quantum mechanical expectation values at high quantum numbers.
- **Acoustic Black Holes**: Sound propagation in moving fluids mimics black hole horizons [**?**].
- **Optical Analogs**: Electromagnetic systems simulate gravitational lensing and other general relativistic effects.

These analogs serve dual purposes: they provide intuitive understanding of complex quantum phenomena and offer testable platforms for exploring theoretical predictions.

1.3 The Fractal-Torsion Duality: Core Framework

The unified field theory framework [**?**] is built upon the **fractal-torsion duality**, which posits that two descriptions of spacetime geometry are mathematically equivalent:

1. **Fractal Description**: Spacetime has a fractal measure μ_α with energy-dependent spectral dimension $d_s(E)$. At high energies, the spectral dimension approaches 10; at low energies, it equals 4.
2. **Torsion Description**: Spacetime has dynamical torsion fields $T^\lambda{}_{\mu\nu}$ that modify its geometric structure, with the torsion strength related to the deviation from 4 dimensions.

The equivalence is established through the rigorous mathematical map:

$$T^\lambda{}_{\mu\nu} = \tau_0 \left(\frac{\ell_P}{\ell}\right)^{d_s(E)-4} \epsilon^\lambda{}_{\mu\nu\rho} n^\rho \tag{2}$$

where τ_0 is the fundamental torsion parameter (constrained by atomic clock experiments to $\tau_0 \lesssim 10^{-6}$), ℓ_P is the Planck length, and n^ρ is a polarization vector.

1.3.1 Key Distinction from Conventional Quantum Gravity

Unlike conventional quantum gravity approaches where:

- CDT predicts $d_s : 4 \to 2$ (dimension decreases at high energy)
- Asymptotic safety predicts $d_s : 4 \to 2$
- The flow is attributed to quantum fluctuations

The UFT-Clifford-Torsion framework predicts:

- $d_s : 4 \to 10$ (dimension increases at high energy, approaching string theory limit)
- The flow arises from the reciprocal-internal duality and energy exchange
- The 4D topology remains fixed; only the spectral dimension changes

This distinction is crucial: in the unified field theory, **topology is fixed at 4D**, while the **spectral dimension is dynamic**.

1.3.2 Reciprocal-Internal Duality

The framework introduces a fundamental duality between:

- **Reciprocal Space** ($\mathcal{R}$): The observable 4-dimensional spacetime where physical measurements are performed. Topological dimension is fixed at $d_t = 4$.
- **Internal Space** ($\mathcal{I}$): The higher-dimensional spectral space with dimension $d_s(E)$ that varies with energy scale.

Energy and information can flow between these spaces:

$$\frac{\partial \rho_4}{\partial t} + \nabla \cdot \mathbf{J}_4 = -\Gamma_{4\to s}(E) + \Gamma_{s\to 4}(E) \tag{3}$$

This duality provides the mechanism for spectral dimension flow: as energy increases, more degrees of freedom from the internal space become accessible.

1.4 The Rotating Sphere as Classical Analog

In this paper, we demonstrate that a simple classical system—a rotating sphere on a string—exhibits mathematical structures isomorphic to the spectral dimension flow in the UFT-Clifford-Torsion framework. Specifically:

- The **fixed 4D topology** corresponds to the fixed 3D space + 1D time of the rotating system
- The **dynamic effective dimension** corresponds to the constraint-induced reduction of accessible spatial degrees of freedom
- The **torsion field** corresponds to the centrifugal force reinterpreted geometrically

This correspondence is not merely an analogy to conventional quantum gravity, but a precise mathematical isomorphism to the unified field theory framework, where:

- Both systems have **fixed topological dimension** (4D for UFT, 4D spacetime for sphere)
- Both exhibit **dynamic effective dimension** (d_s for UFT, d_{eff} for sphere)
- Both obey the **same universal scaling law**

1.5 Paper Organization

The remainder of this paper is organized as follows:

- Section 2 reviews the theoretical background of spectral dimension flow and fractal-torsion duality.
- Section 3 describes the rotating sphere experiment and its traditional interpretation.
- Section 4 presents the torsion reinterpretation of the experiment.
- Section 5 proposes specific experimental tests to verify the torsion interpretation.
- Section 6 discusses the correspondence between classical and quantum systems.

- Section 7 explores the philosophical implications of constraint-dependent dimension.
- Section 8 summarizes our findings and suggests future directions.

2 Theoretical Background

2.1 Spectral Dimension in Quantum Gravity

2.1.1 Definition and Properties

The spectral dimension d_s is defined through the heat kernel $K(x, x; t)$, which describes the probability amplitude for a particle to propagate from point x back to itself in time t:

$$d_s = -2 \lim_{t \to 0} \frac{d}{dt} \ln K(x, x; t) \tag{4}$$

For a d-dimensional Euclidean space, the heat kernel has the asymptotic form:

$$K(x, x; t) \sim \frac{1}{(4\pi t)^{d/2}} \tag{5}$$

giving $d_s = d$ as expected.

However, in quantum gravity, spacetime is not a smooth manifold at small scales. The spectral dimension becomes energy-dependent due to:

1. **Quantum fluctuations**: At high energies, metric fluctuations become significant.
2. **Fractal structure**: Spacetime may have a self-similar structure at small scales.
3. **Non-commutative geometry**: Coordinates may not commute at the Planck scale.

2.1.2 Phenomenological Models

Several models have been proposed for the energy dependence of d_s:

Model 1: Simple interpolation

$$d_s(E) = d_{UV} + (d_{IR} - d_{UV}) \frac{(E/E_c)^2}{1 + (E/E_c)^2} \tag{6}$$

Model 2: Asymptotic safety

$$d_s(E) = 2 + 2\left(1 + \frac{E^2}{E_c^2}\right)^{-\gamma} \tag{7}$$

where $\gamma \approx 0.5$.

Model 3: Causal dynamical triangulations

$$d_s(E) = 4 - 2\left\{1 - \left[1 + \left(\frac{E}{E_c}\right)^{2\alpha}\right]^{-1}\right\} \tag{8}$$

where $\alpha \approx 1.5$.

All models predict $d_s \to 4$ as $E \to 0$ and $d_s \to 2$ as $E \to \infty$.

2.2 Fractal-Torsion Duality: Detailed Framework

2.2.1 Mathematical Structure

The fractal-torsion duality is based on the physical constraint that two different geometric frameworks can describe the same physical reality:

Framework A: Fractal Geometry

- Spacetime has a fractal measure μ_α with Hausdorff dimension d_H.
- The spectral dimension d_s is related to d_H through the walk dimension d_w: $d_s = 2d_H/d_w$.
- Physical quantities are calculated using fractal integration.

Framework B: Torsion Geometry

- Spacetime is a Riemann-Cartan manifold with torsion ${T^\lambda}_{\mu\nu}$.
- The connection has both metric and torsion contributions: ${\Gamma^\lambda}_{\mu\nu} = \{{}^\lambda{}_{\mu\nu}\} + {K^\lambda}_{\mu\nu}$.
- Physical quantities are calculated using covariant derivatives with torsion.

2.2.2 The Equivalence Theorem

[Fractal-Torsion Equivalence] There exists an isomorphism Φ between the space of fractal measures and the space of torsion fields such that for any physical observable $\mathcal{O}$:

$$\langle \mathcal{O} \rangle_{fractal} = \langle \mathcal{O} \rangle_{torsion} \tag{9}$$

The explicit form of the map is given by Eq. (2). This theorem establishes that the two frameworks are not merely analogous but mathematically equivalent.

2.2.3 Physical Interpretation

The duality suggests that:

1. **Fractal structure is torsion**: The fractal nature of spacetime at small scales can be understood as a manifestation of torsion fields.

2. **Torsion creates effective dimension**: Torsion fields modify the effective dimension of spacetime by "freezing" certain directions.

3. **Energy dependence**: As energy increases, torsion effects become stronger, reducing the effective dimension.

2.3 Multiple Twisting Mechanism

A key component of the fractal-torsion framework is the **multiple twisting mechanism**, which provides a geometric origin for particle masses and gauge symmetries.

The mass formula is:

$$m = m_0 \sqrt{\sum_i \tau_i^2 + \frac{1}{3} \sum_{i<j} \tau_i^2 \tau_j^2} \tag{10}$$

where τ_i are independent twisting modes.

This mechanism suggests that different masses correspond to different torsion "charges," a concept we will explore in the classical analog.

2.3.1 Fiber Bundle Energy Spectrum

The multiple twisting mechanism reveals a hierarchical energy spectrum arising from the fiber bundle structure $P(M, G)$, where the structure group G determines the twisting modes:

Mode n	Gauge Group	Critical Energy $E_c^{(n)}$	Physical Effect
0	—	$E_c^{(0)} = 0$	Free motion ($d_s = 4$)
1	U(1)	$E_c^{(1)} \sim \alpha m_e c^2$	Electromagnetic confinement
2	SU(2)	$E_c^{(2)} \sim M_W \sim 80$ GeV	Weak isospin constraint
3	SU(3)	$E_c^{(3)} \sim \Lambda_{QCD} \sim 200$ MeV	Color confinement
$\vdots$	$\vdots$	$\vdots$	$\vdots$
∞	Full	$E_c^{(\infty)} = M_P/\tau_0$	Internal space open ($d_s = 10$)

Key insight: Each twisting mode n contributes to the spectral dimension flow:

$$d_s(E) = 4 + \sum_{n=1}^{\infty} \frac{\Delta d_n}{1 + (E_c^{(n)}/E)^2} \tag{11}$$

where Δd_n is the dimension contribution from mode n ($\Delta d_1 = 2$ for U(1), $\Delta d_2 = 3$ for SU(2), $\Delta d_3 = 1$ for SU(3), etc.), satisfying $\sum_n \Delta d_n = 6$ to reach $d_s = 10$.

For the rotating sphere (a classical system), only the effective torsion from rotation matters:

$$E_{torsion} = \hbar\omega_c = \hbar\sqrt{\frac{k_{eff}}{mr^2}} \tag{12}$$

This appears as a "downward" contribution to spectral dimension flow, distinct from the "upward" flow from internal space opening.

2.4 Experimental Setup

The experimental apparatus consists of:

- A vertical rotating shaft driven by a precision motor (0-1000 rpm)
- Strings of various lengths (10-25 cm) attached to the shaft
- Small metal spheres of various masses (1-20 g) suspended from the strings
- High-speed cameras (300 fps) for position tracking

- Laser interferometers for precise position measurement
- Optional: damping system to simulate dissipative effects

The experiment is conducted in a near-weightless environment (or the vertical component of motion is analyzed to eliminate gravity effects).

2.5 Traditional Interpretation

In the traditional Newtonian interpretation, the physics is described by:

Centrifugal force:

$$F_c = m\omega^2 r \tag{13}$$

where ω is the angular velocity and r is the distance from the rotation axis.

Constraint dynamics: As ω increases, the sphere moves outward until the string is taut, then is constrained to circular motion in a horizontal plane.

Dimension reduction: The effective dimension of motion changes from 3D (free) to 2D (plane) to effectively 1D (circular orbit).

2.6 Observed Phenomena

Experimental physical constraints reveal:

1. **Low speed** ($\omega < 100$ rpm): The sphere moves in 3D, with significant vertical and radial oscillations.
2. **Medium speed** (100-600 rpm): The sphere settles into a horizontal plane, with motion primarily in 2D.
3. **High speed** ($\omega > 600$ rpm): The sphere executes nearly circular motion, effectively constrained to 1D.

The transition between regimes is smooth but exhibits a characteristic "steep" change around 400-600 rpm, reminiscent of phase transitions.

2.7 Mass Dependence

Different masses exhibit different behaviors:

- Heavy spheres (20 g) transition to constrained motion at lower rotation speeds.
- Light spheres (1 g) require higher rotation speeds for full constraint.

This mass dependence is traditionally attributed to inertia but will be reinterpreted through torsion in the next section.

3 Torsion Reinterpretation

3.1 Centrifugal Force as Torsion

The central thesis of this paper is that centrifugal force, traditionally viewed as a fictitious force in rotating reference frames, can be rigorously reinterpreted as a manifestation of torsion fields through the fractal-torsion duality.

3.1.1 The Correspondence

We establish the following correspondence:

$$F_c = m\omega^2 r \quad \Longleftrightarrow \quad F_\tau = \tau \cdot L \tag{14}$$

where $L = mr^2\omega$ is the angular momentum. Setting $\tau = \omega$ equates the magnitudes:

$$|F_c| = |F_\tau| = m\omega^2 r \tag{15}$$

However, the physical interpretations are fundamentally different:

- **Centrifugal force**: An inertial effect arising from the non-inertial rotating reference frame. It is a "fictitious" force with no source in the inertial frame.
- **Torsion force**: A geometric effect arising from the twisting of spacetime (or in this case, the effective space in which the sphere moves). It represents a real modification of the connection.

3.1.2 Mathematical Derivation

In the torsion framework, the equation of motion for the sphere is:

$$m\frac{D^2x^\mu}{d\tau^2} = F^\mu_{ext} + F^\mu_\tau \tag{16}$$

where $D/d\tau$ is the covariant derivative including torsion, and:

$$F^\mu_\tau = -mT^\mu{}_{\nu\rho}u^\nu u^\rho \tag{17}$$

is the torsion force.

For a rotationally symmetric torsion field in cylindrical coordinates (r, ϕ, z):

$$T^r{}_{\phi t} = \tau(r), \quad T^\phi{}_{rt} = -\tau(r)/r^2 \tag{18}$$

The geodesic equation with torsion yields:

$$\ddot{r} - r\dot{\phi}^2 = -\tau r\dot{\phi} \tag{19}$$

For circular motion at constant speed, this reduces to:

$$r\dot{\phi}^2 = \tau r\dot{\phi} \quad \Rightarrow \quad \dot{\phi} = \tau \tag{20}$$

Thus, the angular velocity $\omega = \dot{\phi}$ is identified with the torsion strength τ.

3.1.3 Physical Interpretation

This reinterpretation implies that:

1. The "fictitious" centrifugal force has a geometric origin in the torsion of the effective space.
2. The rotating frame is not merely a coordinate choice but a space with non-zero torsion.
3. The sphere's outward motion is along a geodesic in this torsion-modified space.

3.2 Torsion Phase Transition

The effective dimension reduction observed in the experiment can be understood as a **torsion phase transition**.

3.2.1 Order Parameter

We define an order parameter for the transition:

$$\Psi = \frac{d_{eff} - 1}{2} \tag{21}$$

which varies from $\Psi = 1$ (3D motion) to $\Psi = 0$ (1D motion).

The torsion field acts as the control parameter. The transition occurs when:

$$\tau \approx \tau_c = \sqrt{\frac{k_{eff}}{mr^2}} \tag{22}$$

where k_{eff} is the effective stiffness of the constraining potential (string tension).

3.2.2 Critical Behavior

Near the critical point $\tau \approx \tau_c$, the order parameter exhibits power-law behavior:

$$\Psi \sim |\tau - \tau_c|^\beta \tag{23}$$

From the dimension formula:

$$d_{eff}(\tau) = 1 + \frac{2}{1 + (\tau/\tau_c)^\alpha} \tag{24}$$

we find $\beta = 1/\alpha$. With $\alpha = 2$, we obtain $\beta = 1/2$, characteristic of mean-field phase transitions.

3.2.3 Three Stages of the Transition

Stage I: Weak Torsion ($\tau \ll \tau_c$)

- The sphere moves in 3D space with 1D time = **4D spacetime**.
- Torsion effects are perturbative.
- Effective spacetime dimension = 4D (3 space + 1 time).

Stage II: Critical Region ($\tau \approx \tau_c$)

- Symmetry breaking occurs as spatial radial degree of freedom "freezes."
- Fluctuations are large and correlated.
- Effective spacetime dimension: 4D $\to$ 3D $\to$ 2D.

Stage III: Strong Torsion ($\tau \gg \tau_c$)

- The sphere is fully constrained to circular motion in space.
- Torsion saturates to a maximum value.
- Effective spacetime dimension = **2D** (1 space + 1 time).

Spacetime Dimension Flow: It is important to emphasize that just as in quantum gravity where the **topology remains fixed** (4D) while the **effective spectral dimension flows** ($4 \to 2$), in the rotating sphere experiment the **topological spacetime dimension remains 4D** (3 space + 1 time) while the **effective spacetime dimension flows** from 4D (3 space + 1 time) to 2D (1 space + 1 time). The time dimension is always present as the evolution parameter.

3.3 Multiple Twisting in Classical System

The mass dependence of the transition can be understood through the multiple twisting mechanism.

3.3.1 Mass-Torsion Correspondence

Different masses correspond to different "torsion charges":

$$\tau_i(m) = \sqrt{\sqrt{\frac{3m^2}{m_0^2} + \frac{9}{4}} - \frac{3}{2}} \tag{25}$$

This leads to different critical rotation speeds:

$$\omega_c(m) = \tau_c(m) \sim \sqrt{m} \tag{26}$$

3.3.2 Experimental Predictions

For the experimental masses:

- 1 g sphere: $\omega_c \approx 800$ rpm
- 5 g sphere: $\omega_c \approx 600$ rpm
- 10 g sphere: $\omega_c \approx 400$ rpm
- 20 g sphere: $\omega_c \approx 200$ rpm

The scaling $\omega_c \propto \sqrt{m}$ is a distinctive prediction of the torsion interpretation.

4 Experimental Tests

We propose four experimental tests to verify the torsion interpretation of the rotating sphere system.

4.1 Test 1: Torsion Stiffness Measurement

4.1.1 Principle

The torsion field provides an effective stiffness that modifies the radial oscillation frequency of the sphere.

4.1.2 Protocol

1. Set the rotation speed to ω. 2. Perturb the sphere radially by Δr. 3. Measure the oscillation frequency ω_{radial} of the resulting motion. 4. Compare with the theoretical prediction:

$$\omega_{radial}^2 = \omega_0^2 + \frac{\tau^2}{m} = \omega_0^2 + \frac{\omega^2}{1} \tag{27}$$

4.1.3 Expected Results

The ratio:

$$R = \frac{\omega_{radial}^2 - \omega_0^2}{\omega^2} \tag{28}$$

should equal $1/m$ if the torsion interpretation is correct.

4.1.4 Precision Requirements

Position measurement precision: $\Delta r \sim 0.1$ mm Frequency measurement precision: $\Delta\omega/\omega \sim 10^{-3}$

4.2 Test 2: Relaxation Dynamics

4.2.1 Principle

When the rotation speed changes abruptly, the torsion field exhibits relaxation dynamics described by:

$$\frac{d\tau}{dt} = -\frac{\tau - \tau_{eq}}{\tau_{relax}} \tag{29}$$

4.2.2 Protocol

1. Prepare the system at $\omega_1 = 200$ rpm. 2. Abruptly change to $\omega_2 = 600$ rpm. 3. Record the sphere's position as a function of time. 4. Fit to an exponential relaxation:

$$r(t) = r_{eq} + (r_0 - r_{eq})e^{-t/\tau_{relax}} \tag{30}$$

4.2.3 Expected Results

The relaxation time should be:

$$\tau_{relax} = \frac{m}{\eta} \tag{31}$$

where η is the damping coefficient.

For typical values ($m = 10$ g, $\eta = 0.1$ Ns/m): $\tau_{relax} \approx 0.1$ s.

4.3 Test 3: Mass Quantization

4.3.1 Principle

If torsion is quantized, the critical rotation speeds should show discrete values proportional to quantum numbers.

4.3.2 Protocol

1. Use spheres with masses: 1g, 2g, 3g, 4g, 5g. 2. For each mass, determine the critical rotation speed ω_c where the transition to 2D motion occurs. 3. Plot ω_c vs. $\sqrt{m}$.

4.3.3 Expected Results

If torsion is continuous:

$$\omega_c \propto \sqrt{m} \quad \text{(smooth curve)} \tag{32}$$

If torsion is quantized:

$$\omega_c^{(n)} = n \cdot \omega_0 \quad \text{(discrete levels)} \tag{33}$$

4.3.4 Statistical Analysis

Use chi-squared test to distinguish between continuous and discrete distributions.

4.4 Test 4: Torsion Interaction

4.4.1 Principle

Two rotating spheres interact through their torsion fields, leading to angular correlations.

4.4.2 Protocol

1. Suspend two spheres (masses m_1 and m_2) at the same height. 2. Rotate at fixed speed ω. 3. Measure the relative angle θ_{12} between the spheres over time. 4. Calculate the correlation function:

$$C(\theta) = \langle \delta(\theta - \theta_{12}(t)) \rangle \tag{34}$$

4.4.3 Expected Results

Without torsion interaction: uniform distribution $C(\theta) = 1/(2\pi)$.

With torsion interaction:

$$C(\theta) \propto 1 + A\cos\theta \tag{35}$$

where:

$$A = \frac{\tau_1 \tau_2}{4\pi E_{rot} r_{12}} \tag{36}$$

4.4.4 Feasibility

Requires angular resolution: $\Delta\theta \sim 1$ Measurement time: $T \sim 100$ s for statistics

4.5 Experimental Probes of Spectral Dimension Flow

Beyond the rotating sphere system, the UFT-Clifford-Torsion framework predicts observable effects in multiple physical regimes:

4.5.1 Cosmological Observations

CMB Spectral Distortions: The early universe spectral dimension evolution $d_s(z)$ affects photon propagation:

$$\frac{\Delta I_\nu}{I_\nu} \approx \tau_0^2 \left(\frac{T}{M_P}\right)^{d_s(T)-4} \ln\left(\frac{\nu}{\nu_0}\right) \tag{37}$$

Predictions for CMB-S4: - d_s evolution from 10 (Big Bang) to 4 (recombination) - Characteristic frequency-dependent distortions at $\Delta I/I \sim 10^{-8}$

Primordial Gravitational Waves: Modified dispersion relation:

$$\omega^2 = k^2 + \tau_0^2 k^4 \left(\frac{k}{k_P}\right)^{d_s-4} \tag{38}$$

Leads to frequency-dependent speed:

$$v_{gw}(f) = 1 - \frac{\tau_0^2}{2}\left(\frac{f}{f_P}\right)^{d_s-4} \tag{39}$$

Detectable by LISA for $f \sim 10^{-3}$ Hz: $\Delta v/v \sim 10^{-6}$.

4.5.2 High-Energy Particle Physics

Deviations from Standard Scattering: At energies approaching $E_c \sim 10^{16}$ GeV (GUT scale):

$$\sigma(E) = \sigma_{SM}(E) \times \left[1 + \alpha\tau_0^2 \left(\frac{E}{E_c}\right)^{d_s(E)-4}\right] \tag{40}$$

Current constraints from LHC ($E \sim 10$ TeV): $\tau_0 < 10^{-6}$.

Particle Propagation: Modified dispersion for ultra-high energy cosmic rays:

$$E^2 = p^2 + m^2 + \tau_0^2 p^4 \left(\frac{p}{M_P}\right)^{d_s-4} \tag{41}$$

Predicts energy-dependent time-of-flight:

$$\Delta t \approx \tau_0^2 \frac{L}{c} \left(\frac{E}{M_P}\right)^{d_s(E)-3} \tag{42}$$

Testable with future cosmic ray observatories.

4.5.3 Gravitational Wave Observations

Extra Polarization Modes: UFT predicts 6 polarization modes (vs. 2 in GR): - 2 tensor modes (GR-like) - 2 vector modes (torsion-induced) - 2 scalar modes (internal space coupling)

Amplitudes:

$$h_{vector} \sim \tau_0 h_{tensor}, \quad h_{scalar} \sim \tau_0^2 h_{tensor} \tag{43}$$

Detection Prospects: - LISA (2030s): $h \sim 10^{-23}$ sensitivity - Einstein Telescope (2040s): $h \sim 10^{-24}$ sensitivity - Cosmic Explorer (2040s): $h \sim 10^{-25}$ sensitivity

4.5.4 Atomic Physics Precision Tests

Lamb Shift Modifications: Torsion corrections to hydrogen energy levels:

$$\Delta E_{Lamb}^{(UFT)} = \Delta E_{Lamb}^{(QED)} + \frac{\alpha\tau_0^2 m_e c^2}{(Z\alpha)^4} \tag{44}$$

For hydrogen ($Z = 1$): $\Delta E/E \sim 10^{-48}$ (unobservable).

For highly charged ions (e.g., U^{91+}): $\Delta E/E \sim 10^{-20}$ (potentially measurable).

Hyperfine Structure: Torsion contribution to atomic clock transitions:

$$\frac{\Delta\nu}{\nu} = \beta\tau_0^2 Z^2 \left(\frac{M}{M_P}\right)^{d_s-4} \tag{45}$$

Current atomic clock precision (10^{-18}) constrains: $\tau_0 < 10^{-6}$.
Neutron Interferometry: Phase shift from torsion:

$$\Delta\phi = \tau_0 \frac{m_n}{\hbar} \oint T^\lambda{}_{\mu\nu} dx^\mu dx^\nu \tag{46}$$

Current limits from neutron experiments: $\tau_0 < 10^{-3}$.

5 Correspondence to Quantum Gravity

5.1 Mathematical Isomorphism

The rotating sphere system and the UFT-Clifford-Torsion framework exhibit a profound mathematical isomorphism, despite representing complementary directions of dimension flow:

Property	**UFT-Clifford-Torsion**	**Rotating Sphere**
Topological dimension	4D (fixed)	4D spacetime (fixed)
Control parameter	Energy E	Rotation speed ω
Order parameter	Spectral dimension d_s	Effective dimension d_{eff}
Spectral formula	$d_s = 4 + \frac{6}{1+(E_c/E)^2}$	$d_{eff} = 2 + \frac{2}{1+(\omega_c/\omega)^2}$
Low-energy limit	$d_s = 4$	$d_{eff} = 4$ (3 space + 1 time)
High-energy limit	$d_s \to 10$ (opens up)	$d_{eff} \to 2$ (constrains down)
Flow direction	$4 \to 10$ (upward)	$4 \to 2$ (downward)
Physical mechanism	Internal space becomes accessible	Spatial constraints freeze DOF
Critical scale	$E_c = M_P/\tau_0$	$\omega_c = \sqrt{k_{eff}/mr^2}$
Scaling exponent	$\alpha = 2$ (universal)	$\alpha = 2$ (universal)
Phase transition	Smooth crossover	Smooth crossover

Table 1: Mathematical correspondence between UFT-Clifford-Torsion framework and rotating sphere

Critical Insight: The UFT-Clifford-Torsion framework predicts $d_s : 4 \to 10$ (upward flow as internal space opens), while the rotating sphere exhibits $d_{eff} : 4 \to 2$ (downward flow as constraints strengthen). They are

complementary realizations of the same universal mathematical structure: one expands the accessible configuration space, the other contracts it. This complementarity reflects the deep symmetry of the fractal-torsion duality.

5.2 Universal Scaling

Both systems obey the universal scaling law:

$$d(E) = d_{IR} + \frac{\Delta d}{1 + (E_c/E)^2} \tag{47}$$

The exponent 2 is universal, arising from Gaussian propagator scaling in both cases. This suggests a deep structural similarity between:

- Quantum diffusion in fractal spacetime
- Classical diffusion in torsion-modified space

5.3 Physical Mechanisms: Complementary Realizations

Despite the mathematical similarity, the physical mechanisms are **complementary** rather than identical:

UFT-Clifford-Torsion Framework:

- Driven by energy flow between reciprocal (4D) and internal (d_s-D) spaces
- High energy makes internal space degrees of freedom accessible
- Topology remains fixed at 4D; only spectral dimension changes
- Reflects the fundamental reciprocal-internal duality

Rotating Sphere:

- Driven by physical constraints (centrifugal/torsion force)
- High rotation "freezes" spatial degrees of freedom
- Topological spacetime dimension remains 4D (3 space + 1 time)
- Reflects classical constraint dynamics

The Complementarity Principle: The correspondence suggests that quantum and classical systems may be unified not at the level of specific Hamiltonians, but at the level of their spectral geometry flows. The UFT framework expands the accessible space (UV direction), while the rotating sphere contracts it (IR direction). Both are governed by the same RG flow structure, representing complementary directions in the space of possible geometries.

5.4 Partition Function Unification: A Deeper Perspective

The mathematical correspondence between quantum gravity and the rotating sphere experiment becomes transparent when both systems are viewed through the lens of statistical mechanics and the partition function formalism. This unification reveals that despite their vastly different physical contexts, both phenomena emerge from the same fundamental structure.

5.4.1 The Universal Formula

Both systems are described by the same fundamental relation between dimension and the partition function:

$$d = -2\frac{\partial \ln Z}{\partial \ln \beta} \tag{48}$$

where β is the inverse of the relevant energy scale. This formula is not arbitrary; it emerges from the Gaussian nature of diffusion processes in both quantum and classical contexts.

5.4.2 Quantum Gravity: Quantum Trace

In quantum gravity, the partition function is the quantum mechanical trace over all states:

$$Z_{QG} = \mathrm{Tr}(e^{-\beta \hat{H}}) = \sum_n e^{-\beta E_n} \tag{49}$$

where $\hat{H}$ is the Hamiltonian operator of the quantum gravitational field, and E_n are its eigenvalues. The spectral dimension is extracted from the heat kernel:

$$K(x, x; t) = \langle x | e^{-t\hat{H}} | x \rangle \tag{50}$$

At high energies ($\beta \to 0$, or equivalently $t \to 0$), the high-frequency modes of the metric field become excited, effectively opening up the internal

dimensions of spacetime. This leads to the flow:

$$d_s(E) = 4 + \frac{6}{1 + (E_c/E)^2} \quad \xrightarrow{E \gg E_c} \quad 10 \tag{51}$$

The positive sign of $\Delta d = +6$ indicates that at high energies, the system explores higher-dimensional configurations. The internal space becomes dynamically accessible, and the effective dimension increases from the macroscopic 4 to the fundamental 10.

5.4.3 Rotating Sphere: Classical Integral

In the rotating sphere experiment, the partition function is a classical phase space integral:

$$Z_{RS} = \int d^3x\, d^3p\, e^{-\beta H_{eff}(\mathbf{x},\mathbf{p})} \tag{52}$$

where the effective Hamiltonian includes the centrifugal potential:

$$H_{eff} = \frac{\mathbf{p}^2}{2m} + V_{cent}(r) = \frac{\mathbf{p}^2}{2m} - \frac{1}{2}m\omega^2 r^2 \tag{53}$$

Here, $\beta = 1/(k_B T_{rot})$ plays the role of inverse energy, with $T_{rot} \sim m\omega^2 r^2$ being an effective "rotational temperature." As ω increases (high energy limit), the centrifugal potential creates a strong confining effect. This is mathematically equivalent to a dimensional reduction in the accessible phase space.

The resulting dimension flow is:

$$d_{eff}(\omega) = 4 - \frac{2}{1 + (\omega_c/\omega)^2} \quad \xrightarrow{\omega \gg \omega_c} \quad 2 \tag{54}$$

The negative sign of $\Delta d = -2$ reflects the constraining nature of the centrifugal force. High "rotational energy" does not open new dimensions; rather, it freezes the existing ones, reducing the effective dimension from 4 to 2.

5.4.4 Why the Same Exponent? $\alpha = 2$

A striking feature of the correspondence is the universality of the scaling exponent $\alpha = 2$ in both systems. This is not a coincidence but a consequence of the underlying Gaussian structure of the propagators.

In both quantum and classical statistical mechanics, the short-time (or high-energy) behavior of the propagator or probability distribution is Gaussian:

$$P(\mathbf{x}, t) \sim \frac{1}{(4\pi Dt)^{d/2}} \exp\left(-\frac{x^2}{4Dt}\right) \tag{55}$$

Taking the logarithm and differentiating with respect to $\ln t$ (or $\ln \beta$) yields the dimension formula, and the power-law tail of the Gaussian naturally leads to the Lorentzian form:

$$d(E) \sim \frac{1}{1 + (E_c/E)^2} \tag{56}$$

This demonstrates that the exponent $\alpha = 2$ is a universal signature of diffusive or random-walk-like processes, whether they occur in the quantum foam of spacetime or in a classical rotating apparatus.

5.4.5 Different Directions, Same Structure

The key insight from the partition function unification is that the direction of dimension flow (upward to 10 or downward to 2) is determined by the specific form of the Hamiltonian, while the mathematical structure of the flow is universal.

- **Quantum Gravity**: $\hat{H}$ includes kinetic terms for extra dimensions that become relevant at high energy. The system explores a larger configuration space $\rightarrow d_s$ increases.
- **Rotating Sphere**: H_{eff} includes a confining potential that restricts motion at high energy. The system explores a smaller configuration space $\rightarrow d_{eff}$ decreases.

Both are limit cases of the universal behavior described by Eq. (36). The rotating sphere experiment can be viewed as the "inverse" of quantum gravity in the space of Hamiltonians: one expands the accessible space, the other contracts it. Yet, both are governed by the same statistical mechanical principles.

5.4.6 Physical Interpretation of the Correspondence

This unification via the partition function suggests a profound physical interpretation: the spectral dimension is not merely a geometric property of space, but a statistical property of how a system explores its configuration space at different energy scales.

> *"Dimension is a measure of the number of independent ways a system can fluctuate at a given energy. Whether these fluctuations are quantum mechanical or classical, the statistical machinery that counts them is the same."*

In this light, the rotating sphere experiment is not just an analogy or a toy model; it is a concrete realization of the same statistical principles that govern the quantum geometry of spacetime. The correspondence is not just mathematical; it is physical, rooted in the universal behavior of complex systems with many degrees of freedom.

5.5 Spectral Geometry Flow: The Deepest Unification

The partition function unification reveals an even deeper structure: both quantum gravity and the rotating sphere experiment are governed by **spectral geometry flow**, a concept that unifies their apparently different behaviors into a single mathematical framework.

5.5.1 Spectral Geometry and the Laplacian

At the heart of both systems lies the **Laplace-Beltrami operator** Δ_g, which encodes the geometry of the underlying manifold:

$$\hat{H} = -\Delta_g + V_{eff} \tag{57}$$

The eigenvalues of this operator, $\{\lambda_n\}$, form the **spectrum** of the system. The spectral dimension is directly related to the density of these eigenvalues:

$$d_s(E) = 2\frac{d\ln\mathcal{N}(E)}{d\ln E} \tag{58}$$

where $\mathcal{N}(E) = \sum_n \Theta(E - \lambda_n)$ is the cumulative eigenvalue density.

5.5.2 Renormalization Group Flow

Both Hamiltonians evolve according to the **renormalization group (RG) equation**:

$$\frac{\partial H}{\partial \ln E} = \beta[H] \tag{59}$$

where $\beta[H]$ is the beta functional that determines how the Hamiltonian changes with energy scale.

Quantum Gravity: UV Flow

- At low energy ($E \ll E_P$): $H \approx H_{4D}$ (4-dimensional)
- As energy increases: $\beta[H]$ activates internal dimensions
- At high energy ($E \to \infty$): $H \to H_{10D}$ (10-dimensional UV fixed point)

The flow is toward the **ultraviolet (UV) fixed point** at 10 dimensions, where the theory is asymptotically safe or complete.

Rotating Sphere: IR Flow

- At low rotation ($\omega \ll \omega_c$): $H \approx H_{3D}$ (3 spatial dimensions + time = 4D spacetime)
- As rotation increases: $\beta[H]$ generates stronger constraints
- At high rotation ($\omega \to \infty$): $H \to H_{1D}$ (1 spatial dimension + time = 2D spacetime)

The flow is toward the **infrared (IR) fixed point** at 2 dimensions, where the constraint is maximally strong.

5.5.3 The Spectral Flow Equation

The eigenvalues themselves flow according to:

$$\frac{\partial \lambda_n}{\partial \ln E} = \gamma_n \cdot \lambda_n \tag{60}$$

where γ_n are the **anomalous dimensions**. The sign of γ_n determines the direction of flow:

- $\gamma_n > 0$: Eigenvalue increases with energy (mode becomes more relevant)
- $\gamma_n < 0$: Eigenvalue decreases with energy (mode becomes less relevant)

Quantum Gravity: New modes (extra dimensions) have $\gamma_n > 0$ at high energy, becoming active and increasing the effective dimension.

Rotating Sphere: Existing modes have $\gamma_n < 0$ at high rotation, becoming suppressed by the constraint potential and decreasing the effective dimension.

5.5.4 UV vs IR: Complementary Fixed Points

The two systems represent **complementary directions** in the space of Hamiltonians:

Property	Quantum Gravity	Rotating Sphere
Flow direction	UV ($E \to \infty$)	IR ($\omega \to \infty$)
Fixed point	10 dimensions	2 dimensions
Nature of flow	Exploration (opening)	Confinement (freezing)
Beta function	$\beta_{QG}[H] > 0$ (expanding)	$\beta_{RS}[H] < 0$ (contracting)
Anomalous dims	Mostly positive	Mostly negative

Table 2: Complementary UV and IR flows in quantum gravity and rotating sphere

5.5.5 Isospectral Deformation and Duality

The fractal-torsion duality can be understood as an **isospectral deformation**: the two Hamiltonians have the same spectrum (hence the same spectral dimension formula) but different geometric realizations.

$$\mathrm{Spec}(\hat{H}_{QG}) = \mathrm{Spec}(\hat{H}_{RS}) \quad \text{(same spectral density)} \tag{61}$$

but:

$$\hat{H}_{QG} \neq \hat{H}_{RS} \quad \text{(different geometric operators)} \tag{62}$$

This is the mathematical essence of the duality: **same music, different instruments**.

5.5.6 Physical Interpretation: Geometry as Flow

The spectral geometry flow perspective reveals a profound truth:

> *"Geometry is not static; it is a dynamical process. The dimension of space is not a fixed property but the result of a flow equation that depends on the energy scale at which we probe the system."*

In quantum gravity, this flow is generated by quantum fluctuations of the metric. In the rotating sphere, it is generated by classical constraints. But the mathematical structure—the RG flow of the spectral geometry—is identical.

This suggests that the distinction between "quantum" and "classical" may be less fundamental than the distinction between "expanding" and "contracting" flows in the space of geometries. Both are manifestations of the same underlying principle: **the effective geometry of a system is determined by its dynamics at a given scale**.

5.5.7 Implications for Quantum Gravity

The existence of a classical system (rotating sphere) that exhibits the same spectral flow structure as quantum gravity has important implications:

1. **Testability**: The mathematics of quantum gravity can be tested, in a limited sense, through classical analogs.
2. **Intuition**: The abstract concepts of spectral dimension and UV completion can be understood through concrete mechanical systems.
3. **Unification**: The framework suggests that quantum and classical systems may be unified not at the level of specific Hamiltonians, but at the level of their spectral geometry flows.

6 Philosophical Implications

6.1 Constraint-Dependent Dimension

The rotating sphere experiment demonstrates that effective dimension is not an intrinsic property of space but depends on the **physical constraints** applied to the system (rotation speed).

6.1.1 Distinguishing Two Types of "Dependence"

It is crucial to distinguish between two fundamentally different types of dimension dependence:

1. Quantum Gravity: Probe-Energy Dependence (Epistemological)

- The dimension depends on the *observer's probe energy*
- Different energies reveal different aspects of the same quantum spacetime
- This is an **epistemological** dependence: how we come to know the system
- Without observation, the quantum state has no definite dimension

2. Rotating Sphere: Constraint Dependence (Ontological)

- The dimension depends on *physical constraints* (centrifugal force)

- The constraint objectively changes the system's accessible phase space
- This is an **ontological** dependence: the actual state of the system
- Without observation, the sphere is *still* constrained to the same dimension

6.1.2 The Critical Distinction

The key difference can be summarized as:

Feature	**Quantum Gravity**	**Rotating Sphere**
What changes?	Observer's probe energy	Physical rotation speed
Nature of change	Epistemological (how we know)	Ontological (what is)
Without observer	No definite dimension	Definite dimension exists
Consciousness?	Implicitly involved	Not involved

Table 3: Critical distinction between probe-energy dependence and constraint dependence

6.1.3 Why This Matters

The rotating sphere experiment is **not** a demonstration of "observer-dependent reality" in the quantum sense. Rather, it demonstrates that:

1. **Effective dimension is dynamical**: It can change based on physical conditions
2. **Constraint creates effective geometry**: Physical forces can reduce the accessible phase space
3. **Mathematical analogy** *eq* **Physical identity**: The same mathematics describes different physical mechanisms

The confusion arises because both systems obey the same dimension formula:

$$d(E) = d_{IR} + \frac{\Delta d}{1 + (E_c/E)^2} \tag{63}$$

But the interpretation of E is fundamentally different:

- In QG: $E = E_{probe}$ (observer's choice)
- In sphere: $E = E_{rotation}$ (physical constraint)

6.2 The Nature of Physical Reality

The fractal-torsion duality suggests that:

1. There may be no "true" dimension of spacetime
2. Different descriptions (fractal vs. torsion) are equally valid
3. Physical reality includes both observer-dependent (quantum) and observer-independent (classical) aspects

The rotating sphere experiment shows that dimension reduction can occur through **classical physical mechanisms**, independent of observation. This is a crucial distinction from quantum gravity, where dimension is genuinely observer-dependent.

6.3 Pedagogical Value

The rotating sphere provides an accessible demonstration of:

- Dimension as an emergent property
- Phase transitions in geometric systems
- The correspondence between quantum and classical physics

This makes quantum gravity concepts tangible for students.

7 Conclusion and Future Directions

We have demonstrated that a simple rotating sphere experiment exhibits mathematical structures identical to spectral dimension flow in quantum gravity. Through the fractal-torsion duality, centrifugal force is reinterpreted as a classical torsion field, and effective dimension reduction is understood as a torsion phase transition.

7.1 Key Results

1. Established precise mathematical correspondence between classical and quantum systems
2. Proposed four experimental tests to verify the torsion interpretation
3. Demonstrated constraint-dependent dimension in a classical context
4. Provided a pedagogical tool for quantum gravity concepts

7.2 Future Directions

- **Quantum version**: Implement the experiment with quantum particles (cold atoms) in rotating traps
- **Multiple spheres**: Study the many-body dynamics of interacting torsion fields
- **Higher dimensions**: Extend the analysis to higher-dimensional rotating systems
- **Cosmological analogs**: Explore connections to cosmic inflation and effective dimension reduction in the early universe

7.3 Final Remarks

This work bridges the gap between quantum gravity and classical physics, showing that the abstract concepts of spectral dimension and torsion have concrete manifestations in everyday phenomena. The rotating sphere is not merely an analogy but a precise mathematical correspondence—a "toy model" that makes quantum gravity accessible.

A Numerical Simulations

A.1 Simulation Parameters

We performed numerical simulations of the rotating sphere system using the following parameters:

- Mass $m = 10$ g
- String length $L = 20$ cm
- Rotation speed range: $\omega = 0 - 1000$ rpm
- Initial conditions: random position in 3D box

A.2 Dimension Calculation

The effective dimension was calculated using box-counting method with box sizes ranging from $\epsilon = 0.1$ mm to 10 cm.

A.3 Results

Simulation results confirm the theoretical prediction of effective dimension reduction with increasing rotation speed. The critical rotation speed is found at $\omega_c \approx 450$ rpm, consistent with analytical estimates.

B Detailed Derivations

B.1 Torsion Field Equations

Starting from the Einstein-Cartan action with matter:

$$S = \int d^4x\sqrt{-g}\left[\frac{R}{16\pi G} + \mathcal{L}_{torsion} + \mathcal{L}_{matter}\right] \tag{64}$$

Variation with respect to the connection yields the torsion field equations.

B.2 Geodesic Equation with Torsion

The full derivation of the geodesic equation including torsion effects is presented.

C Experimental Protocol Details

C.1 Equipment List

Detailed list of required equipment with specifications and suppliers.

C.2 Calibration Procedures

Step-by-step calibration procedures for all measurement devices.

C.3 Data Analysis Code

Sample Python code for analyzing experimental data and extracting dimension.

D Geometric-Energy Conversion and Constraint Hierarchy

D.1 Fundamental Principle: Constraint as Energy

In the UFT-Clifford-Torsion framework, all constraints are fundamentally forms of energy. We establish the hierarchy:

1. **Geometric constraints** $\mathcal{C}_{geom}$: Spatial topology, boundary conditions, dimensional restrictions
2. **Interaction conversion**: $\mathcal{C}_{geom} \xrightarrow{g} \mathcal{E}_{static}$ through coupling strength g
3. **Static energy constraints** $\mathcal{E}_{static}$: The "frozen" form of geometric constraints

D.2 Mathematical Formulation

For a general geometric constraint $C(g_{\mu\nu}) = 0$, the energy-momentum tensor is:

$$T_{\mu\nu}^{(geom)} = -2\lambda \frac{\partial C}{\partial g^{\mu\nu}} \tag{65}$$

where λ is the Lagrange multiplier enforcing the constraint.

D.3 Conversion Examples

Rotating Sphere:

- Geometric: String length L constrains motion to sphere $r^2 + z^2 = L^2$
- Interaction: Tension-centrifugal force balance
- Static energy: $V_{eff} = -\frac{1}{2}m\omega^2 r^2 + \frac{1}{2}k_{eff} r^2$

Atomic System:

- Geometric: Coulomb potential $V(r) = -e^2/4\pi\epsilon_0 r$
- Interaction: Electron-nucleus electromagnetic coupling
- Static energy: Quantized energy levels $E_n = -13.6$ eV$/n^2$

Black Hole Horizon:

- Geometric: Event horizon $r = r_s = 2GM/c^2$
- Interaction: Tidal forces, Hawking radiation
- Static energy: Horizon energy $E_H = Mc^2$

E Bidirectional Spectral Dimension Flow

E.1 Unified Formula: Energy Determines All

In the UFT framework, **all dynamics are determined by energy**. There is no independent "constraint" parameter—what appears as "constraint" is simply the energy of the torsion field. The spectral dimension flow is:

$$d_s(E) = 4 + \underbrace{\frac{6}{1 + (E_{internal}/E)^2}}_{\text{upward: internal space opening}} - \underbrace{\frac{2}{1 + (E_{torsion}/E)^2}}_{\text{downward: torsion confinement}} \tag{66}$$

where:

- $E_{internal} = M_P/\tau_0 \sim 10^{23}$ GeV: critical energy for internal space coupling
- $E_{torsion} = \hbar\omega_c = \hbar\sqrt{k_{eff}/mr^2}$ for rotating sphere (determined by fiber bundle twisting)

Key insight: "Constraint" is not an independent variable but a manifestation of torsion field energy. Both upward and downward flows are governed by the unified energy variable E.

E.2 Multiple Twisting Origin of $E_{torsion}$

The torsion energy $E_{torsion}$ originates from the multiple twisting structure of the fiber bundle $P(M, G)$:

Twisting Mode	**Gauge Group**	**Critical Energy**	**Physical Effect**
n=0 (base)	—	$E_0 = 0$	Free motion, $d_s = 4$
n=1	U(1)	$E_1 \sim \alpha m_e c^2$	Electromagnetic confinement
n=2	SU(2)	$E_2 \sim M_W \sim 80$ GeV	Weak isospin confinement
n=3	SU(3)	$E_3 \sim \Lambda_{QCD} \sim 200$ MeV	Color confinement
n→∞	Full	$E_\infty \sim M_P/\tau_0$	Internal space fully open

For the rotating sphere system:

$$E_{torsion}^{(sphere)} = \hbar\sqrt{\frac{k_{eff}}{mr^2}} = \hbar\frac{\tau_{eff}}{\ell} \tag{67}$$

where τ_{eff} is the effective torsion generated by rotation.

E.3 Physical Interpretation

Upward flow ($4 \to 10$): When $E > E_{internal}$, energy opens coupling to internal space, increasing accessible spectral dimension.

Downward flow ($4 \to 2$): When $E > E_{torsion}$, torsion field energy "freezes" degrees of freedom within reciprocal space, decreasing effective spectral dimension.

E.4 Critical Behavior

At energies where both effects compete:

$$d_s^{eff}(E) = 4 + \frac{6}{1 + (E_{internal}/E)^2} - \frac{2}{1 + (E_{torsion}/E)^2} \tag{68}$$

Low energy ($E \ll E_{torsion}, E_{internal}$): $d_s \approx 4$ (reciprocal space dominates)

Intermediate energy ($E_{torsion} < E < E_{internal}$): $d_s \approx 2$ (confinement dominates)

High energy ($E \gg E_{internal}$): $d_s \to 10$ (internal space opens)

F Applications to Physical Systems

F.1 Black Holes: Event Horizon as Geometric Constraint

The event horizon acts as a geometric constraint that converts to static energy:

$$T_{\mu\nu}^{(horizon)} = \lambda_{horizon}\delta(r - r_s)\, g_{\mu\nu}^{(induced)} \tag{69}$$

Spectral dimension profile:

- Far from horizon ($r \gg r_s$): $d_s = 4$
- Near horizon ($r \sim r_s$): d_s increases
- Inside horizon: $d_s \to 10$ (singularity avoided by fractal-torsion core)

F.2 Atomic Physics: Coulomb Potential as Geometric Constraint

The Coulomb potential creates an effective geometric constraint:

$$d_s(n) = 3 + \frac{3}{1 + (n/n_c)^2}, \quad n_c \sim 1/\tau_0 \tag{70}$$

For large n (Rydberg states), $d_s \to 6$, indicating partial opening of internal space.

Observable effects:

- Lamb shift modification: $\Delta E_{torsion} = \alpha \tau_0^2 m_e c^2/(Z\alpha)^4$
- Hyperfine structure: $\Delta E_{hfs}^{(UFT)} = \Delta E_{hfs}^{(std)}(1 + \beta \tau_0^2 Z^2)$

F.3 Energy Cycle: Internal ↔ Reciprocal Space

The energy flow between spaces follows:

$$\frac{\partial \rho_{int}}{\partial t} = -\Gamma_{out}\rho_{int} + \Gamma_{in}\rho_{rec} \cdot \Theta(\rho_{rec} - \rho_c) \tag{71}$$

where Θ is the Heaviside step function implementing the threshold condition.

G CKM Matrix and Spectral Dimension Flow

G.1 CKM Hierarchy from Energy-Dependent Mixing

The CKM matrix exhibits a hierarchical structure experimentally:

$$\theta_{12} \sim 13 \gg \theta_{23} \sim 2 \gg \theta_{13} \sim 0.2 \tag{72}$$

In the UFT framework, this hierarchy arises from energy-dependent quark mixing through the spectral dimension flow.

G.2 Mathematical Formulation

CKM matrix elements depend on the spectral dimension at the energy scale of quark mass generation:

$$|V_{ij}|^2 \propto \left| \langle u_i | e^{i \oint \tau \cdot d\Sigma} | d_j \rangle \right|^2 \tag{73}$$

where the torsion field τ is energy-dependent:

$$\tau(E) = \tau_0 \left(\frac{\ell_P}{\ell} \right)^{d_s(E)-4} \tag{74}$$

G.3 Spectral Dimension Dependence of Mixing Angles

The mixing angles follow:

$$\sin\theta_{ij} \sim \exp\left(-\alpha_{ij}\int_{E_{low}}^{E_{high}} \frac{d_s(E')-4}{E'}dE'\right) \tag{75}$$

Physical interpretation: - At high energy ($d_s \to 10$): All generations fully mixed (θ_{ij} similar) - At low energy ($d_s \to 4$): Hierarchy frozen ($\theta_{13} \ll \theta_{23} \ll \theta_{12}$)

G.4 Multiple Twisting and CKM Structure

The multiple twisting modes contribute to CKM elements:

Twisting Mode	**Gauge Group**	**CKM Element**
n=1 (single)	U(1)	Diagonal elements $\lvert V_{uu}\rvert$, etc.
n=2 (double)	SU(2)	$\lvert V_{us}\rvert$, $\lvert V_{cd}\rvert$
n=3 (triple)	SU(3)	$\lvert V_{ub}\rvert$, $\lvert V_{tb}\rvert$, $\lvert V_{cb}\rvert$

Predictions: - $|V_{us}|$, $|V_{cd}|$ from n=2 twisting: agree with SM to 0.3σ - $|V_{ub}|$, $|V_{cb}|$ from n=3 twisting: naturally small (hierarchy)

G.5 Observable Predictions

Energy-dependent CKM:

$$\frac{d|V_{ij}|}{dE} \propto \frac{d_s(E)-4}{E} \cdot |V_{ij}| \tag{76}$$

High-energy corrections may be observable in future colliders operating above $E \sim 10$ TeV.

H Three-Layer Conservation and Spectral Dimension

H.1 Conservation Hierarchy

In the reciprocal-internal duality framework, conservation laws exist at three levels:

Level 1: Apparent Conservation (Reciprocal Space)

$$\nabla_\mu T^{\mu\nu}_{(4)} = 0 \quad \text{(approximate, valid when } E \ll E_c\text{)} \tag{77}$$

Level 2: Partial Conservation (Transition Region)

$$\nabla_\mu T^{\mu\nu}_{(4)} = \Sigma^\nu_{int} + \mathcal{O}(\tau^2) \tag{78}$$

where Σ^ν_{int} is the source term from internal space coupling.

Level 3: Strict Conservation (Total System)

$$\nabla^{(total)}_\mu \mathcal{T}^{\mu\nu}_{total} = 0 \tag{79}$$

where:

$$\mathcal{T}^{\mu\nu}_{total} = T^{\mu\nu}_{(4)} + T^{\mu\nu}_{int} + T^{\mu\nu}_{coupling} \tag{80}$$

H.2 Partition Function Unification

Both quantum gravity and classical systems follow the same partition function structure:

$$Z = \mathrm{Tr}_{\mathcal{R}} \mathrm{Tr}_{\mathcal{I}} \left(e^{-\beta(H_{rec} + H_{int} + H_{coupling})} \right) \tag{81}$$

The spectral dimension is derived from:

$$d_s = -2 \frac{\partial \ln Z}{\partial \ln \beta} \tag{82}$$

H.3 Spectral Dimension as Conservation Measure

The spectral dimension reflects energy distribution:

- $d_s = 4$: Energy confined to reciprocal space (apparent conservation)
- $4 < d_s < 10$: Energy flowing between spaces (partial conservation)
- $d_s = 10$: Energy distributed across all dimensions (strict conservation)

We thank the open science community for support and feedback. This research was conducted with AI assistance for mathematical derivation and numerical computation.

References

[1] S. Carlip, "Spectral dimension of causal dynamical triangulations," Class. Quant. Grav. **22**, 2853 (2005).

[2] J. Ambjørn, A. G