

AE4889
Special Topics in Astrodynamics
MIMOSA – MIssion to Mars with a sOlar SAil

Daniel Pérez Trias – 6565158
Delft, June 10, 2026

github.com/dptrias/SpecialTopicsAstrodynamics

Collaborated with:

- Andreu Craus Vidal
- Rodrigo Duarte Soares
- Ismael Gallo López

Personal problem specification

Parameter	Value
Sun–Earth Lagrange point	L_2
Sun–Mars Lagrange point	L_2
Shift in Lagrange points	0.0005
Radius Martian parking orbit	306 666 km
Lightness number	0.025

Contents

List of Acronyms	3
WP1 Sun–Earth and Sun–Mars Lagrange points	1
WP2 Artificial Lagrange points	2
WP3 Stability of the (artificial) Lagrange points	5
WP4 Invariant manifolds and application to Mars exploration mission	6
References	7

List of Acronyms

AEP Artificial Equilibrium Point. [2](#)

CR3BP Circular Restricted Three-Body Problem. [1](#)

JPL Jet Propulsion Laboratory. [1](#)

WP1 Sun–Earth and Sun–Mars Lagrange points

1.1a The Lagrange points of the Sun–Earth and Sun–Mars systems are computed using Newton’s method over equation (1), which is derived from the [Circular Restricted Three-Body Problem \(CR3BP\)](#). The resulting dimensionless coordinates of the colinear Lagrange points (L_2) are presented in Table 1.

$$x - \frac{1 - \mu}{(x + \mu)^2} - \frac{\mu}{(x - 1 + \mu)^2} = 0 \quad (1)$$

Table 1: Dimensionless coordinates (x_i, y_i, z_i) of the Sun–Earth and Sun–Mars colinear Lagrange points in their respective synodic reference frames, obtained from the solution of the circular restricted three-body problem using a Newton method.

Lagrange point	x_i [LU]	y_i [LU]	z_i [LU]
Sun–Earth L_2	1.01009044	0	0
Sun–Mars L_2	1.00476311	0	0

1.1b For verification purposes, the dimensionless coordinates of the colinear Lagrange points are also extracted from the [Jet Propulsion Laboratory \(JPL\) Solar System Dynamics three-body periodic orbits tool](#), which are presented in Table 2. It can be observed that for the represented number of digits, the values are identical, with differences on the order of 10^{-9} LU, likely due to shared system parameters. *(50 words)*

Table 2: Dimensionless x coordinates of the Sun–Earth and Sun–Mars colinear Lagrange points in their respective synodic reference frames per JPL and their absolute differences w.r.t. the values obtained from the Newton method.

Lagrange point	x_i [LU]	Δx_i [LU]
Sun–Earth L_2	1.01009044	$4.21585256 \times 10^{-9}$
Sun–Mars L_2	1.00476311	$3.32491457 \times 10^{-9}$

1.2 A greater value of μ means the primary’s gravitational pull weakens ($1 - \mu$) while the secondary’s strengthens (μ). Since L_2 sits close to the secondary, this increase is more strongly felt there and must be offset by moving the point further from it. Increasing this distance has two compensating effects: it reduces the secondary’s pull more sharply than the primary’s (due to proximity), and it increases the centrifugal acceleration, which opposes both gravitational contributions. The net result is that a more massive secondary, as in the Sun–Earth system, pushes L_2 further from the planet than in the Sun–Mars case. *(99 words)*

WP2 Artificial Lagrange points

2.1a See Table 3.

Table 3: Required lightness number to achieve the specified shift in the Lagrange points.

Lagrange point	Shift in Lagrange point along the Sun-Planet line	Required lightness number, β
Sun–Earth L_2	0.0005	$4.76757628 \times 10^{-3}$
Sun–Mars L_2	0.0005	$5.06104189 \times 10^{-3}$

With respect to feasibility, NASA’s *ACS3* (currently in orbit), based on DCB deployable composite boom technology, has a lightness number $\beta = 0.025$ [1], an order of magnitude above the required values. This confirms that the artificial Lagrange point shifts studied are achievable with current solar sail technology. (47 words)

2.1b Since the required shift is along the Sun–planet line, and all forces at the collinear Lagrange points are aligned along this axis, the sail orientation needed to achieve the desired shift is likewise along this line. As the spacecraft moves sunward from L_2 , the gravitational pulls of both bodies strengthen while the centrifugal acceleration diminishes, requiring a compensating force directed away from the Sun. This uniquely determines the sail orientation, with the sail normal away from the Sun. (80 words)

Table 4: Required sail orientation to achieve the required shift in the Lagrange points.

Lagrange point	$n_{x,i}$	$n_{y,i}$	$n_{z,i}$
Sun–Earth L_2	1	0	0
Sun–Mars L_2	1	0	0




Figure 1: Schematic of all acceleration vectors acting on the spacecraft in the solar sail augmented CR3BP.

2.2 To verify the required lightness numbers calculations the colinear Artificial Equilibrium Point (AEP) specified in [2] is calculated using this work’s implementation and compared to the value reported in the reference, To verify the required lightness numbers calculations the colinear AEP specified in [2] is calculated using this work’s implementation and compared to the value reported in the reference, the considered case being the Sun–Earth system with a AEP located at $x_{\text{AEP}} = 0.983867$ LU. The resulting lightness number is $\beta = 0.0363$, which matches the value reported in [2], confirming the correctness of the implementation. (64 words)

2.3 The required area of the solar sail is derived from the definition of the lightness number:

$$\beta \frac{\mu_{\odot}}{r^2} = 2P \frac{A}{m} \rightarrow A = \frac{\beta \mu_{\odot} m}{2Pr^2} \quad (2)$$

where P is the solar radiation pressure at the location of the sail, which can be calculated from the solar constant, $S_{\odot}$ [3], and the speed of light, c , as follows:

$$P = \frac{S_{\odot}}{c} \left(\frac{\text{AU}}{r} \right)^2 \quad (3)$$

The resulting sizes required to achieve the specified shifts in the Lagrange points are shown in Table 5.

Table 5: Required solar-sail side-length to achieve the required shift in the Lagrange points.

Lagrange point	Required solar-sail side-length [m]
Sun–Earth L_2	17.6459
Sun–Mars L_2	18.1809

These sizes are at the upper limit of currently feasible solar sails. The maximum sail size of DCB-based designs in CubeSat form factor is 16.5 m, with larger platforms expected to be capable of deploying sails of up to 22.8 m (*27U HIPERSail*) [1].

2.4 To analyse the sensitivity of the required lightness number to the shift in the Lagrange point, a range of 21 shifts from 0.0005 LU to 0.001 LU is considered, and the corresponding lightness numbers are computed using the formula from Task 2.1a. The results are shown in Figure 2.




Figure 2: Sensitivity of the required lightness number, β , to the shift in the Lagrange point, Δx , for the Sun–Earth and Sun–Mars systems.

As the equilibrium position moves closer to the planet, the gravitational attraction from both bodies strengthens while the centrifugal acceleration weakens. To counterbalance these effects, the sail acceleration must increase, requiring a larger lightness number. The

required β is observed to follow a near-linear increase with decreasing distance from the Sun for both systems. (*54 words*)

WP3 Stability of the (artificial) Lagrange points

3.1a See Table 6.

Table 6: Eigenvalues of the classical and artificial Lagrange points.

Lagrange point	λ_1	λ_2	λ_3	λ_4
Sun–Earth Classical	−2.484281	−2.056992 <i>i</i>	+2.056992 <i>i</i>	+2.484281
Sun–Earth Artificial	−2.673872	−2.174388 <i>i</i>	+2.174388 <i>i</i>	+2.673872
Sun–Mars Classical	−2.496888	−2.064657 <i>i</i>	+2.064657 <i>i</i>	+2.496888
Sun–Mars Artificial	−2.931888	−2.335453 <i>i</i>	+2.335453 <i>i</i>	+2.931888

3.1b From the linearization of the problem around the equilibrium points it can be observed that all cases have a positive real eigenvalue, indicating that the nonlinear system is unstable around the equilibrium point [4]. All eigenvalues take similar values across the 4 cases, with slightly higher magnitudes in the Sun–Mars system, indicating faster local dynamics. The introduction of the solar sail also increases the magnitude of all eigenvalues slightly, but not enough to have a qualitative effect on the stability of the system. (82 words)

3.2 For verification purposes the eigenvalue calculation tool results are compared against values presented in **szebehely1967theory** for a value of $\mu = 3 \times 10^{-6}$. The resulting eigenvalues are shown in Table ??.

Table 7: Eigenvalues of the classical and artificial Lagrange points.

Lagrange point	λ_1	λ_2	λ_3	λ_4
Sun–Earth Classical	−2.484281	−2.056992 <i>i</i>	+2.056992 <i>i</i>	+2.484281
Sun–Earth Artificial	−2.673872	−2.174388 <i>i</i>	+2.174388 <i>i</i>	+2.673872
Sun–Mars Classical	−2.496888	−2.064657 <i>i</i>	+2.064657 <i>i</i>	+2.496888
Sun–Mars Artificial	−2.931888	−2.335453 <i>i</i>	+2.335453 <i>i</i>	+2.931888

WP4 Invariant manifolds and application to Mars exploration mission

Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem ipsum dolor sit amet, consectetur adipiscing elit. In hac habitasse platea dictumst. Integer tempus convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor. Proin tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula.

References

- [1] W. Wilkie, J. Fernandez, O. Stohlman, *et al.*, “An overview of the NASA Advanced Composite Solar Sail (ACS3) technology demonstration project,” Jan. 2021.
- [2] J. Heiligers, B. Diedrich, B. Derbes, and C. McInnes, “Sunjammer : Preliminary end-to-end mission design,” in *AIAA/AAS Astrodynamics Specialist Conference 2014*, USA, Aug. 2014. [Online]. Available: <https://strathprints.strath.ac.uk/49051/>.
- [3] G. Kopp and J. L. Lean, “A new, lower value of total solar irradiance: Evidence and climate significance,” *Geophysical Research Letters*, vol. 38, no. 1, DOI: <https://doi.org/10.1029/2010GL045777>.
- [4] F. Verhulst, *Nonlinear Differential Equations and Dynamical Systems* (Universitext), 2nd ed. Berlin, Heidelberg: Springer-Verlag, 1996, ISBN: 978-3-540-60934-6.