Logistic regression

David Puelz
The University of Austin

okcupid

Have an account? Sign in


DATING DESERVES BETTER

On OkCupid, you're more than just a photo. You have stories to tell, and passions to share, and things to talk about that are more interesting than the weather. Get noticed for who you are, not what you look like. Because you deserve what dating deserves: better.

JOIN okc

By clicking Join, you agree to our Terms of Service. Learn about how we process and use your data in our Privacy Policy and how we use cookies and similar technology in our Cookie Policy.

GET THE APP



The OkCupid data set

- The `okcupid` data set contains information on 59,946 profiles from users of the OkCupid online dating service.
- We have user age, height, sex, income, orientation, education, body type, ethnicity, and more.
- OkCupid has published their own analyses of this kind of data.

Our question: can we predict the sex of a user from their height?

What is wrong with this regression?

$$\widehat{\text{sex}} = \hat{\beta}_0 + \hat{\beta}_1 \cdot \text{height}$$

What is wrong with this regression?


$$\widehat{\text{sex}} = \hat{\beta}_0 + \hat{\beta}_1 \cdot \text{height}$$

The Y variable is **categorical** — two levels, male or female.

Ordinary linear regression is built for a quantitative Y. Nothing stops us from running it, but we should ask what it would even mean.

But what if we just do it anyway?


Code `male = 1`, `female = 0`, and fit a line:



A line is a spectacularly bad fit. And what would it mean to predict male $= 0.7$? Or 1.2?

First, clean the data

There are some impossible values for height:



Keep heights between 55 and 80 inches (4'7" and 6'8"). That drops only 120 of 59,946 cases.

```
my.profiles <- profiles %>% filter(height >= 55 & height <= 80)
```

The idea behind logistic regression

- Instead of predicting male/female, predict the **probability** that a user is male.
- One level is "success" ($Y = 1$), the other "failure" ($Y = 0$). Here male is success.
- Fit a curve that is always between 0 and 1.

To build that curve we need one piece of vocabulary first: **odds**.

Odds

- "Even (1/1) odds" $\Rightarrow$ probability of success is $1/2$

Odds

- "Even (1/1) odds" $\Rightarrow$ probability of success is $1/2$
- "2/1 odds" $\Rightarrow$ probability is $2/3$

Odds

- "Even (1/1) odds" $\Rightarrow$ probability of success is $1/2$
- "2/1 odds" $\Rightarrow$ probability is $2/3$
- In general, the odds of something with probability p are

$$\text{odds} = \frac{p}{1-p}$$

Odds: an example

During March Madness, a \$100 bet on San Diego State paid \$360 in winnings — odds to win of $1/3.6$.

What probability does that imply?

Odds: an example

During March Madness, a \$100 bet on San Diego State paid \$360 in winnings — odds to win of $1/3.6$.

What probability does that imply?

$$\frac{p}{1-p} = \frac{1}{3.6} \quad \Longrightarrow \quad p = \frac{1}{4.6} \approx 0.22$$

The logistic regression model

Logistic regression models the **log odds** of success as a linear function of X:


$$\log\left(\frac{p}{1-p}\right) = \beta_0 + \beta_1 X$$

The left side runs over all of $(-\infty, \infty)$, so a straight line is a sensible thing to put on the right.

Solving back for p gives an S-shaped curve, bounded between 0 and 1:

$$p = \frac{e^{\beta_0+\beta_1 X}}{1 + e^{\beta_0+\beta_1 X}}$$

The same model, two ways to look at it



Straight on the log-odds scale; squashed into $(0, 1)$ on the probability scale.

Let's fit it

```
model <- glm(male ~ height, data=my.profiles, family=binomial)
summary(model)


Call:
glm(formula = male ~ height, family = binomial, data = my.profiles)

Coefficients:
             Estimate Std. Error z value Pr(>|z|)
(Intercept) -44.44861    0.35751    -124   <2e-16 ***
height        0.66190    0.00529     125   <2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

How do we read the fit?

The regression says

$$\log(\text{odds}) = \log\left(\frac{P(\text{male})}{1 - P(\text{male})}\right) = -44.45 + 0.66 \cdot \text{height}.$$

How do we read the fit?

The regression says

$$\log(\text{odds}) = \log\left(\frac{P(\text{male})}{1-P(\text{male})}\right) = -44.45 + 0.66 \cdot \text{height}.$$

Solving for the probability:

$$\widehat{P}(\text{male}) = \frac{e^{-44.45+0.66 \cdot \text{height}}}{1+e^{-44.45+0.66 \cdot \text{height}}}$$


Making predictions

```
predict(model, data.frame(height=69), type="response")
```

```
     1
0.7725
```

Someone who is 5'9" has about a 77% chance of being male.

Visualizing the fitted model



The curve never leaves $[0, 1]$, and it bends where the data actually change over.

Interpreting the coefficients

$$\log\left(\frac{P(\text{male})}{1-P(\text{male})}\right) = -44.45 + 0.66 \cdot \text{height}$$

- At height $= 0$, the predicted log odds are -44.45, so the predicted probability is essentially 0.

Interpreting the coefficients

$$\log\left(\frac{P(\text{male})}{1-P(\text{male})}\right) = -44.45 + 0.66 \cdot \text{height}$$

- At height $= 0$, the predicted log odds are -44.45, so the predicted probability is essentially 0.
- Each extra inch of height adds 0.66 to the log odds.

Interpreting the coefficients

$$\log\left(\frac{P(\text{male})}{1-P(\text{male})}\right) = -44.45 + 0.66 \cdot \text{height}$$

- At height $= 0$, the predicted log odds are -44.45, so the predicted probability is essentially 0.
- Each extra inch of height adds 0.66 to the log odds.

A height of zero is nowhere near the data. The intercept anchors the line; it is not a claim about anybody.

Log odds are awkward. Odds are not.

Exponentiating both sides:

$$\widehat{\text{odds of male}} = e^{-44.45+0.66 \cdot \text{height}}$$

- One more inch **multiplies** the odds by $e^{0.6619} = 1.94$ — a 94% increase.

Log odds are awkward. Odds are not.

Exponentiating both sides:

$$\widehat{\text{odds of male}} = e^{-44.45+0.66 \cdot \text{height}}$$

- One more inch **multiplies** the odds by $e^{0.6619} = 1.94$ — a 94% increase.
- Two more inches multiply the odds by $1.94^2 = 3.76$ — a 276% increase.

In linear regression slopes add. In logistic regression they multiply.

How good is our model?

- The usual R^2 is not available for logistic regression.

How good is our model?

- The usual R^2 is not available for logistic regression.
- There are "pseudo-R^2" measures, but they are harder to interpret and easier to misuse.

How good is our model?

- The usual R^2 is not available for logistic regression.
- There are "pseudo-R^2" measures, but they are harder to interpret and easier to misuse.
- We will focus on something simple and concrete: **how many cases did we get right?**

Turning probabilities into predictions

Rule: predict male if $\widehat{P}(\text{male}) \geq 0.5$, otherwise female.

```
predicted.sex <- ifelse(predict(model, type="response") >= 0.5, "m", "f")
correct <- ifelse(predicted.sex == my.profiles$sex, 1, 0)
sum(correct) / nrow(my.profiles)

[1] 0.8309
```

83% of what?

Any accuracy number needs a baseline. What if we always guessed the majority?

```
xtabs(~ sex, data=my.profiles) %>% prop.table()

sex
     f      m
0.4027 0.5973
```

Always guessing male gets 60% right. Our model gets 83%.

That improvement is the **lift** the model buys us: from 60% to 83%.

The confusion matrix

Pick one category as **positive** (here, male) and the other as **negative** (female). Every case lands in one of four cells.

	Actual	
Predicted	Positive	Negative
Positive	TP	FP
Negative	FN	TN

FP = a woman predicted male. FN = a man predicted female.

The confusion matrix, for real

```
tab_conf <- xtabs(~ predicted.sex + my.profiles$sex)
tab_conf

             my.profiles$sex
predicted.sex     f     m
            f 19466  5494
            m  4623 30243
```

5,494 false negatives (men predicted female) and 4,623 false positives (women predicted male).

Careful reading the quadrants: the schematic above lists positive first, but R prints `f` first.

Rates from the confusion matrix

- **True positive rate** $= \mathrm{TP}/(\mathrm{TP}+\mathrm{FN})$ — of the actual positives, the share we called positive
- **True negative rate** $= \mathrm{TN}/(\mathrm{TN}+\mathrm{FP})$ — of the actual negatives, the share we called negative
- **False positive rate** $= \mathrm{FP}/(\mathrm{TN}+\mathrm{FP})$
- **False negative rate** $= \mathrm{FN}/(\mathrm{TP}+\mathrm{FN})$

Each rate is a column of the confusion matrix, normalized.

The same table, normalized

```
xtabs(~ predicted.sex + my.profiles$sex) %>% prop.table(2)

             my.profiles$sex
predicted.sex      f      m
            f 0.8081 0.1537
            m 0.1919 0.8463
```

`prop.table(2)` divides by the column totals, so each column is one actual group: we catch 81% of the women and 85% of the men.

Evaluating a classifier: three cautions

- The 0.5 threshold is a **choice**. Move it and you trade false positives against false negatives.
- Those two errors rarely cost the same. A missed loan default and a wrongly denied loan are not equivalent; neither are a missed tumor and a false alarm.
- Everything here is in-sample, and therefore optimistic. For an honest number, hold out a test set or cross-validate.

Adding a predictor

Just like linear regression: each coefficient is the effect on the log odds, holding the others constant.

```
xtabs(~ orientation, data=my.profiles)

orientation
bisexual       gay straight
    2763      5568    51495
```

`orientation` has three categories, so R will fit two dummies against a reference level.

The fit

```
model2 <- glm(male ~ height + orientation, data=my.profiles,
              family=binomial)
summary(model2)


Call:
glm(formula = male ~ height + orientation, family = binomial,
    data = my.profiles)

Coefficients:
                   Estimate Std. Error z value Pr(>|z|)
(Intercept)         -46.08076    0.37167  -124.0   <2e-16 ***
height                0.66535    0.00537   124.0   <2e-16 ***
orientationgay        2.09556    0.07209    29.1   <2e-16 ***
orientationstraight   1.39972    0.06068    23.1   <2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Interpreting the coefficients

$$\log\left(\frac{p}{1-p}\right) = -46.08 + 0.67 \cdot \text{height} + 2.10 \cdot \text{gay} + 1.40 \cdot \text{straight}$$

- The intercept is the log odds for a **bisexual** user of height 0 — the reference level.

Interpreting the coefficients

$$\log\left(\frac{p}{1-p}\right) = -46.08 + 0.67 \cdot \text{height} + 2.10 \cdot \text{gay} + 1.40 \cdot \text{straight}$$

- The intercept is the log odds for a **bisexual** user of height 0 — the reference level.
- Within a given orientation, each extra inch multiplies the odds by $e^{0.6653} = 1.95$.

Interpreting the coefficients

$$\log\left(\frac{p}{1-p}\right) = -46.08 + 0.67 \cdot \text{height} + 2.10 \cdot \text{gay} + 1.40 \cdot \text{straight}$$

- The intercept is the log odds for a **bisexual** user of height 0 — the reference level.
- Within a given orientation, each extra inch multiplies the odds by $e^{0.6653} = 1.95$.
- At the same height, gay vs. bisexual multiplies the odds by $e^{2.0956} = 8.13$; straight vs. bisexual, by $e^{1.3997} = 4.05$.


Why those dummies are so large

```
xtabs(~ sex + orientation, data=my.profiles) %>% prop.table(2)

   orientation
sex bisexual    gay straight
  f   0.7217 0.2848   0.3983
  m   0.2783 0.7152   0.6017
```

The bisexual profiles are 72% female; the gay profiles are 72% male. Orientation carries real information about sex in this data, quite apart from height.

Three curves, same slope



Adding a categorical predictor **shifts** the curve sideways. It does not change its steepness.

In linear regression, an interaction means the **slope** of X_1 depends on the value of X_2.

Exactly the same idea here: the effect of height on the odds is allowed to differ by orientation.

$$\texttt{male} \sim \texttt{height} * \texttt{orientation}$$

The interaction fit

```
int.model <- glm(male ~ height * orientation, data=my.profiles,
                 family=binomial)
summary(int.model)


Call:
glm(formula = male ~ height * orientation, family = binomial,
    data = my.profiles)


Coefficients:
                          Estimate Std. Error z value Pr(>|z|)
(Intercept)               -35.3027     1.4050  -25.13  < 2e-16 ***
height                      0.5076     0.0206   24.67  < 2e-16 ***
orientationgay             -6.2727     1.8365   -3.42  0.00064 ***
orientationstraight       -10.2887     1.4596   -7.05  1.8e-12 ***
height:orientationgay       0.1218     0.0271    4.49  7.1e-06 ***
height:orientationstraight  0.1712     0.0214    8.01  1.2e-15 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

The interaction model

$$\log\left(\frac{p}{1-p}\right) = -35.30 + 0.51 \cdot \text{height} \\ - 6.27 \cdot \text{gay} - 10.29 \cdot \text{straight} \\ + 0.12 \cdot \text{height} \cdot \text{gay} + 0.17 \cdot \text{height} \cdot \text{straight}$$

Each orientation now gets its own slope: 0.51 for bisexual, $0.51 + 0.12 = 0.63$ for gay, $0.51 + 0.17 = 0.68$ for straight.

Three curves, three slopes



Now the curves differ in steepness too: height is a sharper signal for straight users than for bisexual ones.

Three hypotheses worth testing

1. **Overall:** $\beta_1 = \beta_2 = \cdots = 0$. The model has no predictive power at all.

Three hypotheses worth testing

1. **Overall:** $\beta_1 = \beta_2 = \cdots = 0$. The model has no predictive power at all.
2. **One quantitative predictor:** $\beta_i = 0$. That X has no effect, holding the others constant.

Three hypotheses worth testing

1. **Overall:** $\beta_1 = \beta_2 = \cdots = 0$. The model has no predictive power at all.
2. **One quantitative predictor:** $\beta_i = 0$. That X has no effect, holding the others constant.
3. **One dummy:** $\beta_i = 0$. That category is no different from the reference level, holding the others constant.

Testing the overall null

The **likelihood ratio test** compares the full model against an intercept-only model.

For `male ~ height + orientation`, the test statistic is $\chi^2 = 36{,}932$ on 3 degrees of freedom, with $p < 0.001$.

With 59,826 profiles, this is not a close call — but note that with a sample this large, almost any non-zero effect will be "significant."

For one quantitative predictor (height) or one dummy (straight vs. bisexual), read the z value and p-value straight off the `summary` table.

	Estimate	Std. Error	z	$\Pr(>\|z\|)$
`height`	0.6653	0.0054	124.0	$< 2 \times 10^{-16}$
`orientationgay`	2.0956	0.0721	29.1	$< 2 \times 10^{-16}$
`orientationstraight`	1.3997	0.0607	23.1	$< 2 \times 10^{-16}$

Same logic as the t tests in linear regression; the reference distribution is normal, not t.

What else is logistic regression for?

- **Finance:** which customers are likely to default on a loan
- **Medicine:** whether a patient has a condition, given test results
- **Advertising:** who will respond to a campaign
- **Marketing:** who will purchase, subscribe, or sign up

Any time the answer is yes or no and you want a probability rather than a guess.

What to take away

- Model the **log odds** as linear; read the fit back as an S-curve in probability.
- Coefficients **multiply the odds**: e^{β} is the interpretable number.
- Evaluate with a **confusion matrix** against a baseline, not with R^2.
- The 0.5 threshold is yours to move, and the two kinds of error rarely cost the same.