
KEGG-mapped subsystems: median shrunk log2FC, DEGs/tested genes
(* DEG enrichment BH FDR < 0.05)
Mycolic Acid & FAS-II Envelope
+0.12
0/15 DEGs (0 up, 0 down)
+0.17
0/15 DEGs (0 up, 0 down)
Peptidoglycan Biosynthesis
+0.13
0/1 DEGs (0 up, 0 down)
+0.05
0/1 DEGs (0 up, 0 down)
Redox Homeostasis & Cofactors
+0.28
0/6 DEGs (0 up, 0 down)
+0.17
0/6 DEGs (0 up, 0 down)
Arginine & Branched-Chain Amino Acid Biosynthesis
+0.08
2/7 DEGs (2 up, 0 down)
+0.20
1/7 DEGs (1 up, 0 down)
Central Carbon & Glyoxylate Shunt
-0.04
2/7 DEGs (1 up, 1 down)
+0.01
1/7 DEGs (0 up, 1 down)
Type VII (ESX) Secretion
+0.17
5/27 DEGs (3 up, 2 down)
+0.35
7/27 DEGs (5 up, 2 down)
DosR Hypoxia & Dormancy
-0.97
2/4 DEGs (0 up, 2 down)
-2.02
2/4 DEGs (0 up, 2 down)
Cellular Respiration & Energy
+0.90
7/19 DEGs (7 up, 0 down)
+1.41*
10/19 DEGs (10 up, 0 down)
3D clinostat vs 1g
RPM 2.0 vs 1g
median shrunk log2FC
2.0
1.5
1.0
0.5
0.0
−0.5
−1.0
−1.5
−2.0