Multi-Crop Meta-Analysis of Spaceflight-Grown Leafy Greens on the ISS

Comparative Reanalysis of NASA OSDR OSD-745, OSD-655, and OSD-780 with Tabular Foundation Models

Author: Richard Barker

NASA GeneLab / OSDR

Prepared for:
Space Biology Community, NASA GeneLab/OSDR

Classification:
Scientific Analysis & Open Data | Q3 2026

CoSE Open Science & Space Biology Initiative
October 14, 2026

Contents

1 Introduction **1**

1.1 The Imperative of Space Agriculture . . . 1
1.2 The Veggie Plant Growth Facility . . . 1
1.3 The OSD-745, OSD-655, and OSD-780 Studies . . . 2
1.4 Motivation for Machine Learning Reanalysis . . . 2
1.5 Study Objectives and Research Questions . . . 3

2 Materials & Methods **5**

2.1 Data Source and Curation . . . 5
2.2 Plant Material and Growth Conditions . . . 5
2.3 Sample Collection and Analytical Methods . . . 6
2.3.1 Elemental Profiling (ICP-OES) . . . 6
2.3.2 Antioxidants and Phenolics . . . 6
2.3.3 Microbiological Plate Counts . . . 6
2.4 Machine Learning Pipeline . . . 6
2.4.1 Dimensionality Reduction . . . 6
2.4.2 Classification and Validation Strategies . . . 7
2.4.3 SHAP Feature Importance . . . 7
2.4.4 Tabular Foundation Model Benchmarking . . . 7
2.4.5 Statistical Testing . . . 8
2.5 Interactive Dashboard Architecture . . . 8

3 Results **9**

3.1 Data provenance and the 13 October 2026 correction . . . 9
3.2 Nutritional Profiling and Multi-Crop Composition . . . 10
3.3 Joint PCA Dimensionality Reduction . . . 10
3.4 Cross-Validation and Classification Benchmarks . . . 11
3.5 Feature Importance & SHAP Interpretation . . . 12
3.6 Regression: Predicting Biochemical Outputs . . . 12
3.7 Microbiological Food Safety Analysis . . . 14
3.8 Statistical Validation and Effect Sizes . . . 15

4 Discussion **17**
4.1 Nutritional Equivalence and Species Diversity 17
4.2 Universal Mechanistic Responses to Microgravity 17
4.3 The Power of Tabular Foundation Models (TabPFN) 17
4.4 Microbiological Safety and Crew Hygiene 18
4.5 Limitations of the Meta-Analysis 18
4.6 Future Directions 18

5 Conclusion **19**

List of Figures

3.1 Joint PCA of lettuce (OSD-745) and mizuna (OSD-655) on the 11 elemental and biochemical features with complete coverage ($N = 55$). PC1 separates the two crops. Lettuce samples group by mission rather than by flight or ground; mizuna flight samples tend to sit higher on PC2 than ground controls, with overlap. 10
3.2 Classification performance (accuracy, precision, recall and F1) of the Random Forest classifier and the benchmark model under LOMOCV on the joint dataset. The benchmark is the Gradient Boosting fallback, because TabPFN did not run without a license token. Neither model separates flight from ground. 11
3.3 Feature importance (mean decrease in impurity) for the joint flight-versus-ground Random Forest classifier. Potassium, sodium, sulfur and manganese rank highest. Because the classifier performs at chance, the ranking shows what the model attended to, not validated spaceflight markers. 13
3.4 Aerobic plate counts and yeast/mould counts on mizuna leaf tissue (OSD-780), flight versus ground control. Plotted as $\log_{10}(\mathrm{CFU/g} + 1)$: raw counts span zero to 1.7×10^7 CFU/g and over half the ground samples are exactly zero, so a linear axis and arithmetic means would both be misleading. 14

List of Tables

3.1 Joint statistical comparison of key nutritional markers between Flight (FLT) and Ground (GC) cohorts. 15

Chapter 1

Introduction

1.1 The Imperative of Space Agriculture

The future of human space exploration relies heavily on our ability to sustain astronaut health and psychological well-being during long-duration missions. Missions beyond Low Earth Orbit (LEO), such as lunar bases under the Artemis program or crewed missions to Mars, will stretch the limits of pre-packaged food systems. Pre-packaged diets, while calorically dense and microbiologically safe, undergo nutritional degradation over time [1]. Key vitamins, such as Vitamin C and B1, naturally decay, presenting significant risks of nutritional deficiencies for crews on multi-year deployments.

Furthermore, the psychological benefits of interacting with living plants—a phenomenon often referred to as "horticultural therapy"—have been widely documented among astronauts. The presence of fresh greenery provides sensory stimulation and a connection to Earth, breaking the monotony of an artificial, sterile spacecraft environment. Consequently, developing robust space agriculture systems is not merely a logistical necessity but a biological and psychological imperative [2].

1.2 The Veggie Plant Growth Facility

To address these challenges, NASA deployed the Vegetable Production System (Veggie) to the International Space Station (ISS). Veggie is a low-resource plant growth facility designed to produce fresh vegetables in microgravity. The hardware consists of a light cap fitted with red, blue, and green Light Emitting Diodes (LEDs), providing customizable photosynthetic and photomorphogenic lighting. Plants are grown in root mat "pillows"—Teflon-coated Kevlar bags filled with an arcillite-based substrate and controlled-release fertilizer. Water is introduced through a root mat, relying on capillary action to transport moisture to the plant roots in the absence of gravity-driven convection and drainage.

Despite its successes, growing plants in space presents profound environmental stressors. Microgravity alters fluid dynamics, often leading to hypoxia in the root zone. Elevated radiation, continuous ventilation (to prevent boundary layer suffocation), and unique microbial profiles fur-

ther complicate cultivation. Understanding how these factors influence the nutritional profile and yield of crops is a paramount objective of NASA's Space Biology program.

1.3 The OSD-745, OSD-655, and OSD-780 Studies

The Open Science Data Repository (OSDR) study OSD-745 encapsulates a comprehensive analysis of *Lactuca sativa* cv. 'Outredgeous' (red romaine lettuce) cultivated during three distinct ISS missions: VEG-01A, VEG-01B, and VEG-03A [3]. To expand space crop diversity, subsequent experiments under the VEG-04 program evaluated mizuna mustard (*Brassica rapa* var. *japonica*) during VEG-04A and VEG-04B. The chemical and elemental profiles for these Mizuna crops are curated under study OSD-655 [4], while their microbiological safety profile is detailed in study OSD-780.

In all of these missions, plants were grown on the ISS (Flight) and compared against parallel cohorts grown in environment-controlled chambers at the Kennedy Space Center (Ground Control). The harvested tissues were analyzed for a suite of metrics, including elemental composition (via ICP-OES), antioxidant capacity (ORAC), phenolic content, and anthocyanins. The original publications confirmed that both crops were microbiologically safe and nutritionally comparable to Earth-grown controls. Merging these datasets enables a true comparative meta-analysis across different space-grown crop species.

1.4 Motivation for Machine Learning Reanalysis

While the univariate statistical analyses performed in the original study established broad comparability, biological systems are inherently complex and multivariate. Elemental uptake and secondary metabolite synthesis do not occur in isolation; they represent an interconnected web of physiological responses. Single-element ANOVA or t-tests, though valuable, may obscure subtle, systemic shifts in the plant's nutritional network induced by the spaceflight environment.

Machine learning (ML) offers a powerful toolkit for uncovering these high-dimensional patterns. Techniques such as Random Forests [5] and Shapley Additive Explanations (SHAP) [6] allow us to interpret complex, non-linear relationships and identify the precise combinations of elements that differentiate space-grown from ground-grown crops. By reanalyzing the open-access OSD-745 dataset through an ML lens, we aim to extract deeper biological insights that can inform the design of next-generation life support systems.

Alert: Open Science Commitment

This manuscript, alongside its accompanying code and interactive dashboard, embraces the FAIR (Findable, Accessible, Interoperable, and Reusable) data principles. By leveraging the NASA OSDR, we demonstrate the compounding value of open space biology data.

1.5 Study Objectives and Research Questions

The primary objective of this thesis is to develop a comprehensive machine learning pipeline and an interactive web-based dashboard to perform a multi-crop comparative meta-analysis of NASA OSDR datasets OSD-745, OSD-655, and OSD-780. We aim to answer the following research questions:

1. **Multivariate Differentiation:** Can machine learning classifiers reliably distinguish between Flight and Ground samples based solely on their multielemental and biochemical profiles across different plant species?

2. **Cross-Crop Generalizability:** Does a classifier trained on one crop species (e.g., Lettuce) generalize successfully to another species (e.g., Mizuna) grown in the same Veggie spaceflight hardware?

3. **Universal Markers:** Which specific nutrients or combinations thereof serve as universal physiological markers of the spaceflight environment across species?

4. **Food Safety Meta-Analysis:** Do microbiological counts (APC and YMC) significantly differ between crop types or spaceflight conditions, and do they correlate with elemental composition?

5. **Data Accessibility:** How can interactive visualization tools lower the barrier to entry for space biology meta-analysis and data exploration?

Chapter 2

Materials & Methods

2.1 Data Source and Curation

The datasets utilized in this meta-analysis include three study entries from the NASA Open Science Data Repository (OSDR): OSD-745 (microbiological and nutritional profiles for space-grown lettuce [3]), OSD-655 (nutritional characterization of space-grown Mizuna mustard [4]), and OSD-780 (microbial plate count assays for Mizuna mustard). All files and associated metadata were programmatically fetched from the OSDR API (`https://osdr.nasa.gov`) utilizing standard REST queries and parsed from ISA-Tab (Investigation, Study, Assay) format sheets.

Agronomy Note: FAIR Principles

The data harmonization pipeline developed for this project aligns with FAIR principles. All raw assay files, metadata sheets, microbiological plate counts, and biochemical readouts across studies were merged into a consolidated, version-controlled analytical matrix.

2.2 Plant Material and Growth Conditions

Two leafy green crops were analyzed: *Lactuca sativa* cv. 'Outredgeous' (red romaine lettuce) grown during experimental epochs VEG-01A (May-Jun 2014), VEG-01B (Jul-Aug 2015), and VEG-03A (Oct-Dec 2016), and mizuna mustard (*Brassica rapa* var. *japonica*) grown during VEG-04A and VEG-04B. Both crops were seeded into Veggie "pillows" filled with an arcillite-based growth substrate and controlled-release fertilizer (Nutricote 18-6-8, type 180).

Cultivation occurred aboard the ISS (Flight) or under KSC ground environment-control chambers (Ground Control). The Veggie LED light cap provided Red (630 nm), Blue (455 nm), and Green (530 nm) lighting, with photoperiods of 16-hour light / 8-hour dark. Telemetry data from the ISS cabin (relative humidity, temperature, and carbon dioxide concentration) was mirrored in the ground control chambers with a 24- to 72-hour delay to isolate microgravity effects.

2.3 Sample Collection and Analytical Methods

Upon reaching maturity (approximately 33 days post-imbibition for lettuce and 28 days for Mizuna), crops were harvested by the crew. A portion of the fresh tissue was consumed in flight, while the remainder was immediately flash-frozen at -80°C using the MELFI (Minus Eighty-Degree Laboratory Freezer for ISS) and subsequently returned to Earth aboard SpaceX Dragon capsules for detailed laboratory analyses.

2.3.1 Elemental Profiling (ICP-OES)

Lyophilized leaf tissue was digested in nitric acid and analyzed for macronutrients and micronutrients using Inductively Coupled Plasma Optical Emission Spectrometry (ICP-OES) via a Thermo Scientific iCAP 7000 series spectrometer. Elements quantified included Iron (Fe), Potassium (K), Sodium (Na), Phosphorus (P), Sulfur (S), Zinc (Zn), Calcium (Ca), Magnesium (Mg), Manganese (Mn), and Copper (Cu).

2.3.2 Antioxidants and Phenolics

Total antioxidant capacity was determined using the Oxygen Radical Absorbance Capacity (ORAC) assay, read on a BioTek Synergy HT microplate reader. Total phenolic content was measured via the Folin-Ciocalteu colorimetric assay, and anthocyanins were quantified using the pH differential method.

2.3.3 Microbiological Plate Counts

For food safety validation, microbial loads on leaves and growth pillows were quantified. Tissues were homogenized, serial diluted, and plated in duplicate. Aerobic Plate Counts (APC) were evaluated on Plate Count Agar incubated at 30°C, and Yeast and Mold Counts (YMC) were evaluated on Potato Dextrose Agar incubated at 25°C. Colony-forming units (CFU) were counted after 48-72 hours and converted to $\log_{10}$ CFU/g dry weight.

2.4 Machine Learning Pipeline

The analytical pipeline was constructed using Python 3.10, leveraging `scikit-learn`, `pandas`, and `shap`.

2.4.1 Dimensionality Reduction

Principal Component Analysis (PCA) was employed for exploratory data analysis and unsupervised clustering. The elemental data matrix X was standard-scaled to zero mean and unit variance. The covariance matrix C was computed, and the eigenvalues λ and eigenvectors v were derived by solving:

$$Cv = \lambda v \tag{2.1}$$

where $C = \frac{1}{n-1} X^T X$. The first two principal components were extracted to visualize the variance captured by the multielemental profile.

2.4.2 Classification and Validation Strategies

To evaluate spaceflight signatures across crops, we trained classifiers to distinguish Flight vs. Ground Control samples based on elements and biochemicals. We implemented two validation designs:

- **Leave-One-Mission-Out Cross-Validation (LOMOCV):** The model was iteratively trained on four spaceflight missions (e.g., VEG-01A, VEG-01B, VEG-03A, VEG-04A) and validated on the remaining mission (e.g., VEG-04B), preventing overfitting to mission-specific microenvironments.
- **Leave-One-Crop-Out Cross-Validation (LOCOCV): not performed.** This item previously described training on one species and evaluating on the other. No such split was implemented: every `LeaveOneGroupOut` call in `analysis/` groups on mission, not crop. The item is kept in place, rather than deleted, so that readers of the earlier version can see what was claimed and that it was withdrawn. See Results for the cross-crop result obtained on the real data elsewhere.

We trained both Random Forest classifiers [5] and TabPFN tabular foundation models [7] under these designs.

The Random Forest utilized an ensemble of decision trees, minimizing Gini impurity:

$$G = 1 - \sum_{i=1}^{c} p_i^2 \tag{2.2}$$

where p_i is the probability of a sample belonging to class i at a given node.

2.4.3 SHAP Feature Importance

To demystify the "black box" nature of the Random Forest, SHapley Additive exPlanations (SHAP) [6] were utilized. SHAP values calculate the marginal contribution of each elemental feature to the model's prediction across all possible coalitions of features:

$$\phi_i(f, x) = \sum_{S \subseteq N \setminus \{i\}} \frac{|S|!(|N| - |S| - 1)!}{|N|!} \left[f(S \cup \{i\}) - f(S) \right] \tag{2.3}$$

where N is the set of all features, S is a subset of features excluding feature i, and f is the model prediction. This provided a ranked list of elements driving the FLT vs. GC distinction.

2.4.4 Tabular Foundation Model Benchmarking

To validate the Random Forest classification results, we implemented an independent benchmarking pipeline utilizing the state-of-the-art TabPFN (Tabular Prior-data Fitted Network) model

[7]. Unlike traditional classification algorithms (e.g., Random Forests, Gradient Boosted Trees, or Multi-Layer Perceptrons) that require backpropagation and parameter tuning on the training dataset, TabPFN is a transformer-based foundation model trained offline using millions of synthetic datasets.

At inference time, the model takes the entire training set (context) and the test set, performing a single forward pass (prior-data fitted learning) to generate zero-shot predictions. This approach is specifically optimized for small-sample classification tasks ($n < 10,000$ samples) and serves as a highly robust baseline to compare and contrast against the Random Forest classifier. Performance was evaluated using the identical LOMOCV strategy.

2.4.5 Statistical Testing

Non-parametric Mann-Whitney U tests were conducted for univariate comparisons. To account for multiple comparisons, p-values were adjusted using the Benjamini-Hochberg False Discovery Rate (FDR) procedure. Effect sizes were computed using Cohen's d:

$$d = \frac{\mu_1 - \mu_2}{s_{pooled}} \tag{2.4}$$

2.5 Interactive Dashboard Architecture

To democratize access to the OSD-745 results, a static interactive dashboard was developed. The application was built utilizing HTML, CSS, and modern JavaScript, leveraging `Chart.js` and `Plotly.js` for responsive, client-side rendering of the PCA, SHAP plots, and boxplots. The dashboard incorporates the CoSE design system's aesthetic guidelines and is hosted via GitHub Pages for persistent, serverless accessibility.

Chapter 3

Results

3.1 Data provenance and the 13 October 2026 correction

Every result in this chapter has been regenerated. The figures in earlier versions did not come from the NASA archive.

`data/fetch_osdr_data.py` previously returned values written as literals in its own source — first generated by `numpy.random.normal()`, later replaced by hand-typed numbers that still did not match OSDR. That table does not exist in the archive: it carried 36 lettuce samples against the archive's 24, calcium about three times the measured value, a copper column for a study that never measured copper, and no boron column for a study that did.

The fetcher now downloads and uses the archive, with no synthetic fallback. The cohort is 24 lettuce samples (11 flight, 13 ground) and 36 mizuna (18, 18), plus one mizuna sample carrying a phenolic measurement but no elemental data. Elements a study did not measure are left empty rather than filled. The five OSD-655 columns whose archived unit label is wrong by a factor of 1000 are corrected explicitly and the override is recorded.

What changed as a result is summarised below and detailed in each section.

- **Classification no longer separates flight from ground.** Random Forest accuracy is 0.490 and the benchmark model 0.572, against 83.3% and 89.6% previously — at or below chance for a two-class problem.
- **Elemental profile does not predict antioxidant capacity.** Scored out-of-fold, R^2 is -0.90 for phenolics and -2.16 for ORAC: worse than predicting the mean.
- **Three of five reported markers lose significance.** Only potassium and sulfur survive FDR correction.
- **The cross-species claim is withdrawn** (Section below), and so is the biochemical regression section.

3.2 Nutritional Profiling and Multi-Crop Composition

Elemental profiling of lettuce (*Lactuca sativa* cv. 'Outredgeous', OSD-745) and mizuna (*Brassica rapa* var. *japonica*, OSD-655) shows the two species differ in baseline composition, as expected for different crops: median calcium is 6.1 mg/g in mizuna against 4.4 mg/g in lettuce, and sulfur 5.0 against 1.7 mg/g. The superseded version of this paragraph reported 22 versus 12 mg/g calcium and 6.5 versus 3 mg/g sulfur; those values came from the literal table, not the archive.

Flight-versus-ground comparisons are given in Table 3.1. The earlier claim here of "depressed Potassium accumulation and increased secondary metabolite synthesis (phenolics) in flight" does not survive: median flight potassium is *higher* than ground in the archive data, and the phenolic difference is not significant after correction.

3.3 Joint PCA Dimensionality Reduction

To determine if spaceflight signatures are conserved across species, we performed a Principal Component Analysis (PCA) on the merged dataset.




Figure 3.1: Joint PCA of lettuce (OSD-745) and mizuna (OSD-655) on the 11 elemental and biochemical features with complete coverage ($N = 55$). PC1 separates the two crops. Lettuce samples group by mission rather than by flight or ground; mizuna flight samples tend to sit higher on PC2 than ground controls, with overlap.

As seen in Figure 3.1, Principal Component 1 (PC1) captures 47.9% of the variance and sepa-

rates the two crops by their baseline composition (mizuna on the left, lettuce on the right). PC2 captures 26.1% and does not show a shared flight response. Lettuce samples group by mission: VEG-01B lies apart from VEG-01A and VEG-03A, and flight and ground samples are interleaved within each mission. In mizuna, flight samples from both VEG-04A and VEG-04B tend to sit higher on PC2 than their ground controls, but the ranges overlap. The superseded version of this paragraph reported 42.0% and 25.0% and a consistent upward flight shift within both crops; the archive data reproduce neither.

3.4 Cross-Validation and Classification Benchmarks

We evaluated classifiers to distinguish FLT vs. GC using the 11 features with complete coverage across both crops ($N = 55$; 6 samples with an incomplete feature vector are dropped), under Leave-One-Mission-Out Cross-Validation (LOMOCV) across 5 independent spaceflights.




Figure 3.2: Classification performance (accuracy, precision, recall and F1) of the Random Forest classifier and the benchmark model under LOMOCV on the joint dataset. The benchmark is the Gradient Boosting fallback, because TabPFN did not run without a license token. Neither model separates flight from ground.

On the archive data the classifiers do not separate flight from ground. Random Forest reaches accuracy 0.490 (F1 0.455) and the benchmark model 0.572 (F1 0.444) under leave-one-mission-out cross-validation — at or below the ≈ 0.5 expected by chance on a two-class problem. The superseded figures were 89.6% and 83.3%.

Two caveats on the benchmark model. TabPFN requires a license token to download its

weights; where that token is absent the pipeline falls back to a `GradientBoostingClassifier`. The result file now records which model actually ran (`model` and `is_tabpfn` fields), because the fallback was previously stored under the `tabpfn` key and reported as a TabPFN result. The 0.572 above is Gradient Boosting, not TabPFN.

The cross-species generalizability claim has been withdrawn. This paragraph previously reported that under Leave-One-Crop-Out Cross-Validation (LOCOCV), a Random Forest trained on lettuce and evaluated on the second crop reached 79.2% accuracy and TabPFN 83.3%, and concluded that this "validates the existence of a universal microgravity-induced physiological signature."

No leave-one-crop-out cross-validation was ever run. Every `LeaveOneGroupOut` call in `analysis/` groups on mission: `ml_pipeline.py` line 181 (`groups = df['mission']`), `meta_ml_pipeline.py` line 97 (`groups_mission = df['mission'].values`) and `tabpfn_pipeline.py` line 162. No crop-based split exists anywhere in the codebase, and the figures 79.2% and 83.3% appear in no result file.

The analysis has since been performed on the real OSDR elemental data, in the SpaceMineralAtlas study. Grouping on crop across 60 samples, cross-crop transfer performs *below* a stratified-dummy baseline: balanced accuracy 0.486 for a random forest and 0.493 for L_2 logistic regression, against a baseline of 0.600 — margins of -0.113 and -0.107 respectively. Training on one crop and testing on the other is therefore worse than chance, which is the opposite of the conclusion withdrawn above. A universal cross-species signature is not supported by these data.

3.5 Feature Importance & SHAP Interpretation

Shapley Additive Explanations (SHAP) and Gini feature importance were extracted from the joint classifiers to identify the primary drivers of the spaceflight signature.

Regenerated from the archive data, the ranking is Potassium (MDI = 0.154), Sodium (0.132), Sulfur (0.106) and Manganese (0.095); iron is sixth at 0.081. The superseded text reported Fe = 0.28, K = 0.18, Ca = 0.15, Na = 0.12, which reproduced the placeholder object in `docs/js/app.js` rather than any computed output. Given that the classifier itself performs at chance, this ranking describes what the model attended to, not a set of validated spaceflight markers. The per-element interpretation that followed here (iron elevated in flight, potassium significantly lower in flight across both crops, calcium a critical classifier parameter) is withdrawn: potassium is higher in flight, and iron and calcium do not differ after FDR correction (Table 3.1).

3.6 Regression: Predicting Biochemical Outputs

This section has been withdrawn. It previously reported that a Random Forest regressor achieved an R^2 of 0.528 for ORAC and 0.491 for phenolics, and that sulfur and manganese were the top predictors of ORAC — read as evidence that sulfur-containing compounds and manganese-dependent superoxide dismutases are central to the spaceflight stress response. None of that is supported by this repository's own output.




Figure 3.3: Feature importance (mean decrease in impurity) for the joint flight-versus-ground Random Forest classifier. Potassium, sodium, sulfur and manganese rank highest. Because the classifier performs at chance, the ranking shows what the model attended to, not validated spaceflight markers.

- **The R^2 values appear in no result file.** They were written directly into `docs/index.html` and `docs/js/ml-results.js`. The regression that is actually computed (`analysis/ml_pipeline.py`) calls `fit` and `predict` on the same rows, so its R^2 of 0.96–0.99 is in-sample and is not a generalisation estimate either. No replacement figure is given here, because none can currently be sourced.

- **No regression feature importances were ever computed.** `meta_ml_results.json` contains `classification_metrics.feature_importance` only; there is no `regression_metrics` entry. In the classification importances that do exist, sulfur ranks fifth (0.068) and manganese eleventh of thirteen (0.023). The highest-ranked feature is phenolic content itself (0.351).

- **The biochemical inputs did not come from the archive.** The ORAC, total phenolic and anthocyanin assays were listed in the OSDR file manifests for OSD-745 and OSD-655 throughout, but had never been downloaded.

Those assays have since been retrieved from OSDR and analysed. Between-mission variation in them exceeds the within-mission flight effect by roughly an order of magnitude, so the comparison must block on mission; under a stratified test that does so, **no assay shows a flight effect at FDR 0.05 in either crop**. Full detail, including the measured values, is in `veggie_correction/CORRECTION_NOTE_1` in the SpaceMineralAtlas repository.

3.7 Microbiological Food Safety Analysis

This section is rebuilt from OSD-780 (LSDS-128 plate counts). The superseded version read per-sample APC and YMC values off the nutrient table; those columns were part of the literal data and were not measurements. It also reported counts for lettuce, and stated that "all harvested leaves fell well within safe food guidelines ($< 5.0 \log_{10}$ CFU/g)". Neither is supportable.

Scope. OSD-780 covers the VEG-04 (mizuna) missions only. **There is no microbiology for lettuce in the archive**, so no lettuce count can be reported. The study samples leaf, soil, root, wick, water and hardware swabs; only leaf is edible tissue, and the comparisons below use leaf alone. Swab and water samples carry no fresh weight and are reported per swab, so they are never pooled with per-gram tissue counts.




Figure 3.4: Aerobic plate counts and yeast/mould counts on mizuna leaf tissue (OSD-780), flight versus ground control. Plotted as $\log_{10}(\text{CFU/g} + 1)$: raw counts span zero to 1.7×10^7 CFU/g and over half the ground samples are exactly zero, so a linear axis and arithmetic means would both be misleading.

Flight leaf tissue carries a substantially higher microbial load. Median aerobic count is 7,380 CFU/g in flight ($n = 36$) against 0 in ground controls ($n = 87$), $p = 7.6 \times 10^{-10}$; yeast and mould 699 against 0, $p = 3.1 \times 10^{-7}$. Zero counts are common on the ground (52.9% of aerobic samples) and rare in flight (8.3%).

That comparison has a confound worth stating: sanitisation is labelled on ground samples only, so the whole-ground pool is roughly half sanitised while flight tissue carries no such label. Restricting the ground arm to unsanitised samples — the matched comparison — the difference persists: aerobic medians 7,380 versus 41 CFU/g, $p = 2.0 \times 10^{-6}$; yeast and mould 699 versus 0, $p = 1.4 \times 10^{-5}$. Sanitisation itself measurably reduces aerobic load on the ground (median 0 cleaned against 41 uncleaned, $p = 0.0025$) but has no detectable effect on yeast and mould ($p = 0.67$).

The earlier safety statement does not hold. Twelve of 123 aerobic leaf measurements (9.8%) exceed $5.0 \log_{10}$ CFU/g — nine flight, three ground — with a flight maximum of $7.23 \log_{10}$ (1.7×10^7 CFU/g). Yeast and mould counts stay below that bound throughout (maximum $4.99 \log_{10}$). Whether $5.0 \log_{10}$ is the applicable acceptance limit for this product is not established here, and

no acceptance threshold is asserted; what can be said is that the previous claim that all leaves fell well within it is contradicted by the archive data.

3.8 Statistical Validation and Effect Sizes

Non-parametric Mann-Whitney U tests with False Discovery Rate (FDR) corrections were performed on the joint dataset to validate the univariate significance of all features.

Table 3.1: Joint statistical comparison of key nutritional markers between Flight (FLT) and Ground (GC) cohorts.

Marker	Median FLT	Median GC	FDR p-value	Cohen's d
Iron (Fe) [mg/g]	0.063	0.051	0.413	0.21
Potassium (K) [mg/g]	36.7	23.9	0.043*	0.41
Sodium (Na) [mg/g]	0.926	0.570	0.052	0.30
Phosphorus (P) [mg/g]	3.6	2.9	0.117	0.60
Sulfur (S) [mg/g]	5.2	3.5	0.043*	0.84
Zinc (Zn) [mg/g]	0.043	0.036	0.457	0.14
Calcium (Ca) [mg/g]	5.6	5.3	0.457	0.37
Magnesium (Mg) [mg/g]	6.4	5.4	0.413	0.38
Manganese (Mn) [mg/g]	0.249	0.239	0.859	-0.05
Phenolics [GAE mg/g]	16.8	12.6	0.457	0.10
ORAC [μmol TE/g]	385.9	168.4	0.457	0.31

Table 3.1 is regenerated from the archive data. **Only potassium and sulfur differ significantly after FDR correction** ($p_{\mathrm{FDR}} = 0.043$ for both); sodium is marginal at 0.052. Iron, calcium and phenolics — all reported as significant in the superseded version of this table — are not ($p_{\mathrm{FDR}} = 0.413$, 0.457 and 0.457). That earlier table's Fe, K, Ca and S rows reproduced the placeholder object in `docs/js/app.js`, and its phenolics and ORAC rows matched neither those placeholders nor any computed output.

These two results should not be over-read either. The test pools both crops and all five missions without blocking on either. The two species differ substantially in elemental composition and the flight/ground split is not balanced identically across them, so a pooled difference can reflect which crop contributed which samples rather than spaceflight. A stratified analysis of the same kind applied to the antioxidant assays of these studies found no flight effect surviving mission-blocking at all. Potassium and sulfur are candidates for a properly blocked test, not established markers.

Chapter 4

Discussion

4.1 Nutritional Equivalence and Species Diversity

The comparative meta-analysis of Lettuce (*Lactuca sativa*, OSD-745 [3]) and Mizuna mustard (*Brassica rapa*, OSD-655 [4]) grown inside Veggie pillow modules demonstrates that both spaceflight crops maintain high nutritional density. Although baseline levels differ significantly—Mizuna naturally accumulates much higher Calcium and Sulfur than Lettuce—neither crop showed severe microgravity-induced nutrient degradation. This confirms that diverse crop selections can be introduced into astronaut diets without compromising food security or nutritional quality, supporting long-duration exploration missions.

4.2 Universal Mechanistic Responses to Microgravity

This section has been withdrawn. It argued that both crops share a universal elemental response to microgravity (iron up, potassium down and calcium shifted in flight) and proposed substrate hypoxia, transpiration and gravitropic signalling as mechanisms. The archive data do not show that pattern: the joint classifier performs at chance, potassium is higher rather than lower in flight, and iron and calcium do not differ after FDR correction (Table 3.1). With no consistent shift to explain, the proposed mechanisms are not discussed further.

4.3 The Power of Tabular Foundation Models (TabPFN)

This study set out to compare the TabPFN tabular foundation model [7] with a traditional Random Forest model. Space biology datasets are notoriously small due to the high cost of orbital flights. Classical machine learning algorithms trained on such small cohorts ($N = 55$ in our meta-analysis) are highly sensitive to overfitting and batch effects.

The accuracy figures this paragraph rested on (89.6% and 83.3%) were computed from data that did not come from the archive. Regenerated, the classifiers reach 0.572 and 0.490 — at or below chance for a two-class problem — so there is no performance difference here to interpret, and the benchmark figure is Gradient Boosting rather than TabPFN in any run without a license

token. **The cross-species sentence that followed here has been withdrawn.** It reported that TabPFN "generalized across species with 83.3% accuracy" under LOCOCV. No leave-one-crop-out split was implemented — every `LeaveOneGroupOut` call in `analysis/` groups on mission — and when the comparison is run on the real OSDR data, cross-crop transfer falls *below* a stratified-dummy baseline (balanced accuracy 0.486 versus 0.600). Whether tabular foundation models suit small space-biology datasets is therefore not something this study is positioned to demonstrate, and the claim that it does is withdrawn with it.

4.4 Microbiological Safety and Crew Hygiene

Comparing microbial safety counts between OSD-745 (lettuce) and OSD-780 (Mizuna) confirmed that both crops were safe to eat. Leaf surface bacterial and fungal loads remained well below food safety thresholds. The slightly elevated microbial counts observed in flight (Figure 3.4) highlight the microenvironmental challenges inside closed space cabins, where higher humidity and lower boundary-layer air convection facilitate localized microbe colonization. This emphasizes the need for continuous cabin sanitization and plant washing protocols before consumption.

4.5 Limitations of the Meta-Analysis

- **Baseline Variations:** Although the environmental parameters are matched via telemetry, batch-to-batch differences in chamber configurations, seed stocks, and analysis timelines between the Lettuce and Mizuna studies introduce potential confounding variables. - **Substrate Differences:** Differences in the arcillite clay blend ratios (100

4.6 Future Directions

Future work should focus on integrating multi-omics datasets (transcriptomics, metabolomics) to link our elemental and phenotypic observations back to specific gene expression pathways. Incorporating additional crops, such as dwarf tomatoes (VEG-05) and radishes (APH-01), will further expand this meta-analysis framework. The interactive dashboard developed here provides a scalable template for hosting and comparing these open science space agriculture datasets.

Chapter 5

Conclusion

The cultivation of fresh crops in microgravity is a critical milestone for long-duration human space exploration. The NASA OSDR OSD-745, OSD-655, and OSD-780 datasets provided an invaluable opportunity to evaluate space agriculture systems.

In this work, we developed a comprehensive machine learning and meta-analysis pipeline to study Lettuce and Mizuna mustard grown aboard the ISS. On the archive data, the Random Forest classifier does not separate flight from ground samples, so this analysis does not demonstrate a conserved multielemental flight fingerprint. Potassium and sulfur differ after FDR correction in a pooled test that is not blocked by crop or mission (Table 3.1), and flight mizuna leaf carries a substantially higher microbial load than ground controls (Figure 3.4).

The cross-species conclusion has been withdrawn. It previously stated that TabPFN showed "strong cross-species generalizability (83.3%) under LOCOCV", and that this "validates the existence of a universal spaceflight physiological signature across plant families".

No leave-one-crop-out cross-validation was performed in this study. Run on the real OSDR elemental data, cross-crop transfer performs below a stratified-dummy baseline (balanced accuracy 0.486 against 0.600), so the data support the opposite reading: the flight signature does not transfer between these two species. The remaining classifier comparison (Random Forest against the Gradient Boosting fallback, as TabPFN did not run) is a leave-one-mission-out result at chance level and should not be read as evidence of cross-species generality.

Beyond the scientific findings, this project actively championed the principles of Open Science. By aggregating, cleaning, and publishing a harmonized meta-analysis matrix, we have enhanced the FAIR properties of the original OSDR data. The accompanying interactive web dashboard democratizes access to these complex machine learning insights, allowing researchers, agronomists, and the public to visually explore the data.

As humanity prepares for lunar bases and crewed missions to Mars, the integration of advanced data science with space biology will become increasingly indispensable. The methodologies and tools developed in this thesis lay a scalable foundation for the analysis of future bioregenerative life support systems, ensuring that astronauts remain healthy, nourished, and connected to Earth.

Bibliography

[1] Raymond M Wheeler. Agriculture for space: People and places paving the way. *Open Agriculture*, 2(1):14–32, 2017.

[2] Paul Zabel, Matthew Bamsey, Daniel Schubert, and Martin Tajmar. Review and analysis of over 40 years of space plant growth systems. *Life sciences in space research*, 10:1–16, 2016.

[3] Christina L Khodadad, Mary E Hummerick, LaShelle E Spencer, Anirudha R Dixit, Jeffrey T Richards, Matthew W Romeyn, Trent M Smith, Raymond M Wheeler, and Gioia D Massa. Microbiological and nutritional analysis of lettuce crops grown on the international space station. *Frontiers in Plant Science*, 11:199, 2020. doi: 10.3389/fpls.2020.00199.

[4] Jess M Bunchek, Mary E Hummerick, LaShelle E Spencer, Matthew W Romeyn, Millennia Young, Robert C Morrow, Cary A Mitchell, Grace L Douglas, Raymond M Wheeler, and Gioia D Massa. Pick-and-eat space crop production flight testing on the international space station. *Journal of Plant Interactions*, 19(1):2292220, 2024. doi: 10.1080/17429145.2023.2292220.

[5] Leo Breiman. Random forests. *Machine learning*, 45(1):5–32, 2001.

[6] Scott M Lundberg and Su-In Lee. A unified approach to interpreting model predictions. In *Advances in Neural Information Processing Systems 30*, pages 4765–4774. Curran Associates, Inc., 2017.

[7] Noah Hollmann, Samuel Müller, Lennart Purucker, Arjun Krishnakumar, Max Körfer, Shi Bin Hoo, Robin Tibor Schirrmeister, and Frank Hutter. Accurate predictions on small data with a tabular foundation model. *Nature*, 637:319–326, 2025. doi: 10.1038/s41586-024-08328-6.