# Supplementary information: PathoFluor-At

Richard J. Barker [PENDING: co-authors]

## Supplementary Note 1. Assigning genotype and plant labels to segmented seedlings

Each row-scan images one row of seedlings along the scan (line) axis. Genotype lanes are columns along that axis, shared by a plate's three rows. For every plate and timepoint we pooled the line positions of all segmented seedlings and clustered them into five lanes (set A) or three (set B) by exact one-dimensional $k$-means. Plant numbers follow reading order within a lane, which is how the 2016 hand ROIs are numbered.

Plates were not always placed in the same orientation on the stage. The 2016 hand ROIs show *cml24-2* (lane D) with the highest $F_{685}/F_{740}$ on every set-A plate, while in the segmented images the high-ratio lane changes position between plates. We therefore chose the scan direction per plate and timepoint. For each direction we paired every hand ROI with the seedling that direction assigns to it, and scored the within-plate Spearman correlation of three features ($F_{685}/F_{740}$, red-peak width, peak intensity). We kept the direction with the higher mean (Table S1). Plates without hand ROIs (set-A Pathogen at 48 h) receive no genotype label and enter only treatment-level analyses. Set-B hand ROIs are per leaf, so set-B lanes follow the order stated in the vendor report (Col-0, *fls2-17*, *fls2-101*; FACTS R9), pending confirmation.

## Supplementary Note 2. The 2016 hand-drawn ROIs

The 2016 annotator drew ROIs on seedlings and exported their mean spectra. Copies were sorted by timepoint, by genotype, and into curated "USE" subsets (1,358 set-A spectra), plus 16 Windows "- Copy" duplicates. Identical copies were merged by content hash. 12 spectra share a design key with a different spectrum (re-drawn ROIs); both are kept and flagged. ROIs carry 25 distinct free-text tags ("bright", "norm", "dark", "side", "dying" and others) whose meaning is pending (FACTS P6). They are retained in `results/roi_features.csv.gz`. No ROIs exist for set-A Pathogen plates at 48 h.

## Supplementary Note 3. The 2016 vendor analyses

Two Middleton Spectral Vision reports analysed subsets of these data by multivariate curve resolution. For the July set (one infected and one control plate, two plants per genotype) the report describes two chlorophyll-like components and a broad autofluorescence component present mainly in plants infected for at least 24 h, with infected and control plants separated by component medians. For the November set (one plate per treatment) it describes a "chlorophyll shift" component that rises in *Pseudomonas* plants from 6 h and in flg22-treated Col-0 but not *fls2*, tested with notched box plots of individual plants. Neither analysis treated the plate as the experimental unit or used a pre-treatment baseline. We treat their conclusions as hypotheses, tested in the main text. The reports are cited, not redistributed (FACTS P11).

**Table S1:** Scan direction per set-A plate and timepoint, chosen by agreement between the 2016 hand ROIs and the seedlings each direction assigns to them (mean within-plate Spearman correlation of three features). Source: results/label_direction_scores.csv.

| Hour | Treatment | Plate | ROIs | Seedlings | Score fwd | Score rev | Chosen |
|---|---|---|---|---|---|---|---|
| 0 | Control | 1 | 29 | 28 | 0.78 | -0.36 | fwd |
| 0 | Control | 2 | 30 | 27 | 0.50 | 0.48 | fwd |
| 0 | Control | 3 | 30 | 23 | 0.28 | 0.17 | fwd |
| 0 | Control | 4 | 28 | 23 | -0.06 | -0.61 | fwd |
| 0 | Pathogen | 1 | 27 | 24 | 0.44 | 0.04 | fwd |
| 0 | Pathogen | 2 | 26 | 29 | -0.32 | 0.58 | rev |
| 0 | Pathogen | 3 | 27 | 26 | -0.10 | 0.35 | rev |
| 0 | Pathogen | 4 | 30 | 28 | 0.27 | 0.06 | fwd |
| 0 | flg22 | 1 | 29 | 28 | 0.84 | 0.37 | fwd |
| 0 | flg22 | 2 | 30 | 23 | 0.07 | 0.28 | rev |
| 0 | flg22 | 3 | 30 | 28 | 0.22 | 0.24 | rev |
| 0 | flg22 | 4 | 29 | 22 | -0.10 | 0.04 | rev |
| 1 | Control | 1 | 29 | 25 | 0.13 | 0.46 | rev |
| 1 | Control | 2 | 30 | 24 | 0.47 | -0.08 | fwd |
| 1 | Control | 3 | 30 | 21 | -0.29 | 0.19 | rev |
| 1 | Control | 4 | 28 | 23 | 0.12 | -0.47 | fwd |
| 1 | Pathogen | 1 | 28 | 19 | 0.04 | 0.32 | rev |
| 1 | Pathogen | 2 | 29 | 26 | 0.83 | 0.09 | fwd |
| 1 | Pathogen | 3 | 29 | 23 | -0.02 | 0.01 | rev |
| 1 | Pathogen | 4 | 26 | 20 | -0.33 | 0.34 | rev |
| 1 | flg22 | 1 | 29 | 28 | 0.21 | 0.38 | rev |
| 1 | flg22 | 2 | 29 | 23 | 0.44 | -0.03 | fwd |
| 1 | flg22 | 3 | 29 | 19 | -0.05 | -0.21 | fwd |

## Supplementary Note 4. Pilot experiment

## Supplementary Table: strongest baseline-adjusted contrasts





**Figure S1:** Pilot (February–March 2016) leaf area, days 0–3, in units of a red reference square labelled 4 cm$^2$ (multiply by 4 for cm$^2$ if the label is right). 98.3% of the tabulated values match that column of the source sheet (FACTS R10, P12). Data: `tables/figS1_pilot_leaf_area.csv`; source sheet in `data/related_drive_2016/`.

**Table S2:** Baseline-adjusted contrasts with $p < 0.01$ (set A; all 270 tests in tables/stats_contrasts_baseline_adjusted.csv).

| Hour | Feature | Contrast | Effect (SD) | Effect (units) | $p$ | $q$ |
|---|---|---|---|---|---|---|
| 2 | red_fwhm_nm | Pathogen vs Control (Col-0) | 2.39 | 1.765 | 4.9e-06 | 0.001 |
| 2 | shortwave_fraction | Pathogen vs Control (Col-0) | 2.26 | 0.0008884 | 4.0e-05 | 0.005 |
| 1 | red_fwhm_nm | Pathogen vs Control (Col-0) | 2.07 | 1.406 | 5.0e-04 | 0.045 |
| 2 | farred_fraction | Pathogen vs Control (Col-0) | 1.87 | 0.02807 | 1.0e-03 | 0.061 |
| 1 | farred_fraction | Pathogen vs Control (Col-0) | 1.95 | 0.02882 | 1.1e-03 | 0.061 |
| 48 | red_fwhm_nm | flg22 vs Control (Col-0) | 1.50 | 0.9629 | 1.6e-03 | 0.073 |
| 2 | centroid_nm | Pathogen vs Control (Col-0) | 1.83 | 1.677 | 2.2e-03 | 0.083 |
| 1 | centroid_nm | Pathogen vs Control (Col-0) | 1.89 | 1.709 | 2.6e-03 | 0.087 |
| 2 | f685_f740 | Pathogen vs Control (Col-0) | -1.74 | -0.8409 | 3.9e-03 | 0.107 |
| 3 | red_fwhm_nm | Pathogen vs Control (Col-0) | 1.88 | 2.085 | 4.1e-03 | 0.107 |
| 2 | red_fwhm_nm | flg22 vs Control (Col-0) | 1.60 | 1.178 | 4.3e-03 | 0.107 |
| 1 | f685_f740 | Pathogen vs Control (Col-0) | -1.75 | -0.8463 | 7.0e-03 | 0.158 |
| 48 | farred_fraction | flg22 vs Control (Col-0) | 1.30 | 0.01536 | 9.0e-03 | 0.181 |
| 48 | f685_f740 | flg22 vs Control (Col-0) | -1.27 | -0.5162 | 9.4e-03 | 0.181 |