Fluorescence hyperspectral imaging detects an early, FLS2-independent signature of *Pseudomonas syringae* challenge in *Arabidopsis* seedlings, but not of flg22 perception

Richard J. Barker*[1,2], Aditya Singh[3], Simon Gilroy[1], and [PENDING: author order; further authors, e.g. Sarah Sapiro, Cullen Vens (FACTS P10)]

[1]Department of Botany, University of Wisconsin–Madison, USA
[2]Porterfield Lab, Purdue University, USA
[3]Department of Forest and Wildlife Ecology, University of Wisconsin–Madison, USA; present address: University of Florida, USA

[PENDING: date]

Abstract

Detecting plant infection before symptoms appear would help both field and controlled-environment agriculture. In 2016 we imaged *Arabidopsis* seedlings by fluorescence hyperspectral imaging (488 nm excitation, 500–800 nm emission) before and 1–48 h after treatment with a mock solution, the flagellin epitope flg22 or *Pseudomonas syringae* pv. tomato DC3000. Five genotypes were imaged: Col-0, two *fls2* receptor mutants and two *cml24* Ca^{2+}-sensor mutants. We reanalysed all 1,982 hand-drawn seedling spectra of the primary experiment, and 400 raw image cubes (396 GB). Recovering the camera metadata showed that exposure time changed between timepoints: the 2016 summaries' 21% fall in intensity after treatment disappears (to within 4%) once intensities are divided by exposure. In addition, 83 exported spectra had been saved without dark subtraction; we corrected every spectrum by its own emission below the excitation wavelength. Treated and control plates also differed *before* treatment, enough for a classifier to tell them apart (plate-level AUC 1.00), so we analysed only change from each plate's own baseline. On that basis, *P. syringae* broadened the red chlorophyll emission peak from 1 h (+1.40 nm; $q = 0.018$) through 24 h, and raised emission below 640 nm from 2 h. Change-from-baseline spectra separated infected from control plates perfectly at 1, 2 and 24 h (leave-plate-out AUC 1.00; exact permutation $p = 0.029$). The pathogen signature appeared in every genotype; at 24 h the rise in short-wavelength emission was smaller in *cml24-4* ($q = 0.036$). A flg22 signal was weaker (AUC 0.89–1.00, $p \geq 0.10$). It was as strong in both *fls2* mutants, which cannot perceive flg22, as in Col-0, so in this experiment it reflects a plate-level effect rather than flg22 perception. We release the raw data, manifests, code and a reproducible pipeline as a FAIR resource, and document every data problem we found.

1 Introduction

Plants perceive bacteria through pattern-recognition receptors long before disease symptoms appear. The best-characterised receptor is FLS2, which binds the conserved flg22 epitope of bacterial flagellin [1–4]. flg22 induces defence-gene expression and resistance in wild-type plants but not in *fls2* mutants. *fls2* plants still respond to bacterial extracts through other patterns, and are more susceptible to *P. syringae* pv. tomato DC3000 sprayed on the leaf surface [5].

*ORCID 0000-0001-5681-9857. Correspondence: richard@cosecloud.com

Pattern-triggered defence quickly lowers non-photochemical quenching [6]. *P. syringae* infection changes photosynthetic chlorophyll fluorescence parameters near the infection site before visible symptoms, within 3 h for an avirulent strain [7]. Chlorophyll fluorescence imaging is therefore a candidate for disease detection and phenotyping [8–11].

Steady-state chlorophyll fluorescence emission has two bands, near 685–690 nm and 730–740 nm. Because chlorophyll reabsorbs the red band, their ratio falls as chlorophyll content rises. The ratio also depends on leaf structure, optics and photosynthetic activity [12]. Fluorescence hyperspectral imaging records the whole emission spectrum at every pixel, so subtle shape changes can be mapped, not just one or two ratios.

CML24 (*TCH2*) encodes a calmodulin-like Ca^{2+} sensor whose transcript is induced by touch, darkness and other stimuli [13]. The *cml24-4* mutant is impaired in pathogen- and pattern-induced nitric-oxide production [14]. This makes CML24 a candidate link between early Ca^{2+} signalling and downstream changes visible in fluorescence.

Here we reanalyse a 2016 experiment designed to ask whether bacterial challenge and flg22 perception leave a hyperspectral fluorescence signature, and whether that signature depends on FLS2 or CML24. Its fls2 mutants supply the key control: an flg22 signature that reflects perception should be absent in them.

2 Results

2.1 Recovering the experiment from its raw files

We inventoried 7,752 files and parsed every file name into a design record, with no exceptions. That gave 216 sample cubes for the primary July experiment (set A), 168 for an independent November replicate (set B) and 5,842 hand-drawn ROI spectra (2,486 unique by content). The headers revealed that exposure changed between timepoints (Fig. 1a). The 2016 graphs plotted raw counts, so their drop in Control intensity from 2,588 to 2,044 counts after hour 0 (Fig. 1b) is largely a camera setting. Exposure-corrected Control intensities at 1–3 h lie within 4% of hour 0 (Fig. 1c). Hour-0 scans also saturated the 12-bit detector in up to 13% of a cube's seedling pixels (median 0.1%). A further 83 hand-ROI spectra, all from Pathogen plates (3 h plates 1–2 and 24 h plate 4), had been exported without dark subtraction. Their emission below the excitation wavelength sat at the camera's dark level (median 109 counts, against 0.1 for all other set-A spectra). Uncorrected, this offset masquerades as broad short-wavelength "autofluorescence". Nothing can fluoresce at wavelengths shorter than the 488 nm excitation, so we subtracted each spectrum's own mean over 420–470 nm. We therefore based the primary analysis on exposure- and offset-independent shape features.

2.2 Plates differed before treatment

Hour 0 was imaged before treatment (FACTS R13). Even so, flg22 plates already differed from Control plates in red-peak width (-1.29 SD; $q = 5.5 \times 10^{-5}$). A classifier separated them perfectly from spectra alone (plate-level AUC 1.00, $p = 0.029$; Fig. 4a). The same held, less strongly, for Pathogen plates (AUC 0.81). Plate-to-plate differences in seedling batch, placement or imaging geometry are thus as large as the effects we want to detect. All inference below uses change from each plate's own hour-0 baseline, with the plate as the experimental unit.

2.3 An early pathogen signature

Relative to baseline, *P. syringae* broadened the red emission peak of Col-0 seedlings by 1.40 nm at 1 h (2.07 SD, $q = 0.018$), 1.76 nm at 2 h (2.39 SD, $q = 6.8 \times 10^{-4}$), 1.02 nm at 3 h ($q = 0.005$) and 3.35 nm at 24 h ($q = 0.005$). It raised the 500–640 nm fraction of emission from 2 h (1.77 SD, $q = 0.018$) to 24 h (2.46 SD, $q = 4.3 \times 10^{-5}$; Figs 2, 3). 13 of 270 baseline-adjusted tests were significant at $q < 0.05$ (Supplementary Table). A classifier on change-from-baseline spectra




Figure 1: The 2016 intensity drop after treatment is an exposure artefact. (a) Exposure time per timepoint (plate mean, Control plates). (b) Median 683–687 nm emission of Control hand ROIs as plotted in 2016 and (c) divided by exposure. Data: `results/roi_spectra_A.parquet`, `metadata/cube_manifest.csv`.




Figure 2: Treatment-induced change in emission-spectrum shape (Col-0, set A; area-normalised spectra, change from each plate's hour 0, treated minus Control).

separated infected from control plates perfectly at 1, 2 and 24 h (AUC 1.00, 1.00, 1.00; exact $p = 0.029$ each, close to the smallest p the 70 plate relabellings allow; Fig. 4b), and less well at 3 h (AUC 0.75, $p = 0.143$). The 2016 vendor analysis of a single infected and a single control plate reported a broad autofluorescence component in infected plants after 24 h [15]. Our short-wavelength fraction measures that component across all plates.

2.4 The flg22 signal does not depend on FLS2

flg22 also broadened the red peak at 3 h (2.18 SD, $q = 0.005$). After baseline adjustment, flg22 plates were separable from Control plates at 1–3 h (AUC 0.89–1.00; Fig. 4b). With three plates per treatment left, however, no result beat $p = 0.10$ (20 relabellings). The genetics decide what this signal is. A model trained on Col-0 flg22-versus-Control shape features gave an average flg22 score difference of 0.18 in Col-0 (out-of-fold, 1–3 h), 0.36 in *fls2-17* and 0.24 in *fls2-101* (Fig. 5a). flg22 cannot act through FLS2 in the mutants, so this signal is a property of the plates (handling, timing or seedling batch), not of flg22 perception. The pathogen signature was likewise present in all genotypes (Fig. 5b). This is expected, because bacteria are perceived through patterns other than flagellin [5].




Figure 3: Time course of three emission-shape features (change from hour 0; mean ± s.e.m. over plates). Data: `tables/fig3_timecourse.csv`.




Figure 4: Plate-level classification of treatment from spectra, Col-0. (a) Absolute spectra; hour 0 is before treatment. (b) Change from hour 0. * exact permutation $p \leq 0.05$.

2.5 The independent replicate

In the November replicate (set B; 405 nm excitation), hand ROIs exist for one plate per treatment, so treatment main effects cannot be separated from plate effects there. Within plates, however, flg22 changed the emission shape of Col-0 differently from *fls2-17* at 6 h (F_{685}/F_{740} interaction 1.78 SD, $q = 0.026$; 4 related features). This is the pattern expected of FLS2-dependent perception. It was not seen for *fls2-101*. [PENDING: Plate-replicated set-B analysis from the raw cubes (2–3 plates per treatment).]

2.6 Seedling-level analysis of the raw cubes

[PENDING: Segmentation of all cubes (Pathogen 48 h, exposure-corrected intensity, spatial heterogeneity) and pixel-level multivariate curve resolution [16]; label-transfer validation in Supplementary Note 1.]

3 Discussion

Three lessons from this reanalysis apply beyond these data. First, acquisition metadata must travel with hyperspectral intensities. A camera setting that changed between timepoints produced the most striking pattern in the 2016 summaries. Second, in plate-based seedling assays the plate, not the seedling, is the treated unit. Plates can differ before treatment by as much as treatments




Figure 5: Genotype specificity of the signatures. Col-0-trained model applied to each genotype; difference in mean predicted probability between treated and Control plates.

differ afterwards, so a baseline image and leave-plate-out validation are essential. Without them, a classifier here would have "detected" flg22 before it was applied. Third, mutants that cannot perceive the stimulus are the decisive control. Here they showed that an apparently promising flg22 signal was not perception.

What survives is a pathogen signature within 1–2 h of inoculation: a broader red emission peak and more short-wavelength emission. The timing is consistent with reports that chlorophyll fluorescence changes precede visible symptoms [7]. Because the signature appears in *fls2* mutants, it does not require flagellin perception. Loss of CML24 may modulate it: at 24 h the short-wavelength rise was smaller in *cml24-4* (-1.74 SD, $q = 0.036$), the allele that is impaired in pattern-induced NO signalling [14]. The second allele, *cml24-2*, did not differ, so this needs replication. [PENDING: seedling-level test of the cml24 interaction] Limitations: inoculum density, seedling age and growth conditions are not recorded with the data [PENDING: FACTS P1, P4]; there were four plates per treatment; and set B used a different excitation wavelength.

4 Methods

4.1 Plant material and treatments

Arabidopsis thaliana seedlings of five genotypes were grown on square agar plates in five lanes per plate: Col-0 (lane A), *fls2-17* (B), *fls2-101* (C), *cml24-2* (D) and *cml24-4* (E). Lane assignments are taken from the 2016 analysis folders (FACTS R8). [PENDING: Allele origins, backgrounds and stock numbers.] *FLS2* encodes the flagellin receptor that perceives the flg22 epitope [1–4]. *CML24* encodes a calmodulin-like Ca^{2+} sensor [13] implicated in flg22-triggered nitric-oxide signalling [14]. Seedlings were grown on [PENDING: medium (protocol draft: ½ LS, 0.3% P), age, light, photoperiod and temperature]. Plates were treated with a mock solution, flg22 (1 μM, the laboratory's standard working concentration; FACTS R17) or *Pseudomonas syringae* pv. tomato DC3000 (FACTS R5, R16) [PENDING: inoculum density and application method, FACTS P1–P2]. The 2016 protocol draft specifies flood inoculation after Ishiga et al. [17], with Silwet L-77 as the mock. Four plates per treatment were imaged in the July experiment (set A).

The independent November experiment (set B; FACTS R7, R9) used Col-0, *fls2-17* and *fls2-101* only, with two or three plates per treatment, three rows per plate and three seedlings per genotype per row. A pilot experiment (February–March 2016) measured rosette leaf area from images of eight genotypes, including *glr3.3* and *glr3.4* alleles (FACTS R10–R11). It is reported in the supplement only.

4.2 Fluorescence hyperspectral imaging

Plates were imaged in the Gilroy Laboratory (UW-Madison Botany) with a MacroPhor™ fluorescence hyperspectral line-scan imager (Middleton Spectral Vision; FACTS R1). It carried an Andor Zyla 5.5 sCMOS camera in 12-bit low-noise mode (FACTS R2). Set A used 488 nm excitation and 500–800 nm emission (FACTS R3). The camera recorded 324 bands from 400.3 to 808.7 nm at about 1.23 nm spacing. Set B used 405 nm excitation and 400–800 nm emission (FACTS R4), recorded as 406 bands from 400.2 to 800.0 nm, each row-scan acquired at two excitation laser powers, 50% and 58% of maximum (`Laser50`, `Laser58`; FACTS R15). Each plate was imaged as three row-scans, one row of seedlings each. Set A was imaged at `Hour0` (11:20–12:26 on 22 July 2016, about 6 h before the 1 h scans; the pre-treatment baseline, FACTS R13) and at 1, 2, 3, 24 and 48 h after infection (FACTS R6). Set B was imaged at 0, 6, 24 and 48 h. Acquisition settings were read from every ENVI header (`metadata/cube_manifest.csv`).

4.3 Data provenance and quality

All 7,752 files on the 2016 drive were inventoried and hashed (`scripts/00_inventory.py`). Every file name was parsed without exception: 400 hyperspectral cubes (216 set-A samples, 168 set-B samples, 8 set-B bacteria-only scans, 6 set-B reference scans, one reference and one artefact scan in set A) and 5,842 hand-drawn region-of-interest (ROI) spectra (2,486 unique by content hash). We found the following problems and corrected for each where possible.

1. **Exposure time changed between timepoints.** It was 0.305 s at `Hour0`, 0.203 s at 1–3 h, 0.222–0.268 s at 24 h (varying within plates) and 0.222 s at 48 h. The 2016 summary graphs plotted uncorrected intensities, so their apparent drop between 0 and 1 h (about 1.5-fold) matches the exposure ratio $0.305/0.203 = 1.50$. We divide every intensity by its own exposure time, and our primary features are exposure-independent ratios (Section 4.4).

2. **`Hour0` plates were scanned "inverted"**, i.e. upside down (FACTS R14), except four rescans labelled "NI" made later the same day at the 1 h settings. `Hour0` is used only as a baseline and is analysed separately from the upright timepoints.

3. **Detector saturation.** At the `Hour0` exposure, up to [PENDING: N]% of seedling pixels reached the 12-bit ceiling (4095 counts). Saturated pixels are excluded from all spectra.

4. **Fixed scan order.** Within each timepoint plates were scanned Pathogen, then flg22, then Control, 20–40 min apart, so scan time is partly confounded with treatment. We report the drift of Control plates between consecutive hours as a bound on this effect.

5. **The plate is the experimental unit.** Treatments were applied per plate (four per treatment in set A), so treatment effects are tested at plate level. Genotype contrasts are within plates.

6. **Inconsistent dark subtraction in the 2016 exports.** 83 hand-ROI spectra (Pathogen 3 h plates 1–2 and 24 h plate 4) were exported with the camera dark level (median 109 counts) still in them; all others were dark-subtracted (median 0.1 counts). Because nothing fluoresces at wavelengths shorter than the excitation, each spectrum's mean below the excitation line (420–470 nm in set A, 400–404 nm in set B) was subtracted from it before features were computed (`scripts/01_import_rois.py`).

7. **Missing cells.** No hand ROIs were drawn for Pathogen plates at 48 h, although the cubes exist. Our automated segmentation covers them.

4.4 Spectral processing

Each raw cube was read once, sequentially, and hashed with SHA-256 (`scripts/02_harvest_cubes.py`). The per-band dark level was taken as the median of background pixels, which sit at about 108 counts. Seedlings were segmented on the dark-subtracted 685 nm image at 10 background standard deviations, then cleaned, and cotyledon and hypocotyl components were grouped (`src/pathofluor/segment.py`). Each segmented seedling was assigned its lane and plant number from the row layout, which was validated against the 2016 hand ROIs (Supplementary Note 1). Spectral features were computed from the mean unsaturated spectrum of each seedling after Savitzky–Golay smoothing (`src/pathofluor/features.py`):

- the red/far-red ratio F_{685}/F_{740} [12];
- the position and full width at half maximum of the red emission peak;
- the spectral centroid;
- the far-red fraction of 650–800 nm emission;
- the short-wavelength fraction (500–640 nm emission relative to 650–800 nm), which captures the broad autofluorescence reported by the 2016 vendor analysis;
- exposure-normalised intensities (counts s^{-1});
- per-pixel heterogeneity of the ratio and of the short-wavelength fraction.

4.5 Statistics and classification

[PENDING: Final model specification, written after the analysis is frozen.] Treatment effects were tested with linear mixed models with plate as a random effect, against time-matched controls, with false-discovery-rate control [18]. Classifiers were evaluated by leave-one-plate-out cross-validation, with a plate-level permutation null.

4.6 Data and code availability

Code (MIT licence), derived data, the 2016 hand-ROI spectra and all manifests are in the repository. Raw cubes are deposited on Zenodo [PENDING: DOIs]. The two 2016 Middleton Spectral Vision analysis reports [15, 19] are cited but not redistributed (FACTS R19). The metadata follow MIAPPE where applicable [20] and the FAIR principles [21].

Acknowledgements

[PENDING: Acknowledgements and funding (FACTS P10).]

References

[1] Georg Felix, Juliana D. Duran, Sigrid Volko, and Thomas Boller. Plants have a sensitive perception system for the most conserved domain of bacterial flagellin. *The Plant Journal*, 18(3):265–276, 1999. ISSN 1365-313X. doi: 10.1046/j.1365-313x.1999.00265.x. URL `http://dx.doi.org/10.1046/j.1365-313x.1999.00265.x`.

[2] Lourdes Gómez-Gómez, Georg Felix, and Thomas Boller. A single locus determines sensitivity to bacterial flagellin in arabidopsis thaliana. *The Plant Journal*, 18(3):277–284, 1999. ISSN 1365-313X. doi: 10.1046/j.1365-313x.1999.00451.x. URL `http://dx.doi.org/10.1046/j.1365-313x.1999.00451.x`.

[3] Lourdes Gómez-Gómez and Thomas Boller. Fls2: an lrr receptor-like kinase involved in the perception of the bacterial elicitor flagellin in arabidopsis. *Molecular Cell*, 5(6):1003–1011, 2000. ISSN 1097-2765. doi: 10.1016/s1097-2765(00)80265-8. URL http://dx.doi.org/10.1016/s1097-2765(00)80265-8.

[4] Delphine Chinchilla, Zsuzsa Bauer, Martin Regenass, Thomas Boller, and Georg Felix. The *Arabidopsis* receptor kinase fls2 binds flg22 and determines the specificity of flagellin perception. *The Plant Cell*, 18(2):465–476, 2005. ISSN 1532-298X. doi: 10.1105/tpc.105.036574. URL http://dx.doi.org/10.1105/tpc.105.036574.

[5] Cyril Zipfel, Silke Robatzek, Lionel Navarro, Edward J. Oakeley, Jonathan D. G. Jones, Georg Felix, and Thomas Boller. Bacterial disease resistance in arabidopsis through flagellin perception. *Nature*, 428(6984):764–767, 2004. ISSN 1476-4687. doi: 10.1038/nature02485. URL http://dx.doi.org/10.1038/nature02485.

[6] Vera Göhre, Alexandra M. E. Jones, Jan Sklenář, Silke Robatzek, and Andreas P. M. Weber. Molecular crosstalk between pamp-triggered immunity and photosynthesis. *Molecular Plant-Microbe Interactions*, 25(8):1083–1092, 2012. ISSN 1943-7706. doi: 10.1094/mpmi-11-11-0301. URL http://dx.doi.org/10.1094/mpmi-11-11-0301.

[7] Katharina B. Bonfig, Ulrich Schreiber, Andrea Gabler, Thomas Roitsch, and Susanne Berger. Infection with virulent and avirulent p. syringae strains differentially affects photosynthesis and sink metabolism in arabidopsis leaves. *Planta*, 225(1):1–12, 2006. ISSN 1432-2048. doi: 10.1007/s00425-006-0303-3. URL http://dx.doi.org/10.1007/s00425-006-0303-3.

[8] Céline Rousseau, Etienne Belin, Edouard Bove, David Rousseau, Frédéric Fabre, Romain Berruyer, Jacky Guillaumès, Charles Manceau, Marie-Agnès Jacques, and Tristan Boureau. High throughput quantitative phenotyping of plant resistance using chlorophyll fluorescence image analysis. *Plant Methods*, 9(1), 2013. ISSN 1746-4811. doi: 10.1186/1746-4811-9-17. URL http://dx.doi.org/10.1186/1746-4811-9-17.

[9] María L. Pérez-Bueno, Mónica Pineda, Francisco M. Cabeza, and Matilde Barón. Multicolor fluorescence imaging as a candidate for disease detection in plant phenotyping. *Frontiers in Plant Science*, 7, 2016. ISSN 1664-462X. doi: 10.3389/fpls.2016.01790. URL http://dx.doi.org/10.3389/fpls.2016.01790.

[10] María Luisa Pérez-Bueno, Mónica Pineda, and Matilde Barón. Phenotyping plant responses to biotic stress by chlorophyll fluorescence imaging. *Frontiers in Plant Science*, 10, 2019. ISSN 1664-462X. doi: 10.3389/fpls.2019.01135. URL http://dx.doi.org/10.3389/fpls.2019.01135.

[11] Anne-Katrin Mahlein. Plant disease detection by imaging sensors – parallels and specific demands for precision agriculture and plant phenotyping. *Plant Disease*, 100(2):241–251, 2016. ISSN 1943-7692. doi: 10.1094/pdis-03-15-0340-fe. URL http://dx.doi.org/10.1094/pdis-03-15-0340-fe.

[12] Claus Buschmann. Variability and application of the chlorophyll fluorescence emission ratio red/far-red of leaves. *Photosynthesis Research*, 92(2):261–271, 2007. ISSN 1573-5079. doi: 10.1007/s11120-007-9187-8. URL http://dx.doi.org/10.1007/s11120-007-9187-8.

[13] Nikkí A. Delk, Keith A. Johnson, Naweed I. Chowdhury, and Janet Braam. *CML24* , regulated in expression by diverse stimuli, encodes a potential ca2+ sensor that functions in responses to abscisic acid, daylength, and ion stress. *Plant Physiology*, 139(1):240–253, 2005. ISSN 0032-0889. doi: 10.1104/pp.105.062612. URL http://dx.doi.org/10.1104/pp.105.062612.

[14] Wei Ma, Andries Smigel, Yu-Chang Tsai, Janet Braam, and Gerald A. Berkowitz. Innate immunity signaling: Cytosolic ca2+ elevation is linked to downstream nitric oxide generation through the action of calmodulin or a calmodulin-like protein. *Plant Physiology*, 148(2): 818–828, 2008. ISSN 1532-2548. doi: 10.1104/pp.108.125104. URL http://dx.doi.org/10.1104/pp.108.125104.

[15] Howland Jones. Arabidopsis imaging, uw arabidopsis (msv091416). Unpublished technical report (slides), Middleton Spectral Vision, 2016. Analysis of the July 2016 experiment; not publicly available.

[16] Anna de Juan and Romà Tauler. Multivariate curve resolution: 50 years addressing the mixture analysis problem – a review. *Analytica Chimica Acta*, 1145:59–78, 2021. ISSN 0003-2670. doi: 10.1016/j.aca.2020.10.051. URL http://dx.doi.org/10.1016/j.aca.2020.10.051.

[17] Yasuhiro Ishiga, Takako Ishiga, Srinivasa R Uppalapati, and Kirankumar S Mysore. Arabidopsis seedling flood-inoculation technique: a rapid and reliable assay for studying plant-bacterial interactions. *Plant Methods*, 7(1), 2011. ISSN 1746-4811. doi: 10.1186/1746-4811-7-32. URL http://dx.doi.org/10.1186/1746-4811-7-32.

[18] Yoav Benjamini and Yosef Hochberg. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 57(1):289–300, 1995. ISSN 1467-9868. doi: 10.1111/j.2517-6161.1995.tb02031.x. URL http://dx.doi.org/10.1111/j.2517-6161.1995.tb02031.x.

[19] Howland Jones. Arabidopsis imaging, uw arabidopsis, october experiment. Unpublished technical report, Middleton Spectral Vision, 2017. Analysis of the November 2016 replicate; not publicly available.

[20] Evangelia A. Papoutsoglou, Daniel Faria, Daniel Arend, Elizabeth Arnaud, Ioannis N. Athanasiadis, Inês Chaves, Frederik Coppens, Guillaume Cornut, Bruno V. Costa, Hanna Ćwiek Kupczyńska, Bert Droesbeke, Richard Finkers, Kristina Gruden, Astrid Junker, Graham J. King, Paweł Krajewski, Matthias Lange, Marie-Angélique Laporte, Célia Michotey, Markus Oppermann, Richard Ostler, Hendrik Poorter, Ricardo Ramírez-Gonzalez, Živa Ramšak, Jochen C. Reif, Philippe Rocca-Serra, Susanna-Assunta Sansone, Uwe Scholz, François Tardieu, Cristobal Uauy, Björn Usadel, Richard G. F. Visser, Stephan Weise, Paul J. Kersey, Célia M. Miguel, Anne-Françoise Adam-Blondon, and Cyril Pommier. Enabling reusability of plant phenomic datasets with miappe 1.1. *New Phytologist*, 227(1):260–273, 2020. ISSN 1469-8137. doi: 10.1111/nph.16544. URL http://dx.doi.org/10.1111/nph.16544.

[21] Mark D. Wilkinson, Michel Dumontier, IJsbrand Jan Aalbersberg, Gabrielle Appleton, Myles Axton, Arie Baak, Niklas Blomberg, Jan-Willem Boiten, Luiz Bonino da Silva Santos, Philip E. Bourne, Jildau Bouwman, Anthony J. Brookes, Tim Clark, Mercè Crosas, Ingrid Dillo, Olivier Dumon, Scott Edmunds, Chris T. Evelo, Richard Finkers, Alejandra Gonzalez-Beltran, Alasdair J.G. Gray, Paul Groth, Carole Goble, Jeffrey S. Grethe, Jaap Heringa, Peter A.C 't Hoen, Rob Hooft, Tobias Kuhn, Ruben Kok, Joost Kok, Scott J. Lusher, Maryann E. Martone, Albert Mons, Abel L. Packer, Bengt Persson, Philippe Rocca-Serra, Marco Roos, Rene van Schaik, Susanna-Assunta Sansone, Erik Schultes, Thierry Sengstag, Ted Slater, George Strawn, Morris A. Swertz, Mark Thompson, Johan van der Lei, Erik van Mulligen, Jan Velterop, Andra Waagmeester, Peter Wittenburg, Katherine Wolstencroft, Jun Zhao, and Barend Mons. The fair guiding principles for scientific data management and stewardship. *Scientific Data*, 3(1), 2016. ISSN 2052-4463. doi: 10.1038/sdata.2016.18. URL http://dx.doi.org/10.1038/sdata.2016.18.