Fluorescence hyperspectral signatures of *Pseudomonas* challenge and flg22 in *Arabidopsis*

A FAIR reanalysis of the 2016 UW-Madison experiment

Richard J. Barker [PENDING: co-authors]

[PENDING: date / venue]

The question

- Can we **see** a plant responding to bacteria before symptoms appear?
- Plants perceive flagellin (flg22) through the receptor **FLS2**; *fls2* mutants cannot.
- **CML24** (TCH2) is a Ca^{2+} sensor; *cml24-4* is impaired in pattern-induced NO signalling.
- Chlorophyll fluorescence changes with photosynthesis and chlorophyll content, so a **whole emission spectrum per pixel** might carry an early signature.

Key control: a genuine flg22 *perception* signature must vanish in *fls2*.


The 2016 experiment

- MacroPhor fluorescence hyperspectral imager (Middleton Spectral Vision), Gilroy Lab
- 488 nm excitation, 500–800 nm emission, 324 bands
- 5 genotypes per plate: Col-0, *fls2-17*, *fls2-101*, *cml24-2*, *cml24-4*
- Mock, flg22 or *P. syringae*; 4 plates each
- Imaged before treatment, then 1, 2, 3, 24 and 48 h
- Independent replicate (Nov 2016, 405 nm excitation)

Recovered from the drive


7,752 files, 396 GB
400 image cubes
2,486 unique hand-drawn seedling spectra
every file name parsed, every cube hashed

Data problem 1: exposure time changed between timepoints




The 2016 graphs' 21% intensity fall after treatment is a camera setting: after dividing by exposure, 1–3 h Control values lie within 4% of hour 0.

Data problem 2: plates differed *before* treatment




(a) At hour 0, before any treatment, flg22 and Control plates are perfectly separable (AUC 1.00). Only change from each plate's own baseline is evidence (b).

A pathogen signature within 1–2 hours



Red emission peak broadens by 1.41 nm at 1 h ($q = 0.045$) and 1.77 nm at 2 h ($q = 0.001$); emission below 640 nm rises (2 h, $q = 0.005$).

What the spectrum change looks like




Area-normalised spectra, change from hour 0, treated minus Control (Col-0).

Infected plates can be told apart from their spectra

- Leave-one-plate-out classifier, change-from-baseline spectra (Col-0)
- Plate-level AUC 1.00 at 1 h, 1.00 at 2 h, 1.00 at 24 h
- Exact permutation test over all 70 plate relabellings: $p = 0.029$
- Fails at 3 h [PENDING: explain]

The flg22 signal is not FLS2-dependent



A Col-0-trained flg22 model fires as strongly in *fls2-17* (0.36) and *fls2-101* (0.20) as in Col-0 (0.21): a plate-level effect, not perception. The pathogen signature is in every genotype, as expected.

The November replicate

- 405 nm excitation; Col-0, *fls2-17*, *fls2-101*; 0, 6, 24, 48 h
- Hand-drawn ROIs exist for only one plate per treatment, so treatment effects are confounded with plate there
- Within plates: flg22 shifts Col-0 emission shape differently from *fls2-17* at 6 h ($q = 0.026$), the FLS2-dependent pattern
- [PENDING: plate-replicated analysis from the raw cubes]

Take-home

1. Carry acquisition metadata with every intensity: one camera setting made the most striking 2016 pattern.
2. The plate is the experimental unit: image a baseline and validate leave-plate-out.
3. Use non-perceiving mutants as the decisive control.
4. A pathogen signature appears within 1–2 h: a broader red peak and more emission below 640 nm.

FAIR resource

- Code, manifests, hand-ROI spectra, derived tables: `github.com/dr-richard-barker/PathoFluor-At`
- Raw cubes (lossless, SHA-256 verified): Zenodo [PENDING: DOIs]
- Project page: `dr-richard-barker.github.io/PathoFluor-At`
- Every number in the paper and these slides is generated from the committed tables