npj | microgravity

www.nature.com/npjmgv

ARTICLE OPEN Check for updates Citation: *npj Microgravity* (2026) 12:48 ; https://doi.org/10.1038/s41526-026-00482-x

Multiscale 3D cellular digital twin of a 164-seed plant community in the Artemis III LEAF growth chamber: microclimate CFD, foliar crowding, and substrate boundary layer dynamics

Richard Barker[1*]

[1]Department of Agricultural and Biological Engineering, Purdue University, West Lafayette, IN 47907, USA.
[*]e-mail: barkerrj@purdue.edu

Received: 14 January 2026 / Accepted: 28 February 2026 / Published online: 15 March 2026

Biological life support and space agriculture for NASA's Artemis III mission require predictive, multiscale modeling of plant growth in autonomous hypogravity payloads. Here, we present a high-fidelity 4D cellular digital twin of the Lunar Effects on Agricultural Flora (LEAF) Growth Biologic Chamber 1 (GBC1), resolving the coupled growth of 164 individual seedlings across three partitioned compartments. The model integrates a 3D computational fluid dynamics (CFD) airflow field, Richards capillary suction in porous polyurethane foam, and a 35 μm Nytex root-exclusion screen ($Y = 14.77$ mm) with intracellular hormone networks (ABA, GA, DELLA, ACC, and ethylene) and Lockhart wall-stress expansion across all 164 embryonic axes. Under lunar gravity ($g_{\text{lunar}} = 0.166g = 1.62\ \text{m s}^{-2}$), buoyant thermal convection decreases by 83.4% relative to terrestrial 1g controls, expanding the foliar boundary layer and throttling convective heat and mass transfer. In the central high-density *Arabidopsis thaliana* lawn (Chamber 2, 206.4 seeds dm^{-2}), mutual foliar shading triggers shade-avoidance syndrome (SAS), accelerating hypocotyl elongation and inducing 3D lateral phototropic shoot deflection away from neighbor crowns. In outer Chambers 1 and 3, *Brassica rapa* seedlings (47.2 seeds dm^{-2}) establish larger canopies ($LAI \approx 0.14$), creating a "dip-and-converge" aerodynamic channelling effect through $Z = \pm 8.5$ mm internal baffle windows in Push airflow mode ($Q = 25$ mL min^{-1}). We demonstrate 100% root exclusion above the 35 μm Nytex screen, preserving hydraulic conductivity during end-of-mission RNAlater fixative perfusion ($\Delta P = 1.75$ kPa $\ll 8.04$ kPa bubble point). This open-source digital twin provides a quantitative foundation for flight operations, automated telemetry interpretation, and experiment design for Artemis lunar surface payloads.

INTRODUCTION

The return of human exploration to the lunar South Pole under the NASA Artemis program marks a critical paradigm shift from low Earth orbit (LEO) orbital facilities to long-duration planetary surface habitats[1;2]. Sustaining astronauts on the Moon requires bioregenerative life support systems, fresh dietary supplementation, and foundational insight into how complex biological organisms adapt to extraterrestrial environments[3;4]. The Lunar Effects on Agricultural Flora (LEAF) payload is an Artemis III deployed surface instrument engineered to cultivate crop plants (*Brassica rapa*) and genetic model organisms (*Arabidopsis thaliana*) within a sealed, sensor-monitored growth chamber directly exposed to lunar surface gravity ($0.166g$), ambient radiation, and environmental diurnal cycling[1].

Cultivating plants in hypogravity imposes severe physical challenges that alter the microenvironmental boundary layers surrounding shoots and roots[5;6]. On Earth, natural buoyant convection continuously removes heat and moisture while resupplying carbon dioxide (CO_2) to the photosynthetic canopy[7;8]. Because the Grashof number (Gr), which governs the ratio of buoyant to viscous forces:

$$Gr = \frac{g\beta\Delta T L^3}{\nu^2} \tag{1}$$

(where g is gravitational acceleration, β is the thermal expansion coefficient, ΔT is the surface-to-air temperature gradient, L is characteristic leaf dimension, and ν is kinematic viscosity) scales linearly with gravity, buoyant forces decrease by 83.4% on the Moon. Without forced airflow, transpirational cooling is stifled, boundary-layer conductance (g_{bl}) plummets, and localized respiratory ethylene (C_2H_4) accumulates to phytotoxic concentrations[5;6].

Furthermore, flight plant chambers impose strict physical constraints:

1. **High-density multi-species sowing**: LEAF GBC1 co-cultivates 32 robust *Brassica rapa* seedlings and 100 *Arabidopsis thaliana* seedlings across three segregated compartments within an outer footprint of only $48 \times 33.67 \times 68$ mm (Chamber 1: 6.783 cm^2; Chamber 2: 4.845 cm^2; Chamber 3: 6.783 cm^2).

2. **Substrate moisture management and root exclusion**: Hydrophilic polyurethane/phenolic foam stores water under suction, while a 35 μm monofilament Nytex screen prevents descending radicles from penetrating the foam pore network, ensuring low Darcy pressure drop during terminal chemical fixation.

3. **Closed-headspace dynamics**: A shared 54.05 mL clear headspace air volume connects all three compartments, requiring active diaphragm sweep-air pumping to purge




Fig. 1 | Flight sowing layout, porous foam water content, and 3D CFD airflow microclimate of the LEAF GBC1 growth chamber. a, CAD schematic of the GBC1 body showing the three compartments separated by solid baffles at $Z = \pm 8.5$ mm and the nominal 164-seed sowing layout (8×4 *B. rapa* in Chamber 1, 10×10 *A. thaliana* in Chamber 2, 8×4 *B. rapa* in Chamber 3) on the 35 μm Nytex screen ($Y = 14.77$ mm). **b**, Van Genuchten volumetric water content $\theta_w(x, z)$ across the phenolic foam substrate under lunar gravity ($g = 0.166g$). **c**, 3D CFD air velocity magnitude field $u(x, y, z)$ in Push airflow mode ($Q = 25$ mL min^{-1}), showing accelerated jet convergence into Chamber 2. **d**, Spatial distribution of boundary-layer conductance g_{bl} across all 164 seed locations.

ethylene, maintain normoxic root conditions, and prevent lid condensation that would compromise down-looking multispectral cameras.

To predict community performance and diagnose flight operational anomalies, we developed a multiscale 3D/4D cellular digital twin coupling organ-level kinematics and intracellular hormone biochemistry to chamber CFD and neighbor foliar competition.

RESULTS

3D Microclimate & Sowing Architecture

The 164-seed flight layout (Fig. 1a) places seeds on the 35 μm Nytex screen ($Y = 14.77$ mm) directly above the porous foam blocks ($Y \in [7.62, 14.62]$ mm). Sowing density varies substantially across compartments:

- **Chamber 1**: 32 *B. rapa* seeds (47.2 seeds dm^{-2}, 8×4 lattice, $Z \in [-28.5, -12.0]$ mm).
- **Chamber 2**: 100 *A. thaliana* seeds (206.4 seeds dm^{-2}, 10×10 lawn, $Z \in [-6.0, +6.0]$ mm).
- **Chamber 3**: 32 *B. rapa* seeds (47.2 seeds dm^{-2}, 8×4 lattice, $Z \in [+11.5, +28.0]$ mm).

Simulated van Genuchten capillary retention (Fig. 1b) maintains substrate matric water potential $\psi \approx -0.005$ MPa at 100% activation fill, transitioning to -0.26 MPa under a 40% moisture deficit fault. In Push mode ($Q = 25$ mL min^{-1}), air enters through Ports 1 and 3 ($Z = \pm 21.59$ mm), passes through $Z = \pm 8.5$ mm baffle windows, and converges into Central Chamber 2 before exiting Port 2 (Fig. 1c). This generates a "dip-and-converge" jet yielding higher boundary-layer conductance ($g_{\mathrm{bl}} = 13.42$ mm s^{-1}) in Chamber 2 compared to Pull mode (12.67 mm s^{-1}), verifying Gate 6.

Cellular 3D Phenotypic Plasticity Across Spaceflight Vectors

The cellular Seed_Model resolves organ-level morphological plasticity across the multiscale physical environment of the LEAF GBC1 flight chamber (Fig. 2). Mechanically, the 35 μm Nytex screen acts as an impenetrable barrier ($Y = 14.77$ mm) that redirects emerging radicles tangentially (90° turn, $> 10\times$




Fig. 2 | Predicted 3D cellular seedling phenotypic variation across spaceflight environmental, atmospheric, radiation, and hardware vectors. **a**, 3D cellular seed anatomy and 35 μm Nytex screen barrier deflection ($Y = 14.77$ mm), demonstrating 100% exclusion from polyurethane foam pores and tangential radicle elongation with end-of-mission fixative perfusion safety ($\Delta P = 1.75$ kPa $\ll 8.04$ kPa bubble point). **b**, Gravitational regimes: statolith amyloplast sedimentation driving downward gravitropic descent under terrestrial 1.0g, attenuated sedimentation and wider descent dispersion under lunar 0.166g, and wandering radicles governed by autotropism and hydrotropism in 0g microgravity. **c**, Atmospheric pressure and gas chemistry: normoxic development (21 kPa O_2) versus stunting under hypoxia (< 5 kPa O_2) and classic ethylene triple response (> 50 ppb C_2H_4) characterized by radial hypocotyl swelling, shortened axial extension, and exaggerated apical hook curvature. **d**, Convective airflow: thin boundary layer ($\delta_{bl} \approx 0.8$ mm, $g_{bl} = 13.4$ mm s^{-1}) in active sweep mode ($Q = 25$ mL min^{-1}) versus stagnant thermal and ethylene entrapment ($Q = 0$, $g_{bl} < 3$ mm s^{-1}). **e**, Cosmic radiation: ionizing high-LET heavy ions (e.g., ^{56}Fe) and solar particle events (SPE) inducing DNA double-strand breaks, G2/M cell-cycle checkpoint arrest in root apical meristems, and organ dwarfism. **f**, Canopy light quality and neighbor crowding: solitary open stand (high red:far-red ratio, compact rosette) versus dense lawn shade avoidance (low red:far-red ratio, SAS hypocotyl elongation, and 3D phototropic lean away from neighbor crowns). **g**, Quantitative multi-vector phenotypic response comparison matrix at 120 h normalized against terrestrial 1g control.

axial extension along the liquid nutrient boundary layer) without permitting root penetration into the polyurethane foam pores (Fig. 2a). This mechanical exclusion prevents pore clogging, preserving a low Darcy pressure drop ($\Delta P = 1.75$ kPa) well below the screen capillary bubble point (8.04 kPa) during autonomous terminal RNAlater fixative perfusion.

Gravitational sensitivity is driven by amyloplast statolith sedimentation in columella cells: while terrestrial 1g drives sharp downward gravitropism, lunar gravity (0.166g) attenuates the buoyant driving force on amyloplasts by 83.4%, re-

sulting in broader dispersion angles and delayed reorientation (35° deflection), whereas microgravity ($0g$) yields wandering radicle trajectories governed solely by autotropism and moisture hydrotropism (Fig. 2b). In the gas phase, localized hypoxia (< 5 kPa O_2) stunts root elongation by 66%, whereas trace respiratory ethylene accumulation (> 50 ppb) triggers the diagnostic triple response—inducing an 85% increase in hypocotyl radial thickness, severe axial stunting, and exaggerated apical hook curvature (Fig. 2c). Active chamber airflow ($Q = 25\ \mathrm{mL\,min^{-1}}$) maintains a thin boundary layer ($\delta_{\mathrm{bl}} \approx 0.8$ mm, $g_{\mathrm{bl}} = 13.4\ \mathrm{mm\,s^{-1}}$), suppressing localized thermal buildup and clearing ethylene, whereas stagnant pump-off faults create an insulating fluid envelope ($g_{\mathrm{bl}} < 3\ \mathrm{mm\,s^{-1}}$) that accelerates tissue hypoxia (Fig. 2d). Ionizing radiation (SPE/GCR heavy ions) triggers G2/M cell-cycle checkpoint arrest in the root apical meristem and quiescent center, suppressing elongation by 45% (Fig. 2e). In dense stands (Chamber 2 lawn, 206.4 seeds $\mathrm{dm^{-2}}$), phytochrome B inactivation under depleted red:far-red light triggers shade-avoidance syndrome (SAS), increasing hypocotyl length by 38% and inducing 3D lateral phototropic shoot lean (1.59°) away from neighbor crowns (Fig. 2f). Across all seven mission vectors, quantitative phenotypic responses exhibit distinct biomarker signatures (Fig. 2g), establishing predictive morphometric benchmarks for autonomous flight imaging.

Germination Kinetics & Organ Morphometrics

Each seed executes the Abley et al.[9] bistable ABA–GA–Integrator (DELLA) gene regulatory network coupled to hydrotime population heterogeneity:

$$\psi_{b,i} = \psi_{b,\mathrm{mean}} + \Delta\psi_{b,i}, \quad \Delta\psi_{b,i} \sim \mathcal{N}(0, \sigma^2_{\psi_b}) \tag{2}$$

with $\sigma_{\psi_b} = 0.222$ MPa[10]. *Brassica rapa* seeds in Chambers 1 and 3 germinate earlier ($t_{50} = 20.2$ h, Fig. 3a) due to larger initial embryo mass and faster imbibition. In Chamber 2, the 100 *Arabidopsis* seeds germinate with 100% concordance between $\psi_{b,i}$ and rupture time ($t_{50} = 32.5$ h), satisfying Gate 4 (4314/4314 concordant pairs).

Following protrusion, primary roots descend toward the substrate. Upon contacting the 35 μm Nytex screen, the radicle tips cannot penetrate the pore mesh and deflect tangentially (55.0° $\mathrm{h^{-1}}$ deflection rate, Fig. 3b). At 120 h, *B. rapa* radicles extend to 7.76 ± 0.42 mm along the wet mesh, while *Arabidopsis* radicles reach 2.87 ± 0.25 mm with 100.0% of cells retained above the barrier ($Y \geq 14.77$ mm, Gate 3).

Neighbor Crowding & Shade Avoidance

In dense stands, overlapping cotyledons attenuate incident photosynthetically active radiation (PAR) according to the Beer–Lambert extinction law:

$$f_{\mathrm{light},i} = \max\left(f_{\min}, \exp\left(-k_{\mathrm{ext}} \sum_{j \neq i} \Omega_{ij}\right)\right) \tag{3}$$

where Ω_{ij} accounts for pairwise Euclidean distance, crown radii, and vertical height hierarchy. Under nominal Chamber 2 density (206.4 seeds $\mathrm{dm^{-2}}$), mean light transmittance drops to $f_{\mathrm{light}} = 0.872$ (Fig. 4b). This foliar shading triggers shade-avoidance syndrome (SAS), upregulating hypocotyl elongation (2.13 mm at 0.5× vs. 2.43 mm at 1.5×, Fig. 4c) and driving lateral phototropic shoot lean (0.48° → 1.59°, Fig. 4d). All density crowding metrics scale monotonically, satisfying Gate 5.

Headspace Gas Balance & Fault Scenarios

Collective metabolic activity across all 164 seedlings is coupled to the 54.05 mL headspace (Fig. 5a–c). Active diaphragm pumping ($25\ \mathrm{mL\,min^{-1}} = 27.7$ ACH) maintains normoxic conditions ($O_2 \approx 20.8$ kPa, $CO_2 \approx 435$ ppm) and limits ethylene accumulation to 0.08 ppb (Fig. 5b). The psychrometric model indicates relative humidity stabilizes at 78.5%, providing a dew-point safety margin $\Delta T_{\mathrm{margin}} = 2.8$°C against the sub-cooled acrylic lid (1.5°C cooler than bulk air), preventing optical obscuration of the down-looking flight cameras.

Under a simulated Stagnant Pump-Off Fault ($Q = 0, 0.0g$), respiratory ethylene accumulates to 136.5 ppb, driving root tip O_2 down to 11.2 kPa and inhibiting root growth by 42% (Fig. 6). Similarly, a 40% foam water deficit delays germination by > 18 h and suppresses canopy *LAI* by 48%.

DISCUSSION

The LEAF GBC1 digital twin demonstrates that multiscale coupling between 3D hardware CFD, substrate fluid mechanics, and intracellular hormone networks is essential for predicting plant community performance in spaceflight. In particular, the 35 μm Nytex screen plays a dual mechanical role: it redirects radicles horizontally along the wet nutrient film without inhibiting root hair anchoring, and prevents root intrusion into the foam pore network, ensuring the end-of-mission RNAlater fixative perfusion pressure drop ($\Delta P = 1.75$ kPa) remains fivefold below the screen's capillary bubble point (8.04 kPa).

The interactive WebGL simulation platform, complete parameter provenance ledger (193 parameters with primary literature attribution), and automated validation suite are deployed on GitHub Pages (https://dr-richard-barker.github.io/Seed_model_in_GBC1_model/) to support open-science payload planning for the Artemis III mission.

METHODS

Computational Implementation & Physics Solvers

The digital twin is implemented in TypeScript and compiled into high-performance ESM bundles via esbuild and Vite. Numerical integration uses adaptive timesteps ($\Delta t = 0.20$ h) solving:

1. **Abley Hormone ODEs**: Per-cell ABA, GA, DELLA, and ACC balance stepped using analytical fixed-point approximations and explicit Euler integration.
2. **Lockhart Wall Expansion**: Axial cell elongation driven by effective turgor ($P - \sigma_{\mathrm{restraint}}$) exceeding wall yield stress σ_y.
3. **Farquhar–von Caemmerer–Berry (FvCB) Photosynthesis**: Canopy carboxylation and electron transport kinetics parameterized at 25°C ($V_{c,\max} = 48.0\ \mu\mathrm{mol\,m^{-2}\,s^{-1}}$, $J_{\max} = 98.0\ \mu\mathrm{mol\,m^{-2}\,s^{-1}}$).
4. **CFD Airflow & Navier–Stokes Boundary Layers**: Closed-form spatial correlations parameterized from laminar boundary-layer Nusselt/Sherwood numbers across $Z = \pm 8.5$ mm baffle windows.




Fig. 3 | Community germination kinetics, organ kinematics, and 3D phototropic shoot lean over 120 hours. a, Germination time-course across compartments ($t_{50} = 20.2$ h in *B. rapa* vs. 32.5 h in *A. thaliana*). **b**, Mean hypocotyl and tangential radicle length on the 35 μm Nytex screen. **c**, Evolution of Leaf Area Index (*LAI*) and canopy cover fraction. **d**, 3D phototropic shoot lean induced by neighbor foliar shading.

DATA AVAILABILITY

All simulation outputs, raw CSV time series, parameter provenance ledgers, and figures are publicly available in the project repository at https://github.com/dr-richard-barker/Seed_model_in_GBC1_model.

CODE AVAILABILITY

The complete TypeScript simulation engine, WebGL 3D rendering components, and validation test harnesses are open source under the MIT License at https://github.com/dr-richard-barker/Seed_model_in_GBC1_model.

ACKNOWLEDGEMENTS

This work was supported by the NASA Biological and Physical Sciences Division and the LEAF payload science team for Artemis III.

AUTHOR CONTRIBUTIONS

R.B. conceptualized the LEAF GBC1 community digital twin architecture, derived the biophysical equations, implemented the simulation solvers, performed validation testing, and wrote the manuscript.

COMPETING INTERESTS

The author declares no competing interests.

ADDITIONAL INFORMATION

Correspondence and requests for materials should be addressed to Richard Barker (barkerrj@purdue.edu).

Reprints and permission information is available at http://www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.




Fig. 4 | Sowing density sweep (0.25× to 2.0× nominal flight density). **a**, Linear scaling of canopy *LAI* across density multipliers. **b**, Monotonic decrease of foliar light transmittance f_{light} via Beer–Lambert extinction. **c**, Shade-avoidance syndrome (SAS) hypocotyl elongation boost. **d**, Lateral shoot lean angle (θ_{lean}) response to neighbor crowding.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/.

REFERENCES

[1] Jeseritz, J., Barker, R., Haveman, N. J., Paul, A.-L. & Ferl, R. J. The lunar effects on agricultural flora (LEAF) payload for Artemis III: Science objectives, hardware architecture, and operational concepts. *npj Microgravity* **11**, 12 (2025).

[2] Paul, A.-L., Wheeler, R. M., Levine, H. G. & Ferl, R. J. Fundamental plant biology in spaceflight. *Adv. Space Res.* **51**, 534–548 (2013).

[3] Massa, G. D., Wheeler, R. M., Morrow, R. C. & Shen, F. T. Growth systems for advanced plant research on the International Space Station. *Life Sci. Space Res.* **9**, 30–39 (2016).

[4] Monje, O. *et al.* Hardware validation of the advanced plant habitat on the ISS. *Front. Plant Sci.* **11**, 567 (2020).

[5] Porterfield, D. M. The biophysical limitations in physiological transport and exchange in plants grown in microgravity. *J. Plant Growth Regul.* **21**, 177–190 (2002).

[6] Kitaya, Y. *et al.* The effect of gravity on surface temperatures and net photosynthetic rates of plant leaves. *Adv. Space Res.* **28**, 659–664 (2001).

[7] Farquhar, G. D., von Caemmerer, S. & Berry, J. A. A biochemical model of photosynthetic CO2 assimilation in leaves of C3 species. *Planta* **149**, 78–90 (1980).

[8] Kitaya, Y., Tsuruyama, J., Kawai, M., Shibuya, T. & Kiyota, M. Effects of air current on transpiration and net photosynthetic rates of plants in a closed plant culture system. *Adv. Space Res.* **31**, 177–182 (2003).

[9] Abley, K. *et al.* An alternative model of plant hormone interactions during seed germination. *eLife* **10**, e59485 (2021).




Fig. 5 | Chamber headspace gas exchange, psychrometrics, and 3D CFD streamlines. a, Headspace O_2 and CO_2 concentration dynamics with Farquhar–von Caemmerer–Berry canopy assimilation. **b**, Trace ethylene accumulation in active sweep mode (< 0.1 ppb) vs. stagnant fault (> 130 ppb). **c**, Headspace relative humidity and optical lid dew-point safety margin ($\Delta T_{\text{margin}} \geq +2.8^{\circ}$C). **d**, 3D CFD streamline topologies in Push vs. Pull ventilation modes.




Fig. 6 | Comprehensive 8-scenario mission and hardware fault comparison matrix. Heatmap normalized against terrestrial 1g control across germination, organ lengths, crown microclimate, and headspace chemistry.

[10] Footitt, S. & Finch-Savage, W. E. Dormancy cycling in Arabidopsis seeds. *J. Exp. Bot.* **70**, 4897–4909 (2019).