

From a procedural fork to a cited resource: building a FAIR, literature-grounded *Arabidopsis thaliana* atlas from an uncredentialed rice viewer

Richard J. Barker*

September 18, 2026

Abstract

`rice-atlas` is an interactive 3D viewer that lets a user select any of several hundred organs of a procedurally-generated rice plant. Its own documentation states plainly that it is a conceptual educational model, not a scan of a specimen, and it cites no *Oryza sativa* dataset, carries no FAIR metadata, and had not been reviewed. We report what it takes to turn that pattern into a genuinely citable scientific resource for a second species, *Arabidopsis thaliana*, rather than simply relabelling the same geometry. We rebuild the organ-selectable viewer from a clean-room implementation – `rice-atlas`’s own license is ambiguous (GitHub records it as `NOASSERTION`) – and ground it in three places real data was available: organ geometry parameters cite peer-reviewed developmental-biology sources (3 of 5 organs carry an explicit geometry citation); developmental gene expression is linked, not redistributed, to the real 79-organ Klepikova/TraVA atlas, whose bulk-reuse terms we could not confirm; and a spaceflight-response layer – absent from `rice-atlas` entirely – is computed directly from real NASA Open Science Data Repository (OSDR) count matrices for root tissue (OSD-120: 32,833 genes, 18 flight vs. 18 ground samples, top flight/ground log2 ratio 4.75 for `AT2G01422`) and whole seedlings under simulated and real microgravity (OSD-314: 5 microgravity vs. 6 1g samples, top ratio 5.54 for `AT4G28520`). We report this as a mean-CPM ratio, explicitly not a statistically tested differential-expression call, and say so everywhere the numbers appear. A second pass rebuilds the viewer’s geometry on the same spline-swept tube construction `rice-atlas` itself actually uses (found by reading its source, not its file names) and wires the plant’s shape to real per-ecotype trait data: 5 ecotypes (`Col-0`, `Ler`, `Ws`, `Tsu-0`, `Edi-0`) carry directly-computed rosette-shape descriptors from a paper’s own raw published data (mean compactness 0.6637 for `Col-0` vs. 0.7617 for `Ler`, independently confirming `Ler`’s known compact habit by an entirely different method than the genetics literature that first described it), `Ler`’s short pedicels and blunt siliques follow the real natural *erecta* mutation, and `Cvi-0` varies by a directly-quoted real leaf-count difference. The result is graded against our own portfolio’s evidence gate before publication, and every claim in this manuscript traces to a file this repository’s own scripts produce. The main finding is procedural: an appealing interactive demo and a FAIR scientific resource are different amounts of work, and the gap between them is exactly the set of citations, provenance notes, and honest limitations this atlas adds.

1 Introduction

Interactive 3D anatomy viewers are an appealing format for plant biology outreach and education: a user rotates a whole organism, clicks an organ, and reads what it is. The format’s origin, however, matters for whether what they read is true. `rice-atlas` (<https://github.com/dr-richard-barker/rice-atlas>) is one such viewer, built on a procedural functional-structural plant model and forked twice from a generic human-anatomy template before being relabelled for

*Center of Space Exploration (CoSE). Correspondence: dr.richard.barker@gmail.com

rice. Its own README is explicit that the geometry is conceptual, and inspecting the repository directly turns up no cited *Oryza sativa* dataset, no license the GitHub license detector can resolve (recorded as NOASSERTION), no FAIR metadata, and no GitHub Pages deployment.

None of that makes the viewer worthless – a clearly-labelled conceptual model is a legitimate teaching tool. It does mean that building "the *Arabidopsis* version" of it is an underspecified request until a decision is made about which of two very different projects is wanted: a relabelled geometric demo, or a resource that earns the adjectives this project set out with – FAIR, reviewed, and scientifically accurate. We chose, and report here, a hybrid: keep the interactive organ-selectable viewer, because that is the part of rice-atlas actually worth learning from, but rebuild its content so that every organ's geometry and every data value shown traces to a real, checkable source.

Arabidopsis thaliana is a fortunate choice of second species for this exercise. Unlike rice, it has a 79-organ, whole-development RNA-seq atlas [1], single-cell developmental atlases for at least two organ systems [2, 3], and – because it is the standard model organism for plant spaceflight biology – a substantial NASA Open Science Data Repository (OSDR) footprint, including studies that profile root and whole-seedling transcriptomes under real and simulated microgravity. That last point gives the resulting atlas a dimension rice-atlas has no counterpart for at all: a spaceflight-response layer, grounded in real flight experiments rather than invented.

We report three things: what we found rice-atlas actually to be, once inspected rather than taken at face value; what real data sources were available to ground an *Arabidopsis* equivalent, and which ones we could and could not actually use; and what building to a pre-registered FAIR/evidence-gate standard changed about the result compared to simply forking the original.

2 Methods

2.1 Characterizing rice-atlas

We inspected `dr-richard-barker/rice-atlas` via the GitHub API and `gh` CLI (repository metadata, full file tree, Pages configuration, and commit history) rather than assuming its content from its name or description. This established that it is a fork (itself forked from a human-anatomy viewer template) with no commits from the account that owns it, no `data/` directory, no cited primary dataset, and a license the GitHub license detector could not resolve to a standard SPDX identifier.

2.2 Provenance discipline

Every external fact used in this repository is recorded in `data/README.md` with its source, verification method, and license status before being used anywhere else. Three literature DOIs (1–3) were each confirmed against the CrossRef API (title, full author list, year, volume, pages) rather than trusted from a search snippet. Four NASA OSDR study accessions were confirmed live against the OSDR metadata API before use; two (OSD-120, OSD-314) had already-fetched raw count matrices available (reused, with permission implicit in common ownership, from the sibling repository `arabidopsis-drem-osdr`, itself sourced from the same public, CC0 OSDR accessions) and were used quantitatively; two (OSD-38, OSD-522) are cited as confirmed, relevant, real studies but are not yet quantitatively integrated, and are documented as pending rather than fabricated.

2.3 Spaceflight response quantification

For each of the two studies with available raw counts, samples were split into their real experimental groups by the labels already present in the count matrix's own sample names (OSD-120: GC ground control vs. FLT flight, Day 13 root tissue; OSD-314: 1g vs. 0g, whole seedling). Raw counts

were converted to counts-per-million (CPM) per sample to correct for sequencing-depth differences, averaged within each group, and reported as $\log_2((\text{mean CPM}_{\text{treatment}} + 1)/(\text{mean CPM}_{\text{control}} + 1))$. This is deliberately a simple mean-ratio summary: no DESeq2/edgeR model was fit, no p -values or multiple-testing correction were computed, and no covariates (genotype, light treatment) were controlled for. The implementation is `scripts/01_compute_spaceflight_response.py`; its output feeds `scripts/02_export_viewer_data.py`, which derives the compact per-organ summary shown in the viewer, so no number reaching the viewer or this manuscript is hand-typed.

2.4 Structural model

Organ geometry for the viewer (root system, rosette leaf, inflorescence axis, flower, silique) is generated procedurally in the client (`app/src/geometry.ts`), not sculpted or scanned. Roots, stems, petioles and pedicels are built as tapering tubes swept along Catmull-Rom splines using a double-reflection rotation-minimizing frame builder (`app/src/organGeometry/tube.ts`) – independently implementing the same class of technique found by reading rice-atlas’s own source (Results), rather than `THREE.TubeGeometry`’s default Frenet frames, which visibly twist on tight or near-planar curves. Leaf blades and petals are a parametric curved surface (`app/src/organGeometry/leafBlade.ts`: a width profile along a midrib plus a cross-width fold) rather than a distorted sphere. Root branching topology and the rosette’s leaf-arrangement angle follow parameters attributed, organ by organ, to the literature sources in `app/src/organs.ts`; where a parameter is illustrative rather than measured, that organ’s on-screen description says so explicitly, matching the disclosure standard rice-atlas itself already sets for its own geometry.

2.5 Ecotype trait sourcing

Per-ecotype shape parameters (`app/src/ecotypes.ts`) come from three independent real sources rather than one uniform dataset, each disclosed as such in the application itself: (1) rosette compactness and size for Col-0, Ler, Ws, Tsu-0 and Edi-0 are direct per-accession means computed by `scripts/04_fetch_ecotype_traits.py` from Camargo et al. [2014]’s own raw published data (their Methods S2 supplementary archive, not a value read off a figure); (2) Ler’s short pedicels and blunt siliques follow the natural *erecta* mutation [5], with a quantitative proxy from Bundy et al. [6], explicitly labelled in-app as a proxy from an induced allele in a different genetic background rather than a direct measurement of natural Ler; (3) Cvi-0’s leaf count and thickness follow a direct quoted comparison in Coneva and Chitwood [7]. Every ecotype/trait combination without a real source found (e.g. Cvi-0’s rosette compactness, since it is not one of the Camargo et al. dataset’s 19 founder accessions) defaults to the Col-0 reference value and states so on screen, rather than interpolating or guessing one.

2.6 Evidence gate

Before this repository’s first push, and again before any push touching publication-relevant content, we ran this portfolio’s own pre-publication check (a blocklist sweep for previously-fabricated strings, CrossRef verification of every DOI, confirmation that every number traces to a producing script with no synthetic-data fallback, and confirmation that every cited file path and accession resolves in the actual repository). Only a check that was actually performed is recorded as an attestation; an attestation is never written to unblock a push it did not actually clear.

	rice-atlas	arabidopsis-atlas (this work)
Primary dataset cited	none found	3/5 organs cite a real structural paper
Spaceflight data	none	OSD-120, OSD-314 (real OSDR counts)
License (GitHub-detected)	NOASSERTION	MIT (this repository’s own)
FAIR metadata	none	<code>LICENSE</code> , <code>CITATION.cff</code> , <code>.zenodo.json</code>
GitHub Pages	not deployed	deployed
Independent review	none found	ABAI evidence-gate attestation on file

Table 1: Attributes compared directly by inspecting both repositories, not by description.

3 Results

3.1 What rice-atlas is, on inspection

3.2 Real spaceflight response, by organ

Of the five organs modelled, one (root) has a directly tissue-matched real spaceflight comparison: OSD-120 profiled root tissue specifically, at Day 13, under 18 flight and 18 ground control samples across 32,833 genes. The largest mean flight/ground CPM ratio in that comparison is $4.75 \log_2$ units, for AT2G01422.

OSD-314 profiled whole seedlings, not a single organ, across the same 32,833 genes, comparing 5 real microgravity samples against 6 1g ground samples; its largest ratio is $5.54 \log_2$ units, for AT4G28520. Because it is not organ-specific, we report it in the data provenance ledger rather than attach it to any single organ in the viewer, to avoid implying a tissue-level claim the data does not support.

The remaining three organs (rosette leaf, inflorescence axis, flower/silique) have no organ-matched spaceflight dataset wired into this version of the atlas; the viewer states this plainly for each rather than substituting the whole-seedling numbers as if they were organ-specific.

3.3 Realistic geometry and real ecotype shape-morphing

Inspecting rice-atlas’s actual source (not just its file names) showed that its visual quality comes mainly from spline-swept tube construction – a Catmull-Rom spline plus a parallel-transport (rotation-minimizing) frame builder – rather than from Blender sculpting, which it uses only for a thin Solidify-modifier pass on leaf blades. We re-implemented the same class of technique independently in the viewer itself (a double-reflection rotation-minimizing frame builder, `app/src/organGeometry/`), replacing the original primitive-shape organs with tapering, curved tube geometry and a parametric curved leaf-blade surface.

We then wired this geometry to real per-ecotype trait data so a viewer can select a natural accession and see the plant’s shape actually respond. 6 ecotypes are available; 5 of them (Col-0, Ler, Ws, Tsu-0, Edi-0) have real, directly-computed rosette-shape descriptors from Camargo et al. [2014]’s own published raw data (60 images per accession): mean rosette Compactness is 0.6637 for Col-0, 0.7617 for Ler, and 0.6729 for Ws – Ler-0 is the most compact and Tsu-0 the least compact of the five measured here, independently corroborating Ler’s known compact habit with an entirely different, image-based method than the genetic literature that first described it. Ler’s short pedicels and blunt siliques reflect the real, natural *erecta* (*er*) mutation [5]; a quantitative proxy from Bundy et al. [6] gives a mature wild-type pedicel of 7.90mm versus a $\sim 2.6\times$ shorter induced *er-105* pedicel, disclosed in the application itself as a proxy from an induced allele in a Columbia background, not a direct field measurement of natural Ler. Cvi-0, absent from the MAGIC founder panel, instead varies by a real, directly-quoted comparison from Coneva and Chitwood [7]: 2 more leaves than Ler in long-day conditions, with thicker leaf blades.

Every gap is disclosed rather than filled with a plausible value: Cvi-0’s rosette-compactness

parameter defaults to the Col-0 value because no measurement exists for it in the dataset used; ERECTA status for Cvi-0/Tsu-0/Edi-0 is stated as an assumed default, not confirmed per-accession; and no accession-labelled reference photo was found for Ws, so its photo link is simply absent rather than substituted.

3.4 Verification of the viewer itself

The viewer was checked interactively, not only by static code review: in both a local development server and the actual production build served at the live GitHub Pages URL, clicking each organ was confirmed to open an information panel showing the correct TraVA category, the correct geometry citation, and – for root – the real gene table described above, with no console errors in either environment.

4 Discussion

The central lesson of this project was not scientific but procedural: an appealing interactive 3D atlas and a FAIR scientific resource are not the same amount of work, and the difference is not visual polish. rice-atlas already had the harder-to-build part – a working, organ-selectable procedural plant viewer. What it lacked was cheaper in engineering terms but slower in practice: a cited dataset behind every organ, a resolvable license, and a record of what had and had not been checked. Recreating the visual experience for *Arabidopsis* took a fraction of the effort spent sourcing, verifying, and honestly scoping the data behind it.

That verification effort surfaced problems worth naming. A CrossRef check was needed because a plausible-looking DOI is not evidence it is correct. A live API check against NASA OSDR was needed because a study accession copied from memory is a guess, not a fact, until confirmed against the record NASA actually serves. And checking travadb.org’s actual interface – rather than assuming a REST-style URL pattern would work – avoided shipping a broken or misleading link in a resource whose entire premise is that its links and numbers are trustworthy.

The decision to compute only a simple mean-CPM ratio, rather than a full differential-expression model, was a scope choice made explicit rather than hidden: a half-rigorous statistical claim dressed as a rigorous one would be worse than a clearly labelled simple one. The same logic applied to organ coverage – three of five organs carry no spaceflight layer at all, and the viewer says so, rather than reusing the one available whole-seedling comparison across organs it was never measured in.

This pattern repeated when extending the atlas to ecotype shape-morphing: the visually appealing part (a plant that reshapes when you pick a different accession) was straightforward once the geometry was parametric; the slow part was again verification. The raw-data archive behind Camargo et al. [2014]’s rosette-shape descriptors turned out to be genuinely reusable – a real, column-labelled CSV, not just a figure to eyeball – but only after an explanatory table in the same paper’s own supplementary material (numbered, unlabelled columns, six example rows) turned out not to be it. The lesson generalizes: "supplementary data exists" and "supplementary data is the specific table you need, in the format you need" are different claims, and only checking the second one is real verification.

4.1 Limitations

No single real dataset gives whole-organism 3D geometry for *Arabidopsis*, just as none exists for rice; every organ shape here is a citation-informed illustrative approximation, not a reconstruction from measurement, and several structural parameters (rosette leaf-blade curvature, inflorescence taper, flower whorl spacing) are not yet tied to a specific measured value even where a general developmental-biology source is cited. TraVA/Klepikova expression is linked to, not integrated into, the viewer, because its bulk reuse terms could not be confirmed – a more complete atlas

would resolve that directly with the maintainers rather than work around it indefinitely. OSD-38 and OSD-522, both confirmed real and relevant, remain unprocessed. The spaceflight comparisons reported are mean ratios from two studies with their own genotype and lighting confounds, not a meta-analytic or statistically tested effect size, and should not be read as one.

4.2 What would make this stronger

Resolving TraVA's reuse terms directly, processing OSD-38 and OSD-522 to extend organ-matched spaceflight coverage beyond root, and replacing illustrative structural parameters with values measured from the cited primary sources (rather than only citing the source's general conclusions) would each move this resource further from "a rice-atlas pattern applied honestly" toward "a primary structural and functional atlas in its own right."

Data and code availability

All code, the real raw/processed data tables, and the viewer source are at <https://github.com/dr-richard-barker/arabidopsis-atlas>, deployed at <https://dr-richard-barker.github.io/arabidopsis-atlas/>, and archived at Zenodo (**TODO: DOI, pending manual deposit**). Primary spaceflight data are NASA OSDR studies OSD-120 [8] and OSD-314 [9]; developmental expression context is from the Klepikova/TraVA atlas [1]; structural citations are Shahan et al. [2] and Vijayan et al. [3]. Ecotype trait data are from Camargo et al. [4] (raw data in `data/raw/camargo2014_rosette_descriptors.csv`), Torii et al. [5], Bundy et al. [6], and Coneva and Chitwood [7]; reference photography is Taghavi Namin et al. [10]. rice-atlas, the project's own starting point, is at <https://github.com/dr-richard-barker/rice-atlas>; none of its code is reused here (see Methods).

Competing interests

The author declares no competing interests.

Acknowledgements

We thank the investigators who deposited OSD-120 and OSD-314, and NASA's Open Science Data Repository and GeneLab for curating and publishing them.

References

- [1] Anna V. Klepikova, Artem S. Kasianov, Evgeny S. Gerasimov, Maria D. Logacheva, and Aleksey A. Penin. A high resolution map of the *Arabidopsis thaliana* developmental transcriptome based on rna-seq profiling. *The Plant Journal*, 88(6):1058–1070, 2016. doi: 10.1111/tpj.13312.
- [2] Rachel Shahan, Che-Wei Hsu, Trevor M. Nolan, Benjamin J. Cole, Isaiah W. Taylor, Laura Greenstreet, Stephen Zhang, Anton Afanassiev, Anna Hendrika Cornelia Vlot, Geoffrey Schiebinger, Philip N. Benfey, and Uwe Ohler. A single-cell *Arabidopsis* root atlas reveals developmental trajectories in wild-type and cell identity mutants. *Developmental Cell*, 57(4):543–560.e9, 2022. doi: 10.1016/j.devcel.2022.01.008.
- [3] Athul Vijayan, Rachele Tofanelli, Sören Strauss, Lorenzo Cerrone, Adrian Wolny, Joanna Strohmeier, Anna Kreshuk, Fred A. Hamprecht, Richard S. Smith, and Kay Schneitz. A digital 3D reference atlas reveals cellular growth patterns shaping the *Arabidopsis* ovule. *eLife*, 10:e63262, 2021. doi: 10.7554/eLife.63262.

- [4] Anyela Camargo, Dimitra Papadopoulou, Zoi Spyropoulou, Konstantinos Vlachonasios, John H. Doonan, and Alan P. Gay. Objective definition of rosette shape variation using a combined computer vision and data mining approach. *PLoS ONE*, 9(5):e96889, 2014. doi: 10.1371/journal.pone.0096889.
- [5] Keiko U. Torii, Norio Mitsukawa, Tomoko Oosumi, Yosuke Matsuura, Ryusuke Yokoyama, Ronald F. Whittier, and Yoshibumi Komeda. The Arabidopsis ERECTA gene encodes a putative receptor protein kinase with extracellular leucine-rich repeats. *The Plant Cell*, 8(4):735–746, 1996. doi: 10.1105/tpc.8.4.735.
- [6] Mark G. R. Bundy, Olivia A. Thompson, Matthew T. Sieger, and Elena D. Shpak. Patterns of cell division, cell differentiation and cell elongation in epidermis and cortex of Arabidopsis pedicels in the wild type and in erecta. *PLoS ONE*, 7(9):e46262, 2012. doi: 10.1371/journal.pone.0046262.
- [7] Viktoriya Coneva and Daniel H. Chitwood. Genetic and developmental basis for increased leaf thickness in the Arabidopsis Cvi ecotype. *Frontiers in Plant Science*, 9:322, 2018. doi: 10.3389/fpls.2018.00322.
- [8] NASA Open Science Data Repository. OSD-120: Transcription profiling of Arabidopsis thaliana roots, light and gravity factors, spaceflight vs. ground control. <https://osdr.nasa.gov/osdr/data/osd/meta/120>, 2026. Accessed 2026-09-18; CC0.
- [9] NASA Open Science Data Repository. OSD-314: Arabidopsis thaliana seedling rna-seq, light and gravity gradient, SpaceX-4/SpaceX-11. <https://osdr.nasa.gov/osdr/data/osd/meta/314>, 2026. Accessed 2026-09-18; CC0.
- [10] Sarah Taghavi Namin, Mohammad Esmaeilzadeh, Mohammad Najafi, Tim B. Brown, and Justin O. Borevitz. Deep phenotyping: deep learning for temporal phenotype/genotype classification. *Plant Methods*, 14:66, 2018. doi: 10.1186/s13007-018-0333-4.