

9/28/2006

E & M


we use Gaussian units (cgs) (Old book edition)

4-5 problems/week from book (Jackson)

Recommended: Look at Feynman Lectures

Electrostatics

Coulomb's law



$$\vec{F} = k \frac{q_1 q_2}{|\vec{r}|^2} \frac{\vec{r}}{|\vec{r}|}$$

this is force
on q_1 by q_2

$$\vec{E}(\vec{r}) = \frac{\vec{F}}{q_1}$$

Field from charge at \vec{r}_1 :

$$\text{Gaussian: } k=1 \quad \text{MKS: } \frac{1}{4\pi\epsilon_0}$$

$$\vec{E}(\vec{r}) = \frac{q_1}{|\vec{r}-\vec{r}_1|^3} (\vec{r}-\vec{r}_1)$$

Superposition: many charges q_i, \vec{r}_i

$$\vec{E}(\vec{r}) = \sum_i \frac{q_i}{|\vec{r}-\vec{r}_i|^3} (\vec{r}-\vec{r}_i) \quad \therefore \text{Linear theory}$$

Continuous distribution of charges

$$dq = \rho(\vec{r}) d^3x$$



$$\vec{E}(\vec{r}) = \int d^3x' \rho(\vec{x}') \frac{(\vec{r}-\vec{x}')}{|\vec{r}-\vec{x}'|^3}$$

$$\rho(\vec{r}) = \sum_i q_i \delta(\vec{r}-\vec{r}_i)$$

$$\delta(x-a) = 0 \quad \text{if } x \neq a$$

$$\int_{-\infty}^{+\infty} \delta(x-a) dx = 1$$

for any $f(x)$


$$\int_{-\infty}^{+\infty} f(x) \delta(x-a) dx = f(a)$$

$$\text{Ex: } \int dx f(x) \delta(x) = -f(0)$$

Integrate
by parts

$$\vec{E}(\vec{r}) = \int d^3x' \sum_i q_i \delta(\vec{r}'-\vec{r}_i) \frac{(\vec{r}-\vec{r}')}{|\vec{r}-\vec{r}'|^3} = \sum_i q_i \frac{(\vec{r}-\vec{r}_i)}{|\vec{r}-\vec{r}_i|^3}$$

Gauss law



$$\vec{E} \cdot \hat{n} ds = ?$$

$$\vec{E} \cdot \hat{n} ds = \frac{q}{r^2} \cos \theta ds$$

$$\cos \theta ds = r^2 d\Omega$$


solid angle

Solid angle = projection of surface \perp to direction
 $\frac{r^2}{r^2}$

$$\Omega = \frac{A_\perp}{r^2}$$

$$d\Omega = \frac{ds \cos \theta}{r^2}$$


$$\therefore \int_S \vec{E} \cdot \hat{n} ds = q \int d\Omega = 4\pi q$$



$$\oint_S \vec{E} \cdot \hat{n} ds = \int_{S_1} d\Omega - \int_{S_2} d\Omega = 0$$

since \hat{n} is
in different direction

Charges outside Gaussian
surface don't contribute.



$$\begin{aligned} \oint_{S_1+S_2+S_3} \vec{E} \cdot \hat{n} ds &= 0 = \int_{S_3} d\Omega - \int_{S_2} d\Omega \\ \oint_{S_1+S_3} \vec{E} \cdot \hat{n} ds &= \int_{S_1} d\Omega + \int_{S_3} d\Omega = \int_{S_1} d\Omega + \int_{S_2} d\Omega \end{aligned} \Rightarrow \int_{S_3} d\Omega = \int_{S_2} d\Omega$$

$$\oint_S \vec{E} \cdot \hat{n} ds = 4\pi \int_V \rho(\vec{x}) d^3x$$

In general: $\oint_S \vec{A}(\vec{x}) \cdot \hat{n} ds = \int_V (\nabla \cdot \vec{A}) d^3x$ S encloses V

$$\int_V d^3x (\nabla \cdot \vec{E} - 4\pi \rho(\vec{x})) = 0 \quad \text{for all } V \Rightarrow \boxed{\nabla \cdot \vec{E} = 4\pi \rho(\vec{x})}$$

Electrostatic boundary

$$\begin{aligned} \vec{E}(\vec{x}) &= \int d^3x' \rho(\vec{x}') \frac{(\vec{x} - \vec{x}')}{|\vec{x} - \vec{x}'|^3} = - \int d^3x' \rho(\vec{x}') \nabla_x \left(\frac{1}{|\vec{x} - \vec{x}'|} \right) \\ &= \nabla_x (\text{something}) \Rightarrow \boxed{\nabla \times \vec{E} = 0} \quad \text{for electrostatics} \quad \nabla \times (\nabla f) = 0 \end{aligned}$$

Given \vec{E} (static) $\exists \phi(\vec{x})$ such that $\boxed{\vec{E} = -\nabla \phi}$

$$\boxed{\phi(\vec{x}) = \int d^3x' \frac{\rho(\vec{x}')}{|\vec{x} - \vec{x}'|}}$$

single charge: $\phi(\vec{x}) = \frac{q}{|\vec{x} - \vec{x}_0|}$

this choice makes $\phi \rightarrow 0$ as $r \rightarrow \infty$

$$\nabla^2 \phi = -4\pi \rho(\vec{x}) \quad \text{Poisson eq.} \quad (\text{solution in free space: no boundary conditions})$$

If $\rho = 0$, $\nabla^2 \phi = 0$ Laplace eq.

Potential Energy of charge in a field

$$W = - \int_A^B \vec{F} \cdot d\vec{l}$$

Work you do to move charge from A to B

$$W = -q \int_A^B \vec{E} \cdot d\vec{l} = q \int_A^B \nabla\phi \cdot d\vec{l} = q[\phi(B) - \phi(A)] \quad \therefore \text{potential energy is } q\phi$$

$$E_{\text{pot}} = q\phi$$

Boundary conditions:

② surface charge

2 different media

① height $\rightarrow 0$

$$\int \vec{E} \cdot \hat{n} ds = (\vec{E}_2 \cdot \hat{n}) S - (\vec{E}_1 \cdot \hat{n}) S = 4\pi\sigma S$$

$(\vec{E}_2 - \vec{E}_1) \cdot \hat{n} = 4\pi\sigma$

Discontinuity in normal component of \vec{E} if $\sigma \neq 0$

tangential component?

\Rightarrow HW | (New Edition) 1.1, 1.3, 1.5, 1.10, 1.11 | \Leftarrow

10/3/2006

② $\int \vec{E} \cdot d\vec{l} = \vec{E}_2 \cdot \hat{t} L - \vec{E}_1 \cdot \hat{t} L = 0$

$$\int \vec{E} \cdot d\vec{l} = \int (\nabla \times \vec{E}) \cdot \hat{n} ds = 0$$

$(\vec{E}_2 - \vec{E}_1) \cdot \hat{t} = 0$



$\nabla^2 \frac{1}{|\vec{x} - \vec{x}'|} = -4\pi \delta(\vec{x} - \vec{x}')$

$\frac{1}{|\vec{x} - \vec{x}'|}$ is $\phi(\vec{x})$ due to unit charge in \vec{x}'

$\nabla^2 \phi = -4\pi\rho$

$$\phi(\vec{x}) = \int d^3 \vec{x}' \frac{\rho(\vec{x}')}{|\vec{x} - \vec{x}'|}$$

$$\nabla_x^2 \phi(\vec{x}) = \int d^3 \vec{x}' \rho(\vec{x}') \nabla^2 \left(\frac{1}{|\vec{x} - \vec{x}'|} \right) = -4\pi \rho(\vec{x})$$



$\sigma d = D$ dipole per unit surface

$E_{in} = 4\pi\sigma$

$\Delta\phi = 4\pi\sigma d = 4\pi D$


Summary:

$$\nabla \cdot \vec{E} = 4\pi\rho \quad \nabla \times \vec{E} = 0 \quad \vec{E} = -\nabla\phi \quad \nabla^2 \phi = -4\pi\rho$$

free space solution (no B.C.s)

$$\phi(\vec{x}) = \int d^3 \vec{x}' \frac{\rho(\vec{x}')}{|\vec{x} - \vec{x}'|}$$

General situation



charge distribution and boundaries

Solve $\nabla^2\phi = -4\pi\rho$ in V with B.C. on S

Green's theorem

$\vec{A}(\vec{x})$ vector field

$$\int_V d^3x \nabla \cdot \vec{A} = \int_S (\vec{A} \cdot \hat{n}) ds$$

$\vec{A} = \phi \nabla \psi$ ϕ, ψ scalar functions

$$\begin{aligned} & \int_V d^3x (\nabla \phi \cdot \nabla \psi + \phi \nabla^2 \psi) = \int_S (\phi \nabla \psi) \cdot \hat{n} ds \\ & \quad \text{notation } \nabla \psi \cdot \hat{n} = \frac{\partial \psi}{\partial n} \\ & = \int_S \phi \frac{\partial \psi}{\partial n} ds \end{aligned}$$

$$\phi \leftrightarrow \psi \text{ then subtract} \rightarrow \int_V d^3x (\phi \nabla^2 \psi - \psi \nabla^2 \phi) = \int_S \left(\phi \frac{\partial \psi}{\partial n} - \psi \frac{\partial \phi}{\partial n} \right) ds$$

Boundary conditions which specify ϕ inside

- 1) ϕ given on S (Dirichlet b.c.)
- 2) $\frac{\partial \phi}{\partial n}$ given on S (Neumann b.c.)

Suppose you have ϕ_1, ϕ_2 that satisfy $\nabla^2\phi = -4\pi\rho$ and $\phi_1 = \phi_2$ on S

$$u = \phi_1 - \phi_2 \quad \nabla^2 u = 0 \quad u = 0 \text{ on } S$$

$$\begin{aligned} \text{Apply } \cancel{\int_V} \quad & \int_V d^3x |\nabla u|^2 = 0 \quad \therefore \nabla u = 0 \text{ inside } V \quad \therefore u = \text{constant} \\ \phi = u & \Rightarrow \phi_1 = \phi_2 + \text{constant} \end{aligned}$$

Green's function for $\nabla^2\phi = -4\pi\rho$

$$\nabla_x^2 G(\vec{x}, \vec{x}') = -4\pi \delta(\vec{x} - \vec{x}') \quad (\text{unit point charge at } \vec{x}')$$

$$\text{Particular solution } \frac{1}{|\vec{x} - \vec{x}'|}$$

$$\text{General solution } G(\vec{x}, \vec{x}') = \frac{1}{|\vec{x} - \vec{x}'|} + F(\vec{x}, \vec{x}') \quad \text{where } \nabla^2 F = 0$$

Green theorem with $\phi = \phi$ $G = G$

$$\int_V d^3x' (\phi \nabla^2 G - G \nabla^2 \phi) = \int_V d^3x' \left((-4\pi)\phi(x') \delta(\vec{x} - \vec{x}') + G(\vec{x}, \vec{x}') 4\pi \Delta(x') \right)$$

$$= \int_S \left(\phi \frac{\partial G}{\partial n'} - G \frac{\partial \phi}{\partial n'} \right) ds$$

$$\boxed{\phi(\vec{x}) = \int_V d^3x' G(\vec{x}, \vec{x}') \Delta(\vec{x}') - \frac{1}{4\pi} \int_S \left(\phi \frac{\partial G}{\partial n'} - G \frac{\partial \phi}{\partial n'} \right) ds}$$

If ϕ given on S : choose $G=0$ on S

$$\phi(\vec{x}) = \underbrace{\int_V d^3x' G(\vec{x}, \vec{x}') \Delta(\vec{x}')}_{\text{volume term}} - \frac{1}{4\pi} \underbrace{\int_S \phi \frac{\partial G}{\partial n'} ds}_{\text{surface term}} \rightarrow \phi|_S = f(\vec{x})$$


This is what we put

G solution to point charge + simple (uniform) b.c. on S
namely $G=0$ on S (Dirichlet)

Ex: Find ϕ for $q=1$ @ x' , b.c. $\phi=0$ on $x=0$... method of images gives

Green funct. for sphere (outside)

you answer \rightarrow Green's function



Charge q @ \vec{y}

Image charge q' @ \vec{y}'

$$\phi(\vec{x}) = \frac{q}{|\vec{x} - \vec{y}|} + \frac{q'}{|\vec{x} - \vec{y}'|} \quad \leftarrow \text{Green function with } (q=1)$$

$$\vec{x} = x \hat{m}_1, \quad \vec{y} = y \hat{m}_2, \quad \vec{y}' = y' \hat{m}_2$$

$$\phi(\vec{x}) = \frac{q}{x |\hat{m}_1 - \frac{y}{a} \hat{m}_2|} + \frac{q'}{y' |\hat{m}_2 - \frac{x}{y'} \hat{m}_1|}$$

$$\phi|_{x=a} = \frac{q}{a |\hat{m}_1 - \frac{y}{a} \hat{m}_2|} + \frac{q'}{y' |\hat{m}_2 - \frac{a}{y'} \hat{m}_1|} = 0 \quad \text{If } \frac{y}{a} = \frac{a}{y'} \text{ then } |m| \text{ are same}$$

$$\text{If } \frac{q}{a} = -\frac{q'}{y'} \text{ and } \frac{y}{a} = \frac{a}{y'}, \quad \phi|_{x=a} = 0$$

$$\therefore y' = \frac{a^2}{y} \quad q' = -\frac{y}{a} q = -\frac{a^2}{ya} q \quad \therefore q' = -\frac{a}{y} q$$


$$G = \frac{1}{|\vec{x} - \vec{y}|} - \frac{a}{y} \frac{1}{|\vec{x} - \frac{a^2}{y} \hat{m}_1|}$$

$$\nabla^2 G = -4\pi \delta(\vec{x} - \vec{y})$$

$$G = \frac{1}{x |\hat{m}_1 - \frac{y}{x} \hat{m}_2|} - \frac{a}{y} \frac{y}{a^2} \frac{1}{|\hat{m}_2 - \frac{x}{a^2} \hat{m}_1|}$$

10/5/2006

$$G(\vec{x}, \vec{x}') = \frac{1}{|\vec{x} - \vec{x}'|} - \frac{\alpha' x'}{|\vec{x} - \frac{\alpha^2}{x'^2} \vec{x}'|}$$



$$\vec{x} \rightarrow x, \theta, \phi$$


$$\vec{x}' \rightarrow x', \theta', \phi'$$

$$\zeta = f(\theta, \phi, \theta', \phi')$$

$$|\vec{x} - \vec{x}'|^2 = x^2 + x'^2 - 2x x' \cos \gamma$$

$$\frac{-1}{\left| \frac{x' x}{a} \vec{x} - \frac{a^2}{x'} \vec{x}' \right|}$$

$$G(\vec{x}, \vec{x}') = (x^2 + x'^2 - 2x x' \cos \gamma)^{-\frac{1}{2}} - \left(\left(\frac{x' x}{a} \right)^2 + a^2 - 2x x' \cos \gamma \right)^{-\frac{1}{2}}$$



$$\frac{\partial G}{\partial x'} = - \left. \frac{\partial G}{\partial x'} \right|_{x=a} = -\frac{1}{a} \frac{x^2 - a^2}{(x^2 + a^2 - 2ax \cos \gamma)^{\frac{3}{2}}}$$

Solution:

$$\phi(x, \theta) = - \int_S a^2 d\Omega' \phi(a, \theta', \phi') \left(-\frac{1}{a} \right) \frac{x^2 - a^2}{[x^2 - 2ax \cos \gamma]^{\frac{3}{2}}} + \int_V d^3 x' G$$

$$d\Omega' = d\phi' d(\cos \theta')$$

Problem: conducting, grounded sphere + point charge
what is surface charge on sphere?



$$\frac{z}{q}$$

$$\boxed{-\left. \frac{\partial \phi}{\partial n} \right|_S = 4\pi \sigma}$$

$$-\left. \frac{\partial G}{\partial x} \right|_{x=a} = -\frac{1}{a} \frac{x^2 - a^2}{(x^2 + a^2 - 2ax \cos \gamma)^{\frac{3}{2}}}$$

total charge on
sphere has to be image charge

$$\sigma(a) = -\frac{2}{4\pi a} \frac{z^2 - a^2}{(z^2 + a^2 - 2az \cos \theta)^{\frac{3}{2}}}$$

Insulated ^{conducting} sphere total charge Q + point charge q $\phi = ?$



$$\cdot q$$

$$\text{use } \frac{Q}{4\pi r^2}$$

and add extra charge $Q - q$ @ center of sphere

$$\frac{Q}{4\pi r^2} + \frac{Q-q}{4\pi r^2} = \frac{Q}{4\pi r^2}$$

Comp. Exam Problem

Spherically symmetric $\phi(r) = \frac{f(r)}{r}$

$$f(r) \rightarrow A \quad r \rightarrow 0$$

$$f(r) \rightarrow B \quad r \rightarrow \infty$$

 f is non singular

(no singularities)

a) total charge?

$$r \rightarrow \infty \quad \phi \sim \frac{B}{r} \Rightarrow Q_{\text{tot}} = B$$

b) identify point charges: f is non singular ϕ has no singularities except at $r=0$

$$r \rightarrow 0 \quad \phi \sim \frac{A}{r} \quad \therefore \text{point charge } q = A \text{ @ } r=0$$

c) find $\rho(x)$

$$\nabla^2 \phi = -4\pi \rho$$

$$\nabla \phi = \frac{1}{r} \nabla f + f \nabla \left(\frac{1}{r} \right) \quad \nabla^2 \phi = \nabla \left(\frac{1}{r} \right) \cdot \nabla f + \frac{1}{r} \nabla^2 f + \nabla f \cdot \nabla \left(\frac{1}{r} \right) + f \nabla^2 \left(\frac{1}{r} \right)$$

$$\nabla^2 \left(\frac{1}{r} \right) = -4\pi \delta(r) \quad \nabla \left(\frac{1}{r} \right) = \frac{\partial}{\partial r} \left(r^{-1} \right) \hat{r} = -\frac{\hat{r}}{r^2} \quad \nabla f(r) = f'(r) \hat{r}$$

$$\nabla^2 f(r) = \frac{1}{r^2} \frac{\partial}{\partial r} r^2 \frac{\partial f}{\partial r} = \frac{1}{r^2} \left(2r \frac{\partial f}{\partial r} + r^2 \frac{\partial^2 f}{\partial r^2} \right) = \frac{2}{r} f' + f''$$

$$2 \nabla \left(\frac{1}{r} \right) \cdot \nabla f = 2 \left(-\frac{\hat{r}}{r^2} \right) \cdot (f' \hat{r}) = -\frac{2}{r^2} f'$$

$$\nabla^2 \phi = -\frac{2}{r^2} f' + \frac{2}{r^2} f'' + \frac{1}{r} f''' - 4\pi f \delta(r) = \frac{1}{r} f''' - 4\pi f(r) \delta(r)$$

$$= \frac{1}{r} f''' - 4\pi A \delta(r) = -4\pi \rho$$

Electrostatic energy



$$W = \frac{q_1 q_2}{|\vec{x}_1 - \vec{x}_2|} = \phi(\vec{x}_2) q_2 = \phi(\vec{x}_1) q_1$$

charge q_i, \vec{x}_i

$$W = \frac{1}{2} \sum_{i \neq j}^N \frac{q_i q_j}{|\vec{x}_i - \vec{x}_j|}$$

continuous \rightarrow

$$W = \frac{1}{2} \int d^3x \int d^3x' \frac{\rho(\vec{x}) \rho(\vec{x}')}{|\vec{x} - \vec{x}'|} = \frac{1}{2} \int d^3x \rho(\vec{x}) \phi(\vec{x})$$

in terms of E :

$$W = \frac{1}{2} \int d^3x \left(-\frac{1}{4\pi} \right) \phi \nabla^2 \phi = \frac{1}{8\pi} \int d^3x \nabla \phi \cdot \nabla \phi = \int d^3x \frac{|E|^2}{8\pi}$$

energy density

$$u = \frac{E^2}{8\pi}$$

$$u = \frac{1}{8\pi} (|E|^2 + |\vec{B}|^2)$$

 $\cdot q$

Force on sphere?

image charge $q' @ y' \leftarrow$ force between charges

$$q' = -\frac{a}{y} q$$

$$y' = \frac{a^2}{y}$$


$$F = \frac{q q'}{|y - y'|^2} = \frac{q^2 a}{y^3} \frac{1}{\left(1 - \frac{a^2}{y^2}\right)^2}$$

$$\propto \frac{1}{y^3}$$

HW 2 : 2.1, 2.7, 2.23, 2.26)

10/10/2006

Read Feynman Lectures on Gauss theory - Electrostatic fields



Pressure on conductor?

$$E = 4\pi\sigma$$

Displace surface by Δx



$$\Delta W = \frac{1}{8\pi} E^2 A \Delta x \quad \text{change in energy}$$

small wrt radius of curvature

$$\Delta W = \frac{(4\pi)^2}{8\pi} \sigma^2 A \Delta x$$

$$F = \frac{\Delta W}{\Delta x}$$

$$P = \frac{F}{A} = \frac{\Delta W}{\Delta x} \frac{1}{A} = 2\pi\sigma^2$$

Orthogonal functions

$$a \rightarrow b \rightarrow x$$

$$\{u_n(x)\}_n$$

Can we represent any $f(x)$ in interval $[a, b]$
in terms of $\{u_n\}$?

Define a scalar product

set of orthonormal functions

$$\langle f, g \rangle = \int_a^b dx f(x) g^*(x)$$

$$\int_a^b dx u_n(x) u_m^*(x) = \delta_{mn} = \begin{cases} 0 & m \neq n \\ 1 & m = n \end{cases}$$

$\{u_n\}$ is complete if $\forall f(x)$ in $[a, b]$: $f(x) = \sum_n a_n u_n(x)$

$$a_n = \int_a^b dx f(x) u_n^*(x)$$

$$\therefore \int_a^b dx \left| f(x) - \sum_{n=1}^N a_n u_n(x) \right|^2 = 0$$

$$f(x) = \sum_n \left(\int_a^b dx f(x) u_n^*(x) \right) u_n(x) \Rightarrow \sum_n u_n(x) u_n^*(x) = \delta(x-x') \quad \text{completeness relation}$$

Example:

$$\left\{ \sqrt{\frac{2}{a}} \sin\left(\frac{2n\pi x}{a}\right), \sqrt{\frac{2}{a}} \cos\left(\frac{2n\pi x}{a}\right) \right\}_{n=1,2,\dots}$$

complete in $[-\frac{a}{2}, \frac{a}{2}]$

$$\int_{-\frac{a}{2}}^{\frac{a}{2}} dx \sin\left(\frac{2n\pi x}{a}\right) \sin\left(\frac{2m\pi x}{a}\right) = \delta_{nm}$$

$$f(x) = \sum_n A_n \sin\left(\frac{2n\pi x}{a}\right) + B_n \cos\left(\frac{2n\pi x}{a}\right) \quad \text{Fourier series}$$

$$A_n = \frac{2}{a} \int_{-\frac{a}{2}}^{\frac{a}{2}} f(x) \sin\left(\frac{2n\pi x}{a}\right) dx$$

6

$$a \rightarrow \infty \xrightarrow[a]{2\pi n} k \text{ (continuous variable)} \quad \left\{ \frac{1}{\sqrt{2\pi}} e^{ikx} \right\}$$

$$f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dk A(k) e^{ikx} \quad \text{Fourier integral} \quad A(k) = \int_{-\infty}^{\infty} dx f(x) e^{-ikx} \frac{1}{\sqrt{2\pi}}$$

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} e^{ikx} e^{-ikx} dx = \delta(k-k') \quad \text{orthonormal}$$

$$\int_{-\infty}^{\infty} dk e^{i k(x-x')} = 2\pi \delta(x-x') \quad \text{completeness relation}$$

$$\text{Integral representations of } \delta: \int dx e^{i(k-k')x} = 2\pi \delta(k-k')$$

Separation of Variables

$\nabla^2 \phi = 0$ in rectangular coord.

$$\frac{\partial^2 \phi}{\partial x^2} + \frac{\partial^2 \phi}{\partial y^2} + \frac{\partial^2 \phi}{\partial z^2} = 0 \quad \phi(x, y, z) = X(x) Y(y) Z(z)$$

$$\Rightarrow \frac{1}{X} \frac{d^2 X}{dx^2} + \frac{1}{Y} \frac{d^2 Y}{dy^2} + \frac{1}{Z} \frac{d^2 Z}{dz^2} = 0 \quad \text{each } \frac{1}{\text{variable}} \frac{d^2 \text{variable}}{d\text{variable}^2} = \text{constant}$$


$$\frac{1}{X} \frac{d^2 X}{dx^2} = -\alpha^2 \quad \frac{1}{Y} \frac{d^2 Y}{dy^2} = -\beta^2 \quad \frac{1}{Z} \frac{d^2 Z}{dz^2} = \gamma^2$$

$$-\alpha^2 - \beta^2 + \gamma^2 = 0$$

Suppose α, β are real

$$X(x) \sim e^{\pm i\alpha x} \quad Y(y) \sim e^{\pm i\beta y} \quad \gamma^2 = \alpha^2 + \beta^2 \quad Z(z) \sim e^{\pm i\gamma z}$$

$$\phi \sim \sum_{\alpha, \beta} A_{\alpha, \beta} e^{\pm i\alpha x} e^{\pm i\beta y} e^{\pm i\gamma z}$$



Solve $\nabla^2 \phi = 0$ inside box

b.c. $\phi = 0$ on all faces except $z=c$ where $\phi = V(x, y)$

$$X \sim \sin\left(\frac{\pi n x}{a}\right) \quad Y \sim \sin\left(\frac{\pi m y}{b}\right)$$

$$\gamma = \sqrt{\left(\frac{\pi n}{a}\right)^2 + \left(\frac{\pi m}{b}\right)^2} = \pi \sqrt{\left(\frac{n}{a}\right)^2 + \left(\frac{m}{b}\right)^2}$$

$$Z \sim e^{\gamma z} - e^{-\gamma z} \Rightarrow Z(0) = 0 \quad Z = \frac{1}{2}(e^{\gamma z} - e^{-\gamma z}) = \sinh(\gamma z)$$

$$\phi = \sum_{n, m} A_{n, m} \sin\left(\frac{\pi n x}{a}\right) \sin\left(\frac{\pi m y}{b}\right) \sinh(\gamma z) \sqrt{\frac{2}{a}} \sqrt{\frac{2}{b}} \quad \text{Err: } 2\pi n \rightarrow \pi n$$

$$n, m = 1, 2, 3, \dots$$

$$n, m = 0 \Rightarrow \phi = 0 \quad \text{true if } V(x, y) = 0$$

Last b.c.


$$\phi(x, y, c) = V(x, y) = \sum_{n,m=1}^{\infty} A_{nm} \sin\left(\frac{\pi n x}{a}\right) \sin\left(\frac{\pi m y}{b}\right) \sinh(\gamma_{nm} c) \sqrt{\frac{2}{a}} \sqrt{\frac{2}{b}}$$

$$A_{nm} \sinh(\gamma_{nm} c) = \iint_{y=0, x=0}^a dx dy V(x, y) \sin\left(\frac{\pi n x}{a}\right) \sin\left(\frac{\pi m y}{b}\right) \sqrt{\frac{2}{a}} \sqrt{\frac{2}{b}}$$

normalization

10/12/2006

Separation of Variables



Cylindrical coords.

interested in ϕ for $r \rightarrow 0$

ϕ finite for $r \rightarrow 0$

$$\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial \phi}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 \phi}{\partial \theta^2} = 0 \quad \text{b.c. } \phi(\theta=0) = \phi(\theta=\pi) = V \neq r$$

$$\phi = R(r) \Psi(\theta)$$

$$r \frac{1}{R} \frac{d}{dr} \left(r \frac{dR}{dr} \right) + \frac{1}{4} \frac{d^2 \Psi}{d\theta^2} = 0$$

$$\frac{r}{R} \frac{d}{dr} \left(r \frac{dR}{dr} \right) = \nu^2$$

$$\frac{1}{4} \frac{d^2 \Psi}{d\theta^2} = -\nu^2$$

$$\Psi(\theta) = A_\nu \sin(\nu\theta) + B_\nu \cos(\nu\theta)$$

try $R \sim r^\alpha$

$$\alpha^2 r^\alpha = \nu^2 r^\alpha \quad \longleftrightarrow \quad \alpha^2 = \nu^2 \quad \alpha = \pm \nu \quad \therefore R(r) = a_\nu r^\nu + b_\nu r^{-\nu}$$

If $\nu = 0$

$$\Psi'' = 0 \implies \Psi(\theta) = A_0 + B_0 \theta$$

$$r \frac{dR}{dr} = \text{const} \quad R(r) = a_0 + b_0 \ln r$$

ϕ finite as $r \rightarrow 0 \implies b_0 = 0$

$$\theta = 0 \quad \phi = A_0 a_0 + \sum_\nu (a_\nu r^\nu + b_\nu r^{-\nu}) B_\nu = V \quad B_\nu = 0 \quad A_0 a_0 = V$$

$$\theta = \beta \quad \phi = (A_0 + B_0 \beta) a_0 + \sum_\nu (a_\nu r^\nu + b_\nu r^{-\nu}) A_\nu \sin(\nu \beta) = V \quad B_0 = 0 \quad \sin(\nu \beta) = 0$$

$$\nu \beta = n\pi \quad n = 1, 2, 3, \dots \quad \nu = \frac{n\pi}{\beta} \quad b_\nu = 0 \quad (\phi \text{ finite as } r \rightarrow 0)$$

$$\phi(r, \theta) = V + \sum_n A_n r^{\frac{n\pi}{\beta}} \sin\left(\frac{n\pi}{\beta} \theta\right)$$

$$\text{for } r \rightarrow 0 \quad \phi \sim V + A_r r^{\frac{\pi}{\beta}} \sin\left(\frac{\pi}{\beta} \theta\right)$$


$$E_r = -\frac{\partial \phi}{\partial r} \quad E_\theta = -\frac{1}{r} \frac{\partial \phi}{\partial \theta} \implies E \sim r^{\frac{\pi}{\beta} - 1}$$

$$\beta = 2\pi \quad \Rightarrow E \sim r^{-\frac{1}{2}}$$

if $\beta > \pi \rightarrow E \text{ diverges}$

$\beta < \pi \rightarrow E \rightarrow 0$

$$\nabla^2 \phi = 0 \quad \text{in spherical coords}$$



$$\frac{1}{r^2} \frac{\partial^2}{\partial r^2} (r \phi) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta \frac{\partial \phi}{\partial \theta}) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 \phi}{\partial \phi^2} = 0$$

$$\phi = \frac{U(r)}{r} P(\theta) Q(\phi)$$

$$r^2 \frac{1}{U} \frac{d^2 U}{dr^2} + \frac{1}{\sin \theta} \frac{1}{P} \frac{d}{d\theta} (\sin \theta \frac{dP}{d\theta}) + \frac{1}{\sin^2 \theta} \frac{1}{Q} \frac{d^2 Q}{d\phi^2} = 0$$

$$\frac{1}{Q} \frac{d^2 Q}{d\phi^2} = -m^2 \Rightarrow Q(\phi) = e^{\pm im\phi} \quad m \text{ integer if } 0 \leq \phi \leq 2\pi$$

$$\frac{r^2}{U} \frac{d^2 U}{dr^2} + \frac{1}{\sin \theta} \frac{1}{P} \frac{d}{d\theta} (\sin \theta \frac{dP}{d\theta}) - \frac{m^2}{\sin^2 \theta} = 0$$

$$\frac{r^2}{U} \frac{d^2 U}{dr^2} = l(l+1) \quad \frac{d^2 U}{dr^2} - \frac{l(l+1)}{r^2} U = 0$$

$$\frac{d}{d\theta} (\sin \theta \frac{dP}{d\theta}) \frac{1}{\sin \theta} + \left(l(l+1) - \frac{m^2}{\sin^2 \theta} \right) P = 0$$

$$U \sim r^\alpha \Rightarrow \alpha(\alpha-1)r^{\alpha-2} - l(l+1)r^{\alpha-2} = 0 \quad \alpha(\alpha-1) = l(l+1)$$

$$\Rightarrow \alpha = l+1 \quad \alpha = -l$$

$$\therefore U = A r^{l+1} + B r^{-l} \quad \frac{U}{r} = A r^l + B r^{-(l+1)}$$

$$\text{change variable: } x = \cos \theta \quad \frac{d}{d\theta} = \frac{dx}{d\theta} \frac{d}{dx} = -\sin \theta \frac{d}{dx} \quad \frac{1}{\sin \theta} \frac{d}{d\theta} = -\frac{d}{dx}$$

$$\frac{d}{dx} \left((1-x^2) \frac{dP}{dx} \right) + \left(l(l+1) - \frac{m^2}{1-x^2} \right) P = 0 \quad \text{case } m=0 \quad \text{azimuthal symmetry}$$

$$P(x) = \sum_{n=1}^{\infty} a_n x^n \quad \frac{d}{dx} \left(n a_n x^{n-1} - n a_n x^{n+1} \right) + l(l+1) a_n x^n = 0$$

$$\underbrace{n(n-1)a_n x^{n-2}}_{n \rightarrow n+2} - n(n+1)a_n x^n + l(l+1)a_n x^n = 0$$

$$(n+2)(n+1)a_{n+2} - (n(n+1) - l(l+1))a_n = 0$$

Recursion relation

$$a_{n+2} = \frac{n(n+1) - l(l+1)}{(n+2)(n+1)} a_n$$

want to converge @ $x=1 \Rightarrow l$ must be integer \Rightarrow finite sum

$P_l(x) \rightarrow$ polynomials of order l

$$P_0(x) = 1 \quad P_1(x) = x \quad P_2(x) = \frac{1}{2}(3x^2 - 1) \quad \text{Legendre Polynomials}$$

Choose arbitrary const. so that $P_l(1) = 1$

$$x \in [-1, 1]$$

Other solution is for $\theta = 0, \pi$ $\sin \theta = 0$ not well behaved @ $\theta = 0, \pi$

$$m=0: \quad \phi(r, \theta) = \sum_{l=0}^{\infty} [A_l r^l + B_l r^{-(l+1)}] P_l(\cos \theta)$$

$\{P_l\}$ is a complete set in $[-1, 1]$ $\therefore f(x) = \sum_{l=0}^{\infty} a_l P_l(x) \quad x \in [-1, 1]$

$$\int_{-1}^1 P_l P_{l'} dx = 0 \quad \text{if } l \neq l' \quad a_l = \int_{-1}^1 f(x) P_l(x) dx$$

normalization Normalization is $(2l+1)/2$

Orthogonality

$$\frac{d}{dx} ((1-x^2) \frac{dP_l}{dx}) + l(l+1) P_l = 0$$

$$x P_l \Rightarrow \int_{-1}^1 dx P_l \frac{d}{dx} \left[\right] + \int_{-1}^1 dx l(l+1) P_l P_{l'} = 0$$

Int by parts

$$0 - \int_{-1}^1 dx \frac{dP_l}{dx} \left[\right] + \int_{-1}^1 0 = 0$$


Copy and least then subtract

$$(l(l+1) - l'(l'+1)) \int_{-1}^1 dx P_l P_{l'} = 0$$

if $l \neq l'$, then $\int = 0$

orthonormal, complete set $\left\{ \sqrt{\frac{2l+1}{2}} P_l(x) \right\}$ Find expression of δ -function
in terms of P_l 's

Example



$$\phi(r=a) = V(\theta)$$

$$\nabla^2 \phi = 0 \text{ solve for } r \leq a$$

ϕ finite for $r \rightarrow 0$ so $B_l = 0$

$$\phi(r, \theta) = \sum_{l=0}^{\infty} A_l r^l P_l(\cos \theta)$$

$$\sum_{l=0}^{\infty} A_l a^l P_l(\cos \theta) = V(\theta) \quad A_l a^l = \frac{2l+1}{2} \int_{-1}^1 d(\cos \theta) V(\theta) P_l(\cos \theta)$$


HW 3: 2.9, 3.1, 3.3, 3.6

10/17/2006

Spherical coord., azimuthal symmetry

$$\phi(r, \theta) = \sum_{l=0}^{\infty} (A_l r^l + B_l r^{-(l+1)}) P_l(\cos \theta)$$

$$\sum_{n=0}^{\infty} x^n = \frac{1}{1-x}$$

Suppose you know solution on z-axis ($\theta=0$) $\phi(z)$ If I can write it as: $\phi(z) \sim \sum_l a_l z^l + b_l z^{-(l+1)}$ Then solution is that times $P_l(\cos \theta)$
(in whole space)Example: $\frac{1}{|\vec{x}-\vec{x}'|}$ is sol. of $\nabla^2 \phi = 0$ for $\vec{x} \neq \vec{x}'$

$$|\vec{x}| > |\vec{x}'| : \frac{1}{|\vec{x}-\vec{x}'|} = \frac{1}{z-z'} = \frac{1}{z} \frac{1}{1-\frac{z'}{z}} = \frac{1}{z} \sum_{l=0}^{\infty} \left(\frac{z'}{z}\right)^l$$

$$a_l = 0 \quad b_l = z'^l$$

$$\sum_{l=0}^{\infty} \frac{z'^l}{z^{2l}} P_l(\cos \theta) = \frac{1}{|\vec{x}-\vec{x}'|}$$

 $m \neq 0$

$$\frac{d}{dx} \left((1-x^2) \frac{dP}{dx} \right) + \left(l(l+1) - \frac{m^2}{1-x^2} \right) P = 0 \quad x \in [-1, 1]$$

$$\text{Try } P \sim x^m \sum_{n=0}^{\infty} a_n x^n$$

Look @ $x \approx 1$: $x = 1-y$ $y > 0$ $y \ll 1$

$$\frac{d}{dx} \rightarrow \frac{d}{dy} \quad 1-x^2 = (1+x)(1-x) \approx 2y$$

$$2y \frac{d}{dy} \left(2y \frac{dP}{dy} \right) + \left(l(l+1) 2y - m^2 \right) P = 0 \quad 4y \frac{d}{dy} \left(y \frac{dP}{dy} \right) - m^2 P = 0$$

$$P \sim y^\alpha \quad 4\alpha^2 - m^2 = 0 \Rightarrow \alpha = \pm \frac{m}{2}$$

$$\text{Ansatz: } P(x) = (1-x)^{\frac{m}{2}} \sum_{n=0}^{\infty} a_n x^n \rightarrow \text{find recursion relation for } a_n$$

 $\ell \rightarrow \text{positive integer} \quad m \rightarrow 0, \pm 1, \pm 2, \dots, \pm \ell$ $P_{lm}(x)$ complete set for any fixed m in $[-1, 1]$; Orthogonal $P_{lm} Q_m = Y_{lm}(\theta, \phi)$ complete set on sphere surface, orthonormal

normalization

Spherical Harmonics

 $0 \leq \theta \leq \pi$ $0 \leq \phi \leq 2\pi$

$$\text{Orthonormal: } \int_0^{2\pi} \int_{-1}^1 Y_{lm}(\theta, \phi) Y_{l'm'}^*(\theta, \phi) d\phi d(\cos \theta) = \delta_{ll'} \delta_{mm'}$$

completeness

$$\sum_{l=0}^{\infty} \sum_{m=-l}^l Y_{lm}(\theta, \phi) Y_{l'm'}^*(\theta, \phi) = \delta(\theta - \theta') \delta(\phi - \phi')$$

Spherical coordinates $\nabla^2 \phi = 0$ $\phi = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} (A_{\ell m} r^{\ell} + B_{\ell m} r^{-\ell-1}) Y_{\ell m}(\theta, \phi)$

Any function $g(\theta, \phi) = \sum_{\ell, m} A_{\ell m} Y_{\ell m}(\theta, \phi)$ $A_{\ell m} = \int_0^{2\pi} \int_{-1}^1 g(\theta, \phi) Y_{\ell m}^*(\theta, \phi) d\phi d(\cos\theta)$

Connection to QM

$$-\frac{\hbar^2}{2m} \nabla^2 \psi + (V(r) - E) \psi = 0 \quad \text{stationary state}$$

H atom $V(r) = -\frac{e^2}{|r|}$

$$\hat{L}^2 Y_{\ell m} = \ell(\ell+1) Y_{\ell m} \quad \hat{L}_z Y_{\ell m} = m Y_{\ell m}$$


$$\hat{L}^2 = -\left(\frac{1}{\sin\theta} \frac{\partial}{\partial\theta} \left(\sin\theta \frac{\partial}{\partial\theta}\right) + \frac{1}{\sin^2\theta} \frac{\partial^2}{\partial\phi^2}\right) \quad \hat{L}_z = -i \frac{\partial}{\partial\phi}$$

$$\vec{L} = \vec{r} \times \vec{p} = -i \vec{r} \times \vec{\nabla}$$

Problem:



2 cylindrical shells, find ϕ inside $\nabla^2 \phi = 0$



$$0 \leq r \leq a$$

$$0 \leq \phi \leq 2\pi$$

$$r=a$$

$$\text{b.c. } 0 < \phi < \pi \quad \phi = V_1$$

$$\pi < \phi < 2\pi \quad \phi = V_2$$

$$\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial \phi}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 \phi}{\partial \phi^2} = 0 \quad \phi(r, \phi) = R(r) P(\phi)$$

$$\frac{r}{R} \frac{d}{dr} \left(r \frac{dR}{dr} \right) + \frac{1}{P} \frac{d^2 P}{d\phi^2} = 0 \quad \frac{1}{P} P'' = -k^2 \quad P \sim A_k \sin(k\phi) + B_k \cos(k\phi)$$

$k \rightarrow$ integer so $P(\phi) = P(\phi + 2\pi)$
single valued soln

$$r \frac{d}{dr} \left(r \frac{dR}{dr} \right) - k^2 R = 0 \quad R \sim r^\alpha \Rightarrow \alpha^2 - k^2 = 0 \quad \alpha = \pm k \quad R \sim r^k + r^{-k}$$

$$k=0, 1, 2, \dots \quad r \rightarrow 0 \quad \phi \text{ is finite so } \cancel{r^k} \Rightarrow R \sim r^k$$

$$\phi(r, \phi) = \sum_{k=0}^{\infty} r^k (A_k \sin(k\phi) + B_k \cos(k\phi))$$

b.c.

$$r=a \quad \sum_{k=0}^{\infty} a^k (A_k \sin(k\phi) + B_k \cos(k\phi)) = \begin{cases} V_1 & 0 < \phi < \pi \\ V_2 & \pi < \phi < 2\pi \end{cases}$$

$$a^k A_k = \frac{1}{\pi} \int_0^\pi V_1 \sin(k\phi) d\phi + \frac{1}{\pi} \int_\pi^{2\pi} V_2 \sin(k\phi) d\phi$$

normalization: $\frac{2}{[\text{interval}]}$ for sin, cos

$$= -\frac{V_1}{\pi k} (\cos(k\pi) - 1) + \frac{V_2}{\pi k} (\cos(k\pi) - 1) = \begin{cases} 0 & k \text{ even} \\ \frac{V_1 - V_2}{\pi k} & k \text{ odd} \end{cases}$$

except Fourier series first term
(extra $\frac{1}{2}$ factor)

$k=\text{odd}$

$$= \frac{2}{\pi k} (V_1 - V_2) \quad \cos(k\pi) = (-1)^k$$

$$k=0 \quad \frac{V_1 + V_2}{2}$$

Repeat for B_k using $\cos(k\phi)$... find $B_k = 0$

@ $r=0$ ϕ is average of surfaces


$$\frac{V_1 + V_2}{2}$$

$$k = 2n+1 \quad n=0, 1, 2, 3, \dots$$

$$\phi(r, \varphi) = \frac{V_1 + V_2}{2} + \sum_{n=0}^{\infty} \frac{2}{\pi} \frac{V_1 - V_2}{(2n+1) a^{2n+1}} \sin((2n+1)\varphi) r^{2n+1}$$

10/19/2006

Solutions of $\nabla^2 \phi = 0$ in Cylindrical coordinates



$$(r, \varphi, z) : \nabla^2 \phi = \frac{\partial^2 \phi}{\partial r^2} + \frac{1}{r} \frac{\partial \phi}{\partial r} + \frac{1}{r^2} \frac{\partial^2 \phi}{\partial \varphi^2} + \frac{\partial^2 \phi}{\partial z^2} = 0$$

$$\phi = R(r) Q(\varphi) Z(z)$$

$$\frac{1}{R} \frac{d^2 R}{dr^2} + \frac{1}{r} \frac{1}{R} \frac{dR}{dr} + \frac{1}{r^2} \frac{1}{Q} \frac{d^2 Q}{d\varphi^2} + \frac{1}{Z} \frac{d^2 Z}{dz^2} = 0$$

$$\frac{1}{Z} \frac{d^2 Z}{dz^2} = +k^2 \quad Z \sim e^{\pm kz}$$

$$\frac{1}{Q} \frac{d^2 Q}{d\varphi^2} = -\nu^2 \quad Q \sim A_\nu \sin(\nu\varphi) + B_\nu \cos(\nu\varphi)$$

ν - integer so that ϕ is single valued

$$\frac{d^2 R}{dr^2} + \frac{1}{r} \frac{dR}{dr} + \left(k^2 - \frac{\nu^2}{r^2} \right) R = 0$$

$$x = rk \quad \frac{d}{dx} = k \frac{d}{dr} \quad \frac{d^2 R}{dx^2} + \frac{1}{x} \frac{dR}{dx} + \left(1 - \frac{\nu^2}{x^2} \right) R = 0$$

try $R \sim x^\alpha \sum_n a_n x^n \rightarrow$ Solution $J_\nu(x), J_{-\nu}(x)$ Bessel functions

If ν is integer, $J_\nu, J_{-\nu}$ are linearly dependent so need second linear indep. solution:

$$\begin{cases} N_\nu(x) = \frac{1}{\sin(\nu\pi)} \left[J_\nu(x) \cos(\nu\pi) - J_{-\nu}(x) \right] \\ J_\nu(x) \end{cases}$$

Neumann function
or Bessel function of 2nd kind

basis vectors in space
of solutions to Bessel eq.

Limiting form: $x \ll 1$ ($x \geq 0$)

$$\frac{d^2 R}{dx^2} + \frac{1}{x} \frac{dR}{dx} - \frac{\nu^2}{x^2} R = 0 \quad R \sim x^\alpha$$

$$\alpha(\alpha-1) + \alpha - \nu^2 = 0 \quad \alpha = \pm\nu \Rightarrow R \sim x^{\pm\nu}$$

$$x \ll 1 : J_\nu \sim x^\nu \quad N_\nu \sim x^{-\nu} \quad \nu \neq 0 \quad N \rightarrow \infty \text{ as } x \rightarrow 0$$

$$N_\nu \sim \ln x \quad \nu = 0$$

$$x \gg 1, v \quad J_v \sim \frac{1}{\sqrt{x}} \cos\left(x - \frac{v\pi}{2} - \frac{\pi}{4}\right) \quad N_v \sim \frac{1}{\sqrt{x}} \sin\left(x - \frac{v\pi}{2} - \frac{\pi}{4}\right)$$

$J_v(x)$ has infinite # of zeros $J_v(x_{vn}) = 0 \quad n=1, 2, \dots$

$$x_{vn} \approx n\pi + (v - \frac{1}{2})\frac{\pi}{2}$$

$\rho \in [0, a]$: $\left\{ J_v\left(\frac{x_{vn}}{a}\rho\right), n=1, 2, 3, \dots\right\}$ is a complete set


$\left\{ \int_0^a J_v\left(\frac{x_{vn}}{a}\rho\right) d\rho \right\}$ are orthogonal

$$\int_0^a J_{vn} J_{vm} d\rho = 0 \quad \text{if } n \neq m \\ = \frac{a^2}{2} \left(J_{v+1}(x_{vn}) \right)^2 \quad \text{if } n = m$$

can write any $f(\rho)$ for $\rho \in [0, a]$

$$f(\rho) = \sum_n A_n J_{vn}\left(\frac{x_{vn}}{a}\rho\right) \quad A_n = \frac{2}{a^2 (J_{vn}(x_{vn}))^2} \int_0^a d\rho f(\rho) J_{vn}\left(\frac{x_{vn}}{a}\rho\right)$$

Example:



$$\nabla^2 \phi = 0 \quad \text{inside cylinder}$$

$$\text{b.c.s: } \phi = 0 \quad \text{on surface}$$

$$\phi = V(\rho, \varphi) \quad @ z=L \quad 0 < \rho < a$$

$$\varphi\text{-dep} \sim A_m \sin(m\varphi) + B_m \cos(m\varphi) \quad m \text{ integer}$$

$$z\text{-dep} \sim \sinh(kz)$$

$$\rho\text{-dep} \sim J_m(k\rho) + N_m(k\rho)$$

$$\phi = 0 \quad \text{for } \rho = a$$

$$\therefore J_m(ka) = 0$$

ϕ finite for $\rho \rightarrow 0$

$$ka = x_{mn} \Rightarrow k = \frac{x_{mn}}{a}$$

$$J_m\left(\frac{x_{mn}}{a}\rho\right)$$


$$\phi = \sum_{m,n} \left(A_{mn} \sin(m\varphi) + B_{mn} \cos(m\varphi) \right) J_m\left(\frac{x_{mn}}{a}\rho\right) \sinh\left(\frac{x_{mn}}{a}z\right)$$

$$\sinh\left(\frac{x_{mn}}{a}z\right) A_{mn} = \frac{2}{\pi a^2 J_{mn}^2(x_{mn})} \int_0^{2\pi} \int_0^a \rho \sin(m\varphi) J_m\left(\frac{x_{mn}}{a}\rho\right) V(\rho, \varphi) d\rho d\varphi$$

Time-dependent fields

10/24/2006

Faraday Induction



$$\oint \vec{E} \cdot d\vec{l} = -\frac{1}{c} \frac{d}{dt} \int_S \vec{B} \cdot \hat{n} ds = -\frac{1}{c} \frac{d}{dt} \Phi_B$$

mass: $m=1$
cgs: $k=1$

(E, B have same dimensions in Gaussian)


flux of B

$$I = \frac{E}{R}$$

current \searrow resistance

(uninterrupted circuit)

Lorentz force $\vec{F} = q(\vec{E} + \frac{1}{c} \vec{v} \times \vec{B})$

Can get Faraday Induction from
Lorentz force + Galilean transformation

Frame of charge: $\vec{v}=0 \quad \vec{E}' = \vec{F} = q \frac{\vec{v}}{c} \times \vec{B}$


Must be \vec{E}' appearing because $\vec{v} \times \vec{B} = 0 \Rightarrow \vec{E}' = \frac{\vec{v}}{c} \times \vec{B}$ so $q\vec{E}' = \vec{F}' = \vec{F} = q \frac{\vec{v}}{c} \times \vec{B}$

In general:

$$\vec{E}' = \vec{E} + \frac{\vec{v}}{c} \times \vec{B}$$

true for $\frac{v}{c} \ll 1$ relativistic: $\vec{E}' = \gamma(\vec{E} + \frac{\vec{v}}{c} \times \vec{B}) + O(\frac{v}{c})^2$

$$\gamma = \frac{1}{\sqrt{1-v^2}} \quad \beta = \frac{v}{c}$$

B static, circuit moving @ \vec{v} rest frame of circuit: $\vec{E} = \frac{\vec{v}}{c} \times \vec{B}$

$$\oint \vec{E} \cdot d\vec{l} = \oint \left(\frac{\vec{v}}{c} \times \vec{B} \right) \cdot d\vec{l} = \int_S \nabla \times \left(\frac{\vec{v}}{c} \times \vec{B} \right) \cdot \hat{n} ds$$

$$\nabla \times (\vec{a} \times \vec{b}) = \vec{a}(\nabla \cdot \vec{b}) - \vec{b}(\nabla \cdot \vec{a}) + (\vec{b} \cdot \nabla)\vec{a} - (\vec{a} \cdot \nabla)\vec{b}$$

$$\nabla \cdot \vec{B} = 0 \quad \vec{v} \text{ is constant} \Rightarrow \nabla \cdot \frac{\vec{v}}{c} = 0 \quad (\vec{B} \cdot \nabla) \frac{\vec{v}}{c} = 0 \quad \Rightarrow \nabla \times \left(\frac{\vec{v}}{c} \times \vec{B} \right) = -\left(\frac{\vec{v}}{c} \cdot \nabla \right) \vec{B}$$

$$= - \int_S \left(\frac{\vec{v}}{c} \cdot \nabla \right) \vec{B} \cdot \hat{n} ds$$

$$\frac{d}{dt} = \frac{\partial}{\partial t} + \vec{v} \cdot \nabla$$

 \vec{B} static $\frac{\partial \vec{B}}{\partial t} = 0$


$$= -\frac{1}{c} \int_S \left(\frac{d\vec{B}}{dt} \right) \cdot \hat{n} ds = -\frac{1}{c} \frac{d}{dt} \int_S \vec{B} \cdot \hat{n} ds = \oint \vec{E} \cdot d\vec{l}$$

$$\oint \vec{E} \cdot d\vec{l} = -\frac{1}{c} \frac{d}{dt} \int_S \vec{B} \cdot \hat{n} ds \quad S \text{ fixed}, \vec{B} = \vec{B}(t)$$

$$\int_S (\nabla \times \vec{E}) \cdot \hat{n} ds = -\frac{1}{c} \int_S \frac{\partial \vec{B}}{\partial t} \cdot \hat{n} ds \quad \Rightarrow \quad \int_S (\nabla \times \vec{E} + \frac{1}{c} \frac{\partial \vec{B}}{\partial t}) \cdot \hat{n} ds = 0$$

$$\boxed{\nabla \times \vec{E} = -\frac{1}{c} \frac{\partial \vec{B}}{\partial t}}$$

Energy Associated with Magnetic Field

turn on \vec{B} : induce

$$\mathcal{E} = -\frac{Id}{Cdt} \Phi(B)$$

$$\frac{dW}{dt} = \mathcal{E}I = \frac{I}{C} \frac{d}{dt} \Phi(B)$$

$$\delta W = \frac{I}{C} \delta \Phi(B) = \frac{I}{C} \delta \int_S (\nabla \times \vec{A}) \cdot \hat{n} dS \quad \vec{B} = \nabla \times \vec{A}$$

$$= \frac{I}{C} \delta \left[\int_T \vec{A} \cdot d\vec{l} \right] \quad W = \frac{I}{C} \oint_T \vec{A} \cdot d\vec{l}$$

Consider 2 loops

 \vec{J} = current density

$$I = \int_J d\sigma \quad \text{cross section}$$

$$W = \frac{1}{C} \int_{T_2} \vec{A} \cdot \vec{J} d\vec{l} d\sigma$$

In general:

$$W = \frac{1}{2C} \int \vec{A} \cdot \vec{J} d^3x \quad \boxed{\text{}}$$

$$\text{Analogous to } W = \frac{1}{2} \int \rho \phi d^3x$$

$$\nabla \times \vec{B} = \frac{4\pi}{C} \vec{J} + \left(\frac{1}{C} \frac{\partial \vec{E}}{\partial t} \right)^{0 \text{ statics}} \quad \frac{1}{2C} \frac{\epsilon_0}{4\pi} \int \vec{A} \cdot (\nabla \times \vec{B}) d^3x$$

$$\int \vec{A} \cdot (\nabla \times \vec{B}) d^3x = \int \vec{B} \cdot (\underbrace{\nabla \times \vec{A}}_{\vec{B}}) d^3x \quad \nabla \cdot (\vec{a} \times \vec{b}) = \vec{b} \cdot (\nabla \times \vec{a}) - \vec{a} \cdot (\nabla \times \vec{b})$$

$$\int \nabla \cdot \vec{A} \times \vec{B} d^3x = \int_S (\vec{A} \times \vec{B}) \cdot \hat{n} dS = 0 \quad \text{take surface to infinity}$$

localized currents & fields

$$W = \frac{1}{8\pi} \int |\vec{B}|^2 d^3x$$

$$\text{energy density } u = \frac{1}{8\pi} (|\vec{E}|^2 + |\vec{B}|^2)$$

Before Maxwell

$$\nabla \cdot \vec{E} = 4\pi \rho$$

$$\nabla \times \vec{B} = \frac{4\pi}{C} \vec{J}$$

$$\nabla \times \vec{E} = -\frac{1}{C} \frac{\partial \vec{B}}{\partial t}$$

$$\nabla \cdot \vec{B} = 0$$

Necessity of new term: $\nabla \cdot \nabla \times \vec{B} = 0 = \frac{4\pi}{C} \nabla \cdot \vec{J}$ $\nabla \cdot \vec{J} = 0$ false with time varying quantities

$$\frac{\partial \rho}{\partial t} + \nabla \cdot \vec{J} = 0$$

$$\frac{4\pi}{C} \left(-\frac{\partial \rho}{\partial t} \right) = -\frac{4\pi}{C} \frac{1}{4\pi} \frac{\partial}{\partial t} (\nabla \cdot \vec{E})$$

⇒

$$\nabla \times \vec{B} = \frac{4\pi}{C} \vec{J} + \frac{1}{C} \frac{\partial \vec{E}}{\partial t} \quad \boxed{\text{}}$$

Static solutions in free space

$$\nabla \times \vec{E} = 0 \Rightarrow \vec{E} = -\nabla \phi$$

$$\nabla^2 \phi = -4\pi \rho$$

$$\phi(\vec{x}) = \int \frac{\rho(\vec{x}')}{|\vec{x} - \vec{x}'|} d^3x'$$

$$\vec{B} = \nabla \times \vec{A}$$

$$\nabla \times (\nabla \times \vec{A}) = -\nabla^2 \vec{A} + \nabla (\nabla \cdot \vec{A})$$

static situation, Coulomb gauge so that $\nabla \cdot \vec{A} = 0$

$$\nabla^2 \vec{A} = -\frac{4\pi}{C} \vec{J}$$

$$\vec{A}(\vec{x}) = \frac{1}{C} \int \frac{\vec{J}(\vec{x}')}{|\vec{x} - \vec{x}'|} d^3x'$$

10/25/2006

Time dependant

$$\nabla \cdot \vec{E} = 4\pi J \quad \nabla \times \vec{B} = \frac{4\pi}{c} \vec{J} + \frac{\partial \vec{E}}{\partial t} \quad \nabla \times \vec{E} = -\frac{1}{c} \frac{\partial \vec{B}}{\partial t} \quad \nabla \cdot \vec{B} = 0$$

$$\vec{B} = \nabla \times \vec{A}$$

$$\nabla \times \vec{E} + \frac{1}{c} \nabla \times \frac{\partial \vec{A}}{\partial t} = 0$$

$$\nabla \times (\vec{E} + \frac{1}{c} \frac{\partial \vec{A}}{\partial t}) = 0$$

$$\vec{E} + \frac{1}{c} \frac{\partial \vec{A}}{\partial t} = -\nabla \phi$$

$$\vec{E} = -\nabla \phi - \frac{1}{c} \frac{\partial \vec{A}}{\partial t}$$

Gauge transformation

$$\vec{A} \rightarrow \vec{A}' = \vec{A} + \nabla \Lambda \quad \nabla \times \vec{A}' = \vec{B} \quad \text{since} \quad \nabla \times \nabla \Lambda = 0$$

$$\phi \rightarrow \phi' = \phi - \frac{1}{c} \frac{\partial \Lambda}{\partial t}$$

$$\vec{E}' = -\nabla \phi + \frac{1}{c} \frac{\partial \vec{A}}{\partial t} - \frac{1}{c} \frac{\partial \Lambda}{\partial t} - \frac{1}{c^2} \frac{\partial^2 \Lambda}{\partial t^2} = \vec{E}$$

$$\nabla \cdot \vec{E} = -\nabla^2 \phi - \frac{1}{c} \frac{\partial}{\partial t} (\nabla \cdot \vec{A}) = 4\pi J$$

$$\nabla \times \vec{B} = -\nabla^2 \vec{A} - \nabla(\nabla \cdot \vec{A}) + \nabla \frac{\partial \phi}{\partial t} + \frac{1}{c^2} \frac{\partial^2 \vec{A}}{\partial t^2} = \frac{4\pi}{c} \vec{J}$$

$$\boxed{\nabla \cdot \vec{A} + \frac{1}{c} \frac{\partial \phi}{\partial t} = 0} \quad \text{Lorentz gauge} \quad \partial_\mu A^\mu = 0 \quad A^\mu = (\phi, \vec{A})$$

$$-\nabla^2 \phi + \frac{1}{c^2} \frac{\partial^2 \phi}{\partial t^2} = 4\pi J$$

$$-\nabla^2 \vec{A} + \frac{1}{c^2} \frac{\partial^2 \vec{A}}{\partial t^2} = \frac{4\pi}{c} \vec{J}$$

2 non homogeneous
wave equations

with transformation

$$\nabla \cdot \vec{A} + \nabla^2 \Lambda + \frac{1}{c} \frac{\partial \phi}{\partial t} - \frac{1}{c^2} \frac{\partial^2 \Lambda}{\partial t^2} = 0 \quad \nabla^2 \Lambda - \frac{1}{c^2} \frac{\partial^2 \Lambda}{\partial t^2} = -\left(\nabla \cdot \vec{A} + \frac{1}{c} \frac{\partial \phi}{\partial t} \right)$$

Still have freedom to make additional gauge transformation such that $\nabla^2 \Lambda - \frac{1}{c^2} \frac{\partial^2 \Lambda}{\partial t^2} = 0$

$$\nabla^2 \Psi(\vec{x}, t) - \frac{1}{c^2} \frac{\partial^2 \Psi}{\partial t^2} = -4\pi f(\vec{x}, t) \quad \text{source function}$$

Fourier Transform

$$\Psi(\vec{x}, t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{\Psi}(\vec{x}, \omega) e^{-i\omega t} d\omega \quad (\nabla^2 + k^2) \Psi(\vec{x}, \omega) = -4\pi f(\vec{x}, \omega)$$

$$\text{Find Green's function: } (\nabla^2 + k^2) G(\vec{x}, \vec{x}; \omega) = -4\pi \delta(\vec{x} - \vec{x}') \quad \text{free space: } G(R, \omega) \quad R = |\vec{x} - \vec{x}'|$$

$$\text{in spherical coords: } \frac{1}{R} \frac{d^2}{dR^2} (RG) + k^2 RG = -4\pi \delta(R)$$

$$\text{for } R \neq 0 \quad (\vec{x} \neq \vec{x}') \quad \frac{d^2}{dR^2} (RG) + k^2 RG = 0 \implies RG = e^{\pm ikR} \quad G(R, \omega) = A_+ \frac{e^{ikR}}{R} + A_- \frac{e^{-ikR}}{R}$$

$$\text{for } R \rightarrow 0 \quad G(R, \omega) = \frac{1}{R} \quad (kR \rightarrow 0 \quad \nabla^2 G = -4\pi \delta(r))$$

spherical wave

Green's function for wave eqn:

$$\nabla^2 G(\vec{x}, \vec{x}; t, t') - \frac{1}{c^2} \frac{\partial^2 G}{\partial t^2} = -4\pi \delta(\vec{x} - \vec{x}') \delta(t - t')$$

$$G = \int \frac{dw}{2\pi} G(\vec{x}, \vec{x}; \omega, t') e^{-i\omega t} \quad \delta(t - t') = \int \frac{dw}{2\pi} e^{-i\omega(t - t')}$$

$$(\nabla^2 + k^2) G(\vec{x}, \vec{x}; \omega, t) = -4\pi \delta(\vec{x} - \vec{x}') e^{i\omega t}$$



Solution: $G = \frac{e^{i k |\vec{x} - \vec{x}'|}}{|\vec{x} - \vec{x}'|} e^{i \omega t'}$

$$G(\vec{x}, \vec{x}'; t, t') = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i \frac{\omega}{c} |\vec{x} - \vec{x}'|} e^{i \omega t'} e^{-i \omega t} dw = \frac{1}{2\pi} \frac{1}{R} \int e^{-i \omega ((t-t') - \frac{1}{c} R)} dw$$

$$G(\vec{x}, \vec{x}'; t, t') = \frac{1}{|\vec{x} - \vec{x}'|} \delta \left[t' - \left(t - \frac{|\vec{x} - \vec{x}'|}{c} \right) \right]$$

time of propagation of disturbance

$$z = t - \frac{|\vec{x} - \vec{x}'|}{c} \quad \text{retarded time} \quad G = \frac{1}{|\vec{x} - \vec{x}'|} \delta(t' - z)$$

fields at \vec{x}, t depend on sources at \vec{x}' and time $t - \frac{|\vec{x} - \vec{x}'|}{c} = z$

$$\nabla^2 \psi - \frac{1}{c^2} \frac{\partial^2 \psi}{\partial t^2} = -4\pi f$$

$$\Psi(\vec{x}, t) = \iiint \frac{\delta(t - (t - \frac{|\vec{x} - \vec{x}'|}{c}))}{|\vec{x} - \vec{x}'|} f(\vec{x}', t') d^3x' dt'$$

$$\Psi(\vec{x}, t) = \int f(\vec{x}', t - \frac{|\vec{x} - \vec{x}'|}{c}) \frac{1}{|\vec{x} - \vec{x}'|} d^3x' = \int \frac{[f(\vec{x}', t')]_{\text{ret}}}{|\vec{x} - \vec{x}'|} d^3x'$$

$$\Phi(\vec{x}, t) = \int \frac{\rho(\vec{x}', t - \frac{|\vec{x} - \vec{x}'|}{c})}{|\vec{x} - \vec{x}'|} d^3x'$$

$$\vec{A}(\vec{x}, t) = \int \frac{\vec{j}(\vec{x}', t - \frac{|\vec{x} - \vec{x}'|}{c})}{|\vec{x} - \vec{x}'|} d^3x'$$

free space solution
(no boundaries)

Conservation of Energy

fields, currents, charges

$$\text{Power transferred to charges} \quad P = \vec{F} \cdot \vec{v} = q \vec{v} \cdot (\vec{E} + \frac{1}{c} \vec{v} \times \vec{B}) = q \vec{v} \cdot \vec{E}$$

$$\vec{j} = n q \vec{v} \quad n: \# \text{ charges / unit volume}$$

$$P = \vec{j} \cdot \vec{E} \implies P = \int \vec{j} \cdot \vec{E} d^3x$$

energy per unit time
transferred to charges

 ① fields with energy density $u = \frac{1}{8\pi} (E^2 + B^2)$

 ② energy current (radiation) \vec{P}

③ kinetic energy of charges ($\frac{d}{dt} \rightarrow \vec{j}, \vec{E}$)

$$\frac{\partial}{\partial t} \int_V u d^3x + \int_S \vec{P} \cdot \hat{n} ds + \int_V \vec{j} \cdot \vec{E} d^3x = 0 \implies \frac{\partial u}{\partial t} + \nabla \cdot \vec{P} + \vec{j} \cdot \vec{E} = 0$$

$$\rightarrow \text{find: } \vec{P} = \frac{c}{4\pi} (\vec{E} \times \vec{B}) \quad \text{Poynting vector}$$

11/2/2006

$$\nabla^2 G = -4\pi \delta(\mathbf{r}-\mathbf{r}') \Rightarrow G \sim \frac{1}{r} \quad \text{3-D}$$

~In r 2-D ← test problem or use 3-D and include $\int_{-\infty}^{\infty} dz$

Conservation of Energy

$$\frac{\partial u}{\partial t} + \nabla \cdot \vec{P} + \vec{j} \cdot \vec{E} = 0 \quad u = \frac{1}{8\pi} (E^2 + B^2)$$

energy
time · area

$$\nabla \times \vec{B} = \frac{4\pi}{c} \vec{j} + \frac{1}{c} \frac{\partial \vec{E}}{\partial t}$$

$$\frac{4\pi}{c} \vec{j} \cdot \vec{E} = \vec{E} \cdot (\nabla \times \vec{B}) - \frac{1}{c} \vec{E} \cdot \frac{\partial \vec{E}}{\partial t}$$

✓

$$-\nabla \cdot (\vec{E} \times \vec{B}) + \vec{B} \cdot (\nabla \times \vec{E}) - \frac{1}{c} \frac{\partial \vec{B}}{\partial t}$$

$$-\frac{1}{c} \frac{1}{2} \frac{\partial (E^2)}{\partial t}$$

$$\nabla \cdot (\vec{E} \times \vec{B}) = \vec{B} \cdot (\nabla \times \vec{E}) - \vec{E} \cdot (\nabla \times \vec{B})$$

$$\frac{\partial}{\partial t} \left(\frac{1}{8\pi} (E^2 + B^2) \right) + \nabla \cdot \vec{P} - \nabla \cdot (\vec{E} \times \vec{B}) \frac{c}{4\pi} - \frac{1}{8\pi} \frac{\partial B^2}{\partial t} - \frac{1}{8\pi} \frac{\partial E^2}{\partial t} = 0$$

$$\nabla \cdot \vec{P} = \frac{c}{4\pi} \nabla \cdot (\vec{E} \times \vec{B}) \Rightarrow \boxed{\vec{P} = \frac{c}{4\pi} (\vec{E} \times \vec{B})} \quad \text{Poynting vector}$$

energy current

Conservation of Momentum

$$\vec{F} = \rho (\vec{E} + \frac{1}{c} \vec{v} \times \vec{B}) \quad \text{Kinetic } \vec{p} \text{ per unit volume}$$

$$\frac{d\vec{P}_{ki}}{dt} = \rho \vec{E} + \frac{1}{c} \vec{j} \times \vec{B} \quad \frac{\text{momentum}}{\text{time · volume}}$$

$$\frac{d\vec{P}_{ki}}{dt} + \frac{\partial}{\partial t} \vec{P}_f + \text{div}(\text{tensor}) = 0$$

/ carried by charges field \vec{p} carried by fields

$$\nabla \cdot \vec{E} = -4\pi \rho \quad \nabla \times \vec{B} = \frac{4\pi}{c} \vec{j} + \frac{1}{c} \frac{\partial \vec{E}}{\partial t}$$

$$\nabla \times \vec{E} = -\frac{1}{c} \frac{\partial \vec{B}}{\partial t}$$

$$\frac{d\vec{P}_{ki}}{dt} = -\frac{1}{4\pi c} \vec{E} (\nabla \cdot \vec{E}) + \frac{1}{4\pi} (\nabla \times \vec{B}) \times \vec{B} - \frac{1}{4\pi c} \frac{\partial \vec{E}}{\partial t} \times \vec{B}$$

↓
 $\vec{E} (\nabla \cdot \vec{E}) + \vec{B} (\nabla \cdot \vec{B})$

$$\frac{\partial}{\partial t} (\vec{E} \times \vec{B}) - \vec{E} \times \frac{\partial \vec{B}}{\partial t}$$

$$c \vec{E} \times (\nabla \times \vec{B})$$

$$\frac{d\vec{P}_{ki}}{dt} = -\frac{1}{4\pi c} \frac{\partial}{\partial t} (\vec{E} \times \vec{B}) + \frac{1}{4\pi} \left[\vec{E} (\nabla \cdot \vec{E}) + \vec{B} (\nabla \cdot \vec{B}) - \vec{E} \times (\nabla \times \vec{E}) - \vec{B} \times (\nabla \times \vec{B}) \right]$$

totally anti-symmet. tensor

to show: $[\vec{a}] = \text{div}(\text{tensor})$

$$\epsilon_{ijk} : \quad \epsilon_{123} = 1 \quad \epsilon_{ijk} = -\epsilon_{jik} \quad (\vec{a} \times \vec{b})_i = \epsilon_{ijk} a_j b_k \quad (\nabla \times \vec{a})_i = \epsilon_{ijk} \partial_j a_k$$

$$\epsilon_{ijk} \epsilon_{ilm} = \delta_{il} \delta_{km} - \delta_{jm} \delta_{kl}$$

$$\vec{a} \cdot \vec{b} = a_i b_i \delta_{ik} = a_i b_i$$

$$(\text{div}(\text{tensor}))_i = \partial_k T_{ik}$$

$$[\vec{a}]_i = E_i \partial_k E_k + B_i \partial_k B_k - \epsilon_{ijk} E_j \epsilon_{klm} \partial_l E_m - \epsilon_{ijk} B_j \epsilon_{klm} \partial_l B_m$$

$$= \partial_k (E_i E_k) - (\partial_k E_i) E_k + \partial_k (B_i B_k) - (\partial_k B_i) B_k + (\underbrace{\delta_{il} \delta_{jm} - \delta_{im} \delta_{jl}}_{}) E_j \partial_l E_m$$

$$E_j \partial_i E_j - E_j \partial_i E_i$$

$$4\pi T_{ik} = E_i E_k + B_i B_k - \frac{1}{2} (E^2 + B^2) \delta_{ik} \quad \text{Maxwell stress tensor}$$

HW 4 : 6.1, 6.11, 6.14a, 6.15, 6.18

11/7/2006

Cons. momentum

$$\vec{P}_{\text{mech}} = \frac{\text{momentum of charges}}{\text{volume}}$$

$$-\frac{d\vec{P}_{\text{mech}}}{dt} = \frac{\partial \vec{P}_{\text{fields}}}{\partial t} + \text{div}(\text{Maxwell Stress Tensor})$$

$$\vec{P}_{\text{mech}} = \vec{P}_{\text{ki}} \quad \vec{P}_{\text{fields}} = \vec{P}_f \quad \text{momentum carried by fields}$$

$$\frac{d\vec{P}_{\text{ki}}}{dt} = \rho \vec{E} + \frac{1}{c} \vec{J} \times \vec{B}$$

$$\vec{P}_f = \frac{1}{4\pi c} (\vec{E} \times \vec{B}) = \frac{1}{c^2} \vec{S}$$

Pointing
current

$$u = \frac{1}{8\pi} (E^2 + B^2)$$

$$\frac{u}{c} \sim \vec{P}_f$$


$$u c \sim \vec{S}$$

$$[\vec{J}]_i = \partial_k (E_i E_k + B_i B_k - \frac{1}{2} (E^2 + B^2) \delta_{ik}) = \partial_k T_{ik} 4\pi$$

$$T_{ik} = \frac{1}{4\pi} (E_i E_k + B_i B_k - \frac{1}{2} (E^2 + B^2) \delta_{ik}) \quad \boxed{\text{Maxwell Stress Tensor}}$$

$$\frac{\partial}{\partial t} \left[\frac{1}{4\pi c} (\vec{E} \times \vec{B}) \right]_i + \partial_k T_{ik} = - \left[\frac{d\vec{P}_{\text{ki}}}{dt} \right]_i$$

$$\int_V d^3x \left[-\frac{d\vec{P}}{dt} \right]_i = \int_V d^3x \left[\frac{1}{4\pi} (\vec{E} \times \vec{B}) \right]_i + \int_S T_{ik} n_k dS$$



$T_{ik} n_k$: i th component of momentum flowing in direction n_k

Transformation: rotation R

$$\text{scalar: } S \xrightarrow{R} S' = S \quad \text{vector: } A_i \xrightarrow{R} A'_i = R_{ik} A_k \quad x'_i = R_{ik} x_k$$

$$\text{tensor of rank 2: } T_{ik} \xrightarrow{R} T'_{ik} = R_{il} R_{km} T_{lm}$$

 \vec{A} is vector if it transforms like a vector

$$R_{ik} = \frac{\partial x'_i(x_1, x_2, x_3)}{\partial x_k}$$

$$\text{parity } P: \vec{x} \rightarrow -\vec{x} \quad \therefore \vec{E} \rightarrow -\vec{E} \quad \phi \rightarrow \phi \quad \rho \rightarrow \rho$$

$$\vec{A} \rightarrow -\vec{A} \quad \vec{B} \rightarrow \vec{B} \quad (\text{pseudovector}) \quad \vec{J} \rightarrow -\vec{J}$$

Maxwell's Eqn invariant under parity

$$T: t \rightarrow -t \quad \therefore \vec{E} \rightarrow \vec{E} \quad \vec{A} \rightarrow -\vec{A} \quad \vec{J} \rightarrow -\vec{J} \quad \vec{B} \rightarrow -\vec{B} \quad \rho \rightarrow \rho \quad \phi \rightarrow \phi$$


$$\nabla \times \vec{B} = \frac{4\pi}{c} \vec{J} + \frac{1}{c} \frac{\partial \vec{E}}{\partial t} \quad \underbrace{\quad}_{+ - - -} \quad \text{M.E. invariant under CPT}$$

$$C: \text{charge inversion} \quad q \rightarrow -q$$

$$\vec{E} \rightarrow -\vec{E} \quad \rho \rightarrow -\rho \quad \phi \rightarrow -\phi$$

$$\vec{B} \rightarrow -\vec{B} \quad \vec{J} \rightarrow -\vec{J}$$

Problem:

steady state current I , pot. diff V

a) Calculate \vec{S} @ surface of wire $E = \frac{V}{L}$


$$\oint_S \vec{B} \cdot d\vec{s} = \int_r^L B dl \quad \left\{ \begin{array}{l} \vec{B} = \frac{2I}{cr} \\ \vec{S} = \frac{c}{4\pi} (\vec{E} \times \vec{B}) \end{array} \right.$$

$$\vec{S} = \frac{c}{4\pi} (\vec{E} \times \vec{B}) \quad S = -\frac{V}{L} \frac{2I}{c b} \hat{r} \frac{c}{4\pi} \text{ into wire}$$

b) Power dissipated: $\frac{VI}{L}$ energy current
 $S 2\pi b = \frac{V}{L} \frac{2I}{cb} \frac{c}{4\pi} 2\pi b = \frac{VI}{L}$

since steady state Energy from V converted to heat (dissipated)
otherwise charges would accelerate

Multipole Expansion (Chpt 4)



$$\phi = ? \quad |\vec{x}| > |\vec{x}'|$$

first term $\sim \frac{1}{r}$ (total charge)

second term $\sim \frac{1}{r^2}$ (dipole)

Spherical

$$\phi(r, \theta, \varphi) = \sum_{l,m} (A_{lm} r^l + B_{lm} \bar{r}^{l+1}) Y_{lm}(\theta, \varphi)$$

$\lim_{r \rightarrow 0, r \rightarrow \infty}$

$$\boxed{\phi(r, \theta, \varphi) = \sum_{l,m} \frac{4\pi}{2l+1} Q_{lm} \frac{1}{r^{l+1}} Y_{lm}(\theta, \varphi)}$$

$$\sum = \sum_{l=0}^{\infty} \sum_{m=-l}^l$$

$l=0$ monopole $l=1$ dipole

Find q_{lm} in terms of ρ $\phi(\vec{x}) = \int \frac{\rho(\vec{x}')}{|\vec{x} - \vec{x}'|} d^3 \vec{x}'$

$$r = |\vec{x}'| \quad \frac{1}{|\vec{x} - \vec{x}'|} = \sum_{lm} \frac{4\pi}{2l+1} \frac{r^l}{r^{l+1}} Y_{lm}(\theta, \varphi) Y_{lm}^*(\theta', \varphi')$$

$$\phi(\vec{x}) = \sum_{lm} \frac{4\pi}{2l+1} \left[\int Y_{lm}^*(\theta, \varphi') r^l \rho(\vec{x}') d^3 \vec{x}' \right] \frac{Y_{lm}(\theta, \varphi)}{r^{l+1}}$$

$$\therefore q_{lm} = \int \rho(\vec{x}') r^l Y_{lm}^*(\theta, \varphi') d^3 \vec{x}'$$

$$q_{l-m} = (-1)^m q_{l-m}^*$$

$$Y_{l-m} = (-1)^m Y_{l-m}^*$$

$$Y_{00} = \frac{1}{\sqrt{4\pi}} \quad Y_{11} = -\sqrt{\frac{3}{8\pi}} \sin\theta e^{i\varphi} \quad Y_{10} = \sqrt{\frac{3}{4\pi}} \cos\theta$$

$$Q_{00} = \frac{1}{\sqrt{4\pi}} \int \rho(\vec{x}') d^3x' = \frac{Q}{\sqrt{4\pi}}$$

$$x = r \sin\theta \cos\varphi \quad y = r \sin\theta \sin\varphi \quad z = r \cos\theta$$

$$\sin\theta e^{i\varphi} = \sin\theta (\cos\varphi - i\sin\varphi) = \frac{1}{r} (x - iy)$$

$$\vec{p} = \int d^3x' \rho(\vec{x}') \vec{x}'$$

$$Y_{11} = - \int d^3x' \rho(\vec{x}') r^1 \frac{1}{r^1} (x^1 - iy^1) \sqrt{\frac{3}{8\pi}} = -\sqrt{\frac{3}{8\pi}} (p_x - ip_y) \quad Q_{11} = \sqrt{\frac{3}{8\pi}} (p_x + ip_y)$$

$$Y_{22} = \frac{1}{4} \sqrt{\frac{15}{2\pi}} \sin^2\theta e^{i2\varphi} = \frac{1}{4} \sqrt{\frac{15}{2\pi}} (x+iy)^2 \frac{1}{r^2}$$

Quadrupole moment

$$Q_{ij} = \int \rho(\vec{x}') (3x_i x_j - r^2 \delta_{ij}) d^3x'$$

Cartesian Coordinates

$$\Phi(\vec{x}) = \int \frac{\rho(\vec{x}')}{|\vec{x} - \vec{x}'|} d^3x' \quad (|\vec{x}| \gg |\vec{x}'|)$$

$$\frac{1}{|\vec{x} - \vec{x}'|} = \frac{1}{|\vec{x}|} \left[1 - \frac{1}{1 - 2 \frac{x^1 x'^1 + y^1 y'^1 + z^1 z'^1}{x^2 + y^2 + z^2} \dots} \right] \approx \frac{1}{|\vec{x}|} \left[1 + \frac{\vec{x} \cdot \vec{x}'}{|\vec{x}|^2} \dots \right]$$

$$\Phi(\vec{x}) \approx \frac{1}{|\vec{x}|} \left(\underbrace{\int \rho(\vec{x}') d^3x'}_Q + \underbrace{\frac{\vec{x}}{|\vec{x}|^3} \cdot \int \rho(\vec{x}') \vec{x}' d^3x'}_{\vec{p} \cdot \vec{x}} \right)$$

$$\boxed{\Phi(\vec{x}) = \frac{Q}{|\vec{x}|} + \frac{\vec{p} \cdot \vec{x}}{|\vec{x}|^3} + \frac{1}{2} \sum_{ij} Q_{ij} \frac{x_i x_j}{|\vec{x}|^5} + \dots}$$

11/9/2006

$$E_r = -\frac{\partial \Phi}{\partial r} = \sum_{lm} 4\pi \frac{(l+1)}{(2l+1)} Q_{lm} \frac{Y_{lm}(\theta, \varphi)}{r^{l+2}}$$

$$E_\theta = -\frac{1}{r} \frac{\partial \Phi}{\partial \theta} = \dots \quad \text{take care of dipole } \vec{p} \text{ along } \hat{z} \quad l=1 \quad m=0$$


$$Y_{10} = \sqrt{\frac{3}{4\pi}} \cos\theta$$

$$E_r = 4\pi \frac{2}{3} Q_{10} \sqrt{\frac{3}{4\pi}} \frac{\cos\theta}{r^3} \quad Q_{10} = \sqrt{\frac{3}{4\pi}} p_z \quad \therefore E_r = \frac{2p \cos\theta}{r^3}$$

$$E_\theta = \frac{p \sin\theta}{r^3} \quad E_\varphi = 0$$

$$\vec{E}(\vec{x}) = \frac{3\hat{x}(\vec{p} \cdot \hat{x}) - \vec{p}}{|\vec{x}|^3} \quad \text{field of dipole in cartesian}$$


Quadrupole

monopole: $Q=0$ dipole: $p_x = a(-q) + (-a)(-q) = 0 = p_y = p_z$

quadrupole:

$$Q_{11} = -q(3a^2 - a^2) + -2a^2q = -4a^2q$$

Electrostatic Energy

slowly varying \vec{E} (across region with ρ)No $\frac{1}{2}$ since this is not energy of charges
It is energy by field

$$W = \int \phi(\vec{x}) \rho(\vec{x}) d^3x$$


summation notation

$$\phi(\vec{x}) = \phi(\vec{0}) + \vec{x} \cdot \nabla \phi(\vec{0}) + \frac{1}{2} \frac{\partial^2 \phi}{\partial x_i \partial x_j}(\vec{0}) x_i x_j$$

$$\frac{\partial E_j}{\partial x_i} \delta_{ij} = \nabla \cdot \vec{E} = 0 \quad \begin{matrix} \text{external field} \\ \text{not produced} \\ \text{by } \rho \end{matrix}$$

$$W = q \phi(\vec{0}) - \vec{E}(\vec{0}) \cdot \vec{p} - \frac{1}{2} \frac{1}{3} \frac{\partial E_j}{\partial x_i}(\vec{0}) \int d^3x \rho(x) [3x_i x_j - x^2 \delta_{ij}]$$

$$W = q \phi(\vec{0}) - \vec{p} \cdot \vec{E}(\vec{0}) - \frac{1}{6} \frac{\partial E_j}{\partial x_i}(\vec{0}) Q_{ij}$$

energy of dipole
in external fieldenergy of quadrupole
in external field

dielectric

molecules in $\vec{E} \rightarrow$ develop \vec{p} material in $E \rightarrow$ develops $\frac{\text{dipole moment}}{\text{volume}} = \vec{P}$ (polarization)write ϕ in material : contribution from "extra charges" $\sim \int \frac{\rho(\vec{x}')}{|\vec{x} - \vec{x}'|} d^3x'$
(free charges)+ contribution from polarization $\sim \int \frac{\vec{P}(\vec{x}') \cdot (\vec{x} - \vec{x}')}{|\vec{x} - \vec{x}'|^3} d^3x'$

$$\phi(\vec{x}) = \int \left\{ \frac{\rho(\vec{x}')}{|\vec{x} - \vec{x}'|} + \frac{\vec{P}(\vec{x}') \cdot (\vec{x} - \vec{x}')}{|\vec{x} - \vec{x}'|^3} \right\} d^3x'$$

Integration by parts...

$$= \int \left\{ \frac{\rho(\vec{x}')}{|\vec{x} - \vec{x}'|} + \vec{P}(\vec{x}') \cdot \nabla' \frac{1}{|\vec{x} - \vec{x}'|} \right\} d^3x' = \int \frac{1}{|\vec{x} - \vec{x}'|} \underbrace{\left[\rho(\vec{x}') - \nabla' \cdot \vec{P}(\vec{x}') \right]}_{\text{effective } \rho_{\text{eff}}} d^3x'$$

effective ρ_{eff}

$$\nabla \cdot \vec{E} = 4\pi \rho \xrightarrow{\text{inside}} \nabla \cdot \vec{E} = 4\pi (\rho - \nabla \cdot \vec{P})$$


$$\nabla \cdot (\vec{E} + 4\pi \vec{P}) = 4\pi \rho_{\text{extra}} \quad \text{Electrostatics in the material} \quad \nabla \times \vec{E} = 0$$

usual situation $\vec{P} \propto \vec{E} \Rightarrow \boxed{\vec{P} = \chi \vec{E}}$
 χ susceptibility

$$\nabla \cdot [(1+4\pi\chi) \vec{E}] = 4\pi \rho_{\text{extra}} \quad \boxed{1+4\pi\chi = \epsilon \text{ dielectric const.}}$$

$$\epsilon \vec{E} = \vec{D} \Rightarrow \boxed{\nabla \cdot \vec{D} = 4\pi \rho_{\text{extra}}}$$


b.c.



$$(\vec{D}_2 - \vec{D}_1) \cdot \hat{n} = 0 \quad \text{if } \rho_{\text{extra}} = 0 \quad \text{Normal component of } \vec{D} \text{ continuous}$$

$$(\vec{D}_2 - \vec{D}_1) \cdot \hat{n} = 4\pi \sigma_{\text{extra}}$$

$$\boxed{(\vec{E}_2 - \vec{E}_1) \times \hat{n} = 0} \quad \text{Tangential component (from } \nabla \times \vec{E} = 0)$$

Dielectric Sphere in Uniform external field $(\epsilon = \epsilon)$ no free charges $\rho_{\text{extra}} = 0$

$$\text{b.c. } r \rightarrow \infty \quad \vec{E} = E_0 \hat{z} \quad \phi \rightarrow -E_0 r \cos \theta$$

$$r \rightarrow 0 \quad \phi \text{ finite}$$

$$\frac{1}{r} \frac{\partial \phi_{\text{out}}}{\partial r} \Big|_{r=a} = \frac{1}{r} \frac{\partial \phi_{\text{in}}}{\partial r} \Big|_{r=a} \quad \begin{matrix} \text{tangential component} \\ \text{normal component} \end{matrix}$$

$$\phi = \sum_{l=0}^{\infty} (A_l r^l + B_l \bar{r}^{l+1}) P_l(\cos \theta)$$

$$\phi_{\text{out}} = A_1 r \cos \theta + A_0 + \sum_{l=0}^{\infty} \frac{B_l}{r^{l+1}} P_l(\cos \theta) \quad A_1 = -E_0$$

$$\phi_{\text{in}} = \sum_{l=0}^{\infty} C_l r^l P_l(\cos \theta)$$

$$\frac{\partial \phi_{\text{out}}}{\partial r} \Big|_{r=a} = -E_0 \cos \theta - \sum_l (l+1) B_l a^{-l-2} P_l(\cos \theta) = \epsilon \sum_l l C_l a^{l-1} P_l(\cos \theta) = \epsilon \frac{\partial \phi_{\text{in}}}{\partial r} \Big|_{r=a}$$

$$l=1 \quad -E_0 - 2B_1 a^{-3} = \epsilon C_1 \quad B_l = C_l = 0 \quad \text{for } l \neq 1$$

$$\frac{1}{a} \frac{\partial \phi}{\partial r} \Big|_{r=a} = E_0 \sin \theta - B_1 a^{-3} \sin \theta = -C_1 \sin \theta$$

$$-2B_1 a^{-3} - \epsilon C_1 = E_0$$

$$-B_1 a^{-3} + C_1 = -E_0$$

$$-2(E_0 + C_1) - \epsilon C_1 = E_0 \quad -C_1(2 + \epsilon) = 3E_0$$

$$C_1 = \frac{-3E_0}{2 + \epsilon}$$

$$B_1 = a^3 (C_1 + E_0) = a^3 \left(\frac{-3E_0}{2 + \epsilon} + E_0 \right)$$

$$-3E_0 + 2E_0 + \epsilon E_0$$

$$-E_0 \quad B_1 = \frac{\epsilon - 1}{\epsilon + 2} a^3 E_0$$

$$\vec{P} = \frac{\vec{p}}{\frac{4}{3}\pi a^3} = \frac{3}{4\pi} E_0 \frac{\epsilon - 1}{\epsilon + 2} \hat{z}$$

$$\phi_{\text{in}} = \frac{-3E_0}{2 + \epsilon} r \cos \theta$$

$$\phi_{\text{out}} = -E_0 r \cos \theta + \frac{\epsilon - 1}{\epsilon + 2} \frac{a^3}{r^2} E_0 \cos \theta$$


$$\vec{p} = \frac{\epsilon - 1}{\epsilon + 2} a^3 E_0 \hat{z}$$

11/14/2006

$$\vec{E}_{in} = \frac{3}{\epsilon+2} E_0 \hat{z} \quad (\epsilon < E_0 \text{ for } \epsilon > 1) \quad \vec{P} = \frac{3}{4\pi} E_0 \frac{\epsilon-1}{\epsilon+2} \hat{z}$$

macroscopically $\vec{P} = \chi \vec{E} \quad \epsilon = 1 + 4\pi \chi$

microscopically $\vec{p} = \gamma \vec{E}$ dipole moment acquired by molecule in \vec{E}
 ↗ molecular polarizability
 Relation $\chi \leftrightarrow \gamma$?



Field in a hole in a dielectric:

$$\vec{E}^e = \vec{E}_{hole} + \vec{E}_{plug} \quad \text{Superposition}$$

$$\vec{E} = \vec{E}_{hole} + \vec{E}_{plug} \quad (\text{uniformly polarized sphere})$$

 \vec{E} inside uniformly polarized sphere

$$\vec{E}_{plug} = \vec{E}_{in} - \vec{E}_0 = E_0 \left(\frac{3}{\epsilon+2} - 1 \right) \hat{z} = \frac{1-\epsilon}{2+\epsilon} E_0 \hat{z} = -\frac{4\pi}{3} \vec{P}$$

$$\vec{E}_{hole} = \vec{E} + \frac{4\pi}{3} \vec{P} \quad \vec{P} = \gamma \vec{E}_{hole} \quad n \underset{\text{volume}}{\neq \text{molecules}} \quad \vec{P} = n \gamma \vec{E}_{hole}$$

$$\vec{P} = n \gamma \left(\vec{E} + \frac{4\pi}{3} \vec{P} \right) \quad \chi = n \gamma \left(1 + \frac{4\pi}{3} \chi \right)$$

$$\chi \left(1 - \frac{4\pi}{3} n \gamma \right) = n \gamma$$

$$\boxed{\chi = \frac{n \gamma}{1 - \frac{4\pi}{3} n \gamma}} \quad \text{Clausius-Mossotti}$$

Magnetostatics (Chapter 5)

$$\nabla \cdot \vec{B} = 0 \quad \nabla \times \vec{B} = \frac{4\pi}{c} \vec{j} \quad \nabla \cdot \vec{j} = 0 \quad \vec{B} = \nabla \times \vec{A}$$

$$\nabla \times (\nabla \times \vec{A}) = \frac{4\pi}{c} \vec{j} \quad \text{gauge: } \nabla \cdot \vec{A} = 0 \quad (\text{previously } \frac{1}{c} \frac{\partial \phi}{\partial t} + \nabla \cdot \vec{A} = 0)$$

$$\nabla^2 \vec{A} = -\frac{4\pi}{c} \vec{j}$$

Solution:
$$\boxed{\vec{A}(\vec{x}) = \frac{1}{c} \int \frac{\vec{j}(\vec{x}')}{|\vec{x} - \vec{x}'|} d^3x'} \quad \text{in Cartesian coords.}$$

$$B_i = (\nabla \times A)_i = \epsilon_{ikm} \partial_k A_m = \epsilon_{ikm} \frac{1}{c} \int J_m(\vec{x}') \frac{1}{|\vec{x} - \vec{x}'|} d^3x'$$

$$\partial_k \frac{1}{|\vec{x} - \vec{x}'|} = -\frac{(\vec{x} - \vec{x}')_k}{|\vec{x} - \vec{x}'|^3}$$

(acts on x (not x')) $\frac{\partial}{\partial x_k}$

$$\boxed{\vec{B} = \frac{1}{c} \int \frac{\vec{j}(\vec{x}') \times (\vec{x} - \vec{x}')}{|\vec{x} - \vec{x}'|^3} d^3x'} \quad \text{Biot-Savart Law}$$

$$d\vec{B} = \frac{I}{c} \frac{d\vec{l} \times (\vec{x})}{|\vec{x}|^3} \quad \text{integrate over wire}$$

Special case: no currents in region of interest

$$\nabla \cdot \vec{B} = 0 \quad \nabla \times \vec{B} = 0 \quad \text{and boundary conditions}$$

$$\vec{B} = -\nabla \phi_m \Rightarrow \nabla^2 \phi_m = 0$$

Force between loops of current $\vec{F} = \mu_0 (\vec{E} + \frac{\vec{v}}{c} \times \vec{B})$

$$\vec{F} = \frac{1}{c} \int d^3x \times (\vec{J}(x) \times \vec{B}(x))$$

$$\vec{B} = \frac{I_1}{r_1} \int \frac{d\vec{l}_1 \times (\vec{x}_{12})}{|\vec{x}_{12}|^3}$$

\vec{B} from a circular current loop



$$\vec{J} = J \delta(r-a) \delta(\cos\theta) \hat{\phi}$$

$$\delta(\theta - \pi)$$

$$dA$$

$$A_\phi = \frac{1}{c} \int r^2 dr \underbrace{d(\cos\theta) d\phi}_1 \frac{I}{a} \delta(r-a) \delta(\cos\theta) [r]^{-\frac{1}{2}}$$

$$I = \int J \delta(r-a) \delta(\cos\theta) r d(\cos\theta) dr$$

$$I = Ja \quad J = \frac{I}{a}$$

$$\vec{F}_{12} = \frac{I_1 I_2}{c^2} \int \int \frac{d\vec{l}_1 \times (d\vec{l}_2 \times (\vec{x}_{12}))}{r_1 r_2 |\vec{x}_{12}|^3}$$

Spherical:

$$|\vec{x} - \vec{x}'| = \left(r^2 + r'^2 - 2rr'(\cos\theta \cos\theta' + \sin\theta \sin\theta' \cos(\phi - \phi')) \right)^{\frac{1}{2}}$$

$$= \frac{Ia}{c} \int_0^{2\pi} d\phi' \left(a^2 + r^2 - 2ar(\sin\theta \cos(\phi - \phi')) \right)^{-\frac{1}{2}}$$

WRONG

since $\vec{A} = \int d^3x' \frac{\vec{J}(x')}{|\vec{x} - \vec{x}'|}$ is in cartesian

$$\nabla^2 \vec{A} = \frac{4\pi}{c} \vec{J} \quad \text{Ex:} \quad \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial A_r}{\partial r} \right) = \frac{4\pi}{c} J_r$$



$$\vec{J} = -J \sin\phi \hat{x} + J \cos\phi \hat{y}$$

Azimuthal symmetry so choose $\phi = 0$

(calculate \vec{A} on x-z plane)

$$A_z = 0 \quad A_x \sim \int_0^{2\pi} d\phi' \frac{\sin\phi'}{[\dots \cos\phi']^{\frac{1}{2}}} = 0$$

$$A_y = \frac{Ia}{c} \int_0^{2\pi} d\phi' \frac{\cos\phi'}{(a^2 + r^2 - 2ar \sin\theta \cos\phi')^{\frac{1}{2}}} \quad A_x = A_z = 0 \quad A_y \rightarrow A_\phi$$

$$\text{far from loop } r \gg a \quad A_\phi = \frac{Ia}{c} \int_0^{2\pi} d\phi' \frac{1}{r} \cos\phi' \left(1 + \frac{a}{r} \sin\theta \cos\phi' \right)$$

$$\int_0^{2\pi} d\phi' \cos^2\phi' = \pi$$

$$A_\phi = \frac{I\pi}{c} \frac{a^2}{r^2} \sin\theta$$

$$\vec{B} = \nabla \times \vec{A}$$

$$\sim (1 - 2 \frac{a}{r} \sin\theta)^{-\frac{1}{2}}$$

$$B_r = \frac{1}{r \sin\theta} \frac{\partial}{\partial \theta} (\sin\theta A_\phi) = \frac{I\pi}{c} \frac{a^2}{r^3} 2 \cos\theta$$

$$B_\theta = \frac{-1}{r} \frac{\partial}{\partial r} (r A_\phi) = \frac{I\pi}{c} \frac{a^2}{r^3} \sin\theta \quad B_\phi = 0$$

Field of dipole $\vec{m} = \pi a^2 \frac{I}{c} \hat{z}$

$$B_r = 2m \frac{\cos\theta}{r^3}$$

$$B_\theta = m \frac{\sin\theta}{r^3}$$

$$B_\phi = 0$$

$$m = \frac{1}{c} I \text{ (area)}$$

11/16/2006

Multipole Expansion (localized current distrib.)

 $|\vec{x}| \gg |\vec{x}'|$

$$\vec{A}(\vec{x}) = \frac{1}{c} \int d^3x' \frac{\vec{J}(\vec{x}')}{|\vec{x} - \vec{x}'|}$$

$$\frac{1}{|\vec{x} - \vec{x}'|} = \frac{1}{|\vec{x}'|} + \frac{\vec{x} \cdot \vec{x}'}{|\vec{x}'|^3} + \dots$$

Monopole term

$$\vec{A}(\vec{x}) = \frac{1}{c} \frac{1}{|\vec{x}'|} \int \vec{J}(\vec{x}') d^3x' = 0 \quad \nabla \cdot \vec{J} = 0 \leftarrow \text{static} \quad \frac{\partial \vec{J}}{\partial t} = 0$$

$$\int \vec{J}(\vec{x}') d^3x' = 0 \quad \int x_j \frac{\partial J_i}{\partial x_i} d^3x' = - \int J_i \frac{\partial x_j}{\partial x_i} d^3x' + \dots = - \int J_j d^3x' = 0$$

Integration by parts

$$\vec{A}(\vec{x}) = \frac{1}{c} \frac{1}{|\vec{x}'|^3} \int d^3x' (\vec{x} \cdot \vec{x}') \vec{J}(\vec{x}')$$

$$A_i = \frac{1}{c} \frac{\vec{x}}{|\vec{x}'|^3} \cdot \int d^3x' \vec{x}' J_i = -\frac{1}{2} \left[\vec{x} \times \left(\int (\vec{x}' \times \vec{J}) d^3x' \right) \right], \quad \frac{1}{c} \frac{1}{|\vec{x}'|^3}$$

$$\vec{m} = \frac{1}{2c} \int \vec{x}' \times \vec{J}(\vec{x}') d^3x' \quad \therefore \vec{A}(\vec{x}) = \frac{\vec{m} \times \vec{x}}{|\vec{x}'|^3} \quad \text{dipole term}$$

magnetic dipole moment

Define magnetization \vec{M} (mag dipole / unit volume)

$$\boxed{\vec{M}(\vec{x}) \equiv \frac{1}{2c} (\vec{x} \times \vec{J})}$$

$$\vec{m} = \int \vec{M} d^3x$$

$$\vec{B} = \nabla \times \vec{A} = \frac{3\vec{x}(\vec{x} \cdot \vec{m}) - \vec{m}}{|\vec{x}'|^3}$$

$$\int d^3x \left[x_i J_k \frac{\partial x_j}{\partial x_{k'}} + x_j J_k \frac{\partial x_i}{\partial x_{k'}} \right] = \int d^3x \left[-x_j \frac{\partial}{\partial x_{k'}} (x_i J_k) + x_j J_k \frac{\partial x_i}{\partial x_{k'}} \right] = 0$$

Integrate by parts (first term) $- (x_j J_k \frac{\partial x_i}{\partial x_{k'}} + x_j x_i \frac{\partial J_k}{\partial x_{k'}})$

$$= \int d^3x (x_i J_j + x_j J_i) \quad \text{stuff in } \frac{\partial}{\partial x_{k'}} \quad J_k \frac{\partial x_i}{\partial x_{k'}} = \frac{\partial}{\partial x_{k'}} (x_i J_k) - x_i \frac{\partial J_k}{\partial x_{k'}} = 0$$

$$\int (\vec{x} \cdot \vec{x}') \vec{J} d^3x' \rightarrow x_k \int d^3x' J_i x'_k = \frac{1}{2} x_k \left(\underbrace{(J_i x'_k - J_k x'_i)}_{J_m x'_n (\delta_{im} \delta_{kn} - \delta_{km} \delta_{in})} d^3x' \right)$$

$$= \epsilon_{ijk} \epsilon_{lmn} J_m x'_n$$

$$\therefore = \frac{1}{2} \epsilon_{ijk} \epsilon_{lmn} x_k \int J_m x'_n d^3x'$$

$$= \epsilon_{ijk} \epsilon_{lmn} J_m x'_n$$

$$= -\frac{1}{2} \epsilon_{ilm} x_k \int (\vec{J} \times \vec{x}')_l d^3x' = \frac{1}{2} (\vec{x} \times \left(\int (\vec{J} \times \vec{x}') d^3x' \right)) = \vec{x} \cdot \int \vec{x}' J_i d^3x'$$

Gaussian units:

$$[E] = [B] = [M] = [\rho]$$

$$\vec{M} = \frac{1}{2c} (\vec{x} \times \vec{J}) \quad [M] = \frac{t}{L} \leftarrow \frac{e}{t L^2} = \frac{e}{L^2} = [E] \quad \vec{E} = \frac{Qq}{r^2} \hat{r}$$

Exercise: for any plane current loop



$$m = \frac{I}{c} (\text{area of loop})$$

in direction \perp loop

$$\text{Energy of dipole } U = -\vec{m} \cdot \vec{B}$$

Moving (discrete) charges

$$\vec{\pi} = \sum_i q_i \vec{v}_i \delta(\vec{x} - \vec{x}_i)$$

$$\vec{m} = \frac{1}{2c} \left\langle d^3x \sum_i q_i (\vec{x} \times \vec{v}_i) \delta(\vec{x} - \vec{x}_i) \right\rangle = \frac{1}{2c} \sum_i q_i (\vec{x}_i \times \vec{v}_i)$$

$$q_i = q \quad m_i = m \quad (\text{mass}) \quad \therefore \quad \vec{m} = \frac{q}{2cm} \sum_i (\vec{x}_i \times \vec{p}_i) = \frac{q}{2mc} \vec{L}$$

Relation between orbital angular momentum and \vec{m} : $\vec{m} = \frac{e}{2mc} \vec{L}$

Material acquires \vec{M} in presence of \vec{B}

$$\text{Contributions to } \vec{A}: \sim \int d^3x' \frac{\vec{j}(x')}{|x-x'|} \sim \int d^3x' \frac{\vec{M} \times (\vec{x}-\vec{x}')}{|\vec{x}-\vec{x}'|^3}$$

excess (free)
currents

internal dipoles

$$\vec{A}(\vec{x}) = \frac{1}{c} \int d^3x' \left\{ \frac{\vec{J}(x')}{|x-x'|} + c \frac{\vec{M}(x') \times (\vec{x}-\vec{x}')}{|x-x'|^3} \right\}$$

$$= \frac{1}{c} \left\{ d^3 x' \left\{ \frac{\vec{J}(x')}{|x-x'|} + c \vec{M}(x') \times \nabla \left(\frac{1}{|x-x'|} \right) \right\} \right.$$

$$\epsilon_{ijk} M_j \vec{d}_k \frac{1}{|\vec{x} - \vec{x}'|} \xrightarrow[\text{Parts}]{\text{Integrate by}} -\epsilon_{ijk} (\vec{d}_k M_j) \frac{1}{|\vec{x} - \vec{x}'|} = \frac{\nabla' \times \vec{M}(\vec{x}')}{|\vec{x} - \vec{x}'|}$$

$$\vec{A}(\vec{x}) = \frac{1}{c} \int d^3x' \frac{\vec{j}(x') + c \nabla' \times \vec{M}(x')}{|x - x'|} \quad \text{solution to} \quad \nabla^2 \vec{A} = -\frac{4\pi}{c} (\vec{j} + c \nabla \times \vec{M})$$

$$\nabla \times \vec{B} = \frac{4\pi}{c} (\vec{j} + c \nabla \times \vec{M}) \quad \vec{j}: \text{free current}$$

$$\vec{H} = \vec{B} - 4\pi\vec{M} \implies \nabla \times \vec{H} = \frac{4\pi}{c} \vec{J}$$


need to calculate \vec{M} (relation between \vec{B} & \vec{H})

$$\vec{B} = \mu \vec{H} \quad \text{for 1) paramagnetic materials} \quad \mu > 1 \quad (\text{permanent dipoles align to field})$$

2) diamagnetic materials $\mu < 1$ (no permanent dipoles)

Small effect: $\mu = 1 \pm \sim 10^{-5}$

3) Ferromagnetic materials



Boundary conditions: B_n continuous normal, H_t continuous tangent for ($\vec{J}_f = 0$)

Solving Magnetostatic problems

$$1) \vec{B} = \mu \vec{H} \quad \vec{B} = \nabla \times \vec{A} \quad \nabla \times \vec{H} = \frac{4\pi}{c} \vec{J} \quad \nabla \times \vec{B} = \frac{4\pi}{c} \mu \vec{J}$$

$$\nabla^2 \vec{A} = -\frac{4\pi}{c} \mu \vec{J}$$

$$\text{Special case } \vec{J} = 0 \quad \nabla \times \vec{H} = 0 \quad \vec{H} = -\nabla \phi_m \quad \nabla^2 \phi_m = 0$$

2) Ferromagnets with fixed magnetization given $\vec{M}(x)$


$$\vec{J} = 0 \quad \nabla \cdot \vec{B} = 0 \quad \nabla \cdot \vec{H} = -4\pi \nabla \cdot \vec{M} \quad \nabla \times \vec{H} = 0 \quad \vec{H} = -\nabla \phi_m \quad A_m = -\nabla \cdot \vec{M}$$

$$\nabla \cdot \vec{H} = 4\pi A_m \quad \nabla^2 \phi_m = -4\pi A_m$$

$$\sigma_m = \vec{M} \cdot \hat{n} / s$$

Problem: uniformly magnetized sphere

11/21/2006



$$A_m = -\nabla \cdot \vec{M} = 0 \quad \text{in volume} \quad \sigma_m = \vec{M} \cdot \hat{r} = M_0 \cos \theta$$

$$\nabla^2 \phi_m = 0 \quad H = -\nabla \phi_m \quad \phi_m = \int_{\text{Sphere}} d\Omega' \frac{\sigma_m(\theta')}{|\vec{x} - \vec{x}'|}$$

$$\phi_m = a^2 \int d\Omega' \frac{\sigma_m(\theta')}{|\vec{x} - \vec{x}'|} \quad \frac{1}{|\vec{x} - \vec{x}'|} = 4\pi \sum_{l,m} \frac{1}{2l+1} \frac{r'_l}{r''^{l+1}} Y_{lm}^{(0),0} Y_{lm}^{*(0),0}$$

Azimuthal symmetry $m=0$ $Y_{l0} \sim \cos \theta \Rightarrow$ only have $l=1 m=0$

$$\phi_m(r, \theta, \psi) = a^2 \frac{4\pi}{3} \frac{r_c}{r_s^2} \underbrace{\sqrt{\frac{3}{4\pi}} \cos \theta}_{Y_{10}} \left\{ \begin{array}{l} M_0 \cos \theta \underbrace{\sqrt{\frac{3}{4\pi}} \cos \theta}_{Y_{10}^*} d\Omega' \\ \end{array} \right.$$


$$\phi_m = \frac{4\pi}{3} M_0 a^2 \frac{r_c}{r_s^2} \cos \theta \quad \text{inside } r_c = r \quad r_s = a \quad \phi_m = \frac{4\pi}{3} M_0 r \cos \theta \quad \vec{H}_m = -\frac{4\pi}{3} M_0 \hat{z}$$

$$\vec{B}_{in} = \vec{H}_{in} + 4\pi \vec{M}_{in} = -\frac{4\pi}{3} M_0 \hat{z} + 4\pi M_0 \hat{z} = \frac{8\pi}{3} M_0 \hat{z}$$


outside $r_c = a \quad r_s = r$

$$\phi_m = \frac{4\pi}{3} M_0 \frac{a^3}{r^2} \cos \theta \quad \text{dipole} \quad \vec{m} = \frac{4\pi}{3} M_0 a^3 \hat{z} \quad \vec{B}_{out} = \vec{H}_{out}$$

B lines




H lines



SPCM to originate from magnetization charges



Problem 6.8



$$\vec{\omega} = \omega \hat{z}$$

$$\vec{E} = E_0 \hat{x}$$

$$\mathcal{J} \times 2\pi r \delta \cos \varphi = \omega p \cos \varphi \quad (\text{macroscopic})$$

$$\nabla \cdot \vec{D} = 4\pi \rho_{\text{free}}$$

$$-\nabla \cdot \vec{P} = \rho_{\text{pol}}$$

$$\vec{D} = \vec{E} + 4\pi \vec{P}$$

$$\nabla \cdot \vec{E} = 4\pi (\rho_{\text{free}} + \rho_{\text{pol}})$$

$$\oint_{\partial V} \vec{P} \cdot d\vec{S} = -q = \int_S \vec{P} \cdot \hat{n} dS = -\vec{P} \cdot \hat{n} A \Rightarrow \vec{P} \cdot \hat{n} = \sigma_{\text{pol}}$$

$$\mathcal{J} = \sigma_m V \delta(r-a)$$

$$\vec{M} = \frac{1}{2c} \vec{x} \times \vec{J}$$

$$-\nabla \cdot \vec{M} = \rho_m$$

$$\vec{M} \cdot \hat{n} = \sigma_m$$

$$\Phi_m = a^2 \int \frac{\sigma_m}{|\vec{x} - \vec{x}'|} d\Omega'$$

$$\nabla \cdot \vec{E} = 4\pi \rho$$

$$\nabla \cdot \vec{B} = 0$$

$$\nabla \times \vec{E} = -\frac{1}{c} \frac{\partial \vec{B}}{\partial t}$$

$$\nabla \times \vec{B} = \frac{4\pi}{c} \vec{J} + \frac{1}{c} \frac{\partial \vec{E}}{\partial t}$$

$$\vec{B} = \nabla \times \vec{A} \quad \vec{E} = -\nabla \phi - \frac{1}{c} \frac{\partial \vec{A}}{\partial t}$$

$$\text{gauge} \quad \nabla \cdot \vec{A} + \frac{1}{c} \frac{\partial \phi}{\partial t} = 0$$

$$\rightarrow \text{gave us wave eq} \quad \nabla^2 \phi - \frac{1}{c^2} \frac{\partial^2 \phi}{\partial t^2} = -4\pi \rho$$

free space solution

$$\phi(\vec{x}, t) = \int \frac{J(\vec{x}', t - \frac{|\vec{x} - \vec{x}'|}{c})}{|\vec{x} - \vec{x}'|} d^3x'$$

$$\vec{S} = \frac{c}{4\pi} (\vec{E} \times \vec{B}) \quad \text{energy current}$$

$$u = \frac{1}{8\pi} (|\vec{E}|^2 + |\vec{B}|^2) \quad \text{energy density}$$

Plane Wave Solutions (in vacuum)

$$\rho = 0 \quad \vec{J} = 0 \quad \nabla \times (\nabla \times \vec{E}) = -\nabla^2 \vec{E} + \nabla (\nabla \cdot \vec{E}) = -\frac{1}{c} \frac{\partial}{\partial t} (\nabla \times \vec{B}) = -\frac{1}{c^2} \frac{\partial^2 \vec{E}}{\partial t^2}$$

$$\nabla^2 \vec{E} = \frac{1}{c^2} \frac{\partial^2 \vec{E}}{\partial t^2} \quad \text{same for } \vec{B}$$

$$\nabla^2 u(\vec{x}, t) - \frac{1}{c^2} \frac{\partial^2 u}{\partial t^2} = 0 \quad \text{Plane wave soln: } u(\vec{x}, t) = A e^{i(\vec{k} \cdot \vec{x} - \omega t)} \quad |\vec{k}| = \frac{\omega}{c}$$

$$\vec{k} \cdot \vec{x} = \text{const}$$

$$A \in \mathbb{C}$$

11/28/2006

Maxwell's Eq. in vacuum

$$\nabla \cdot \vec{E} = 0 \quad \nabla \cdot \vec{B} = 0 \quad \nabla \times \vec{E} = -\frac{1}{c} \frac{\partial \vec{B}}{\partial t} \quad \nabla \times \vec{B} = \frac{1}{c} \frac{\partial \vec{E}}{\partial t}$$


$$\nabla^2 \vec{E} - \frac{1}{c^2} \frac{\partial^2 \vec{E}}{\partial t^2} = 0 \quad \text{same for } \vec{B}$$

Plane wave solution: $\vec{E} = \vec{E}_0 e^{i(\vec{k} \cdot \vec{x} - \omega t)}$ $|\vec{k}| = \frac{\omega}{c}$
 $\vec{B} = \vec{B}_0 e^{i(\vec{k} \cdot \vec{x} - \omega t)}$ $\vec{E}_0, \vec{B}_0 \in \mathbb{C}^3$

physical fields $\operatorname{Re}(\vec{E})$

$$\nabla \times \vec{E} = -\frac{1}{c} \frac{\partial \vec{B}}{\partial t} \Rightarrow i\vec{k} \times \vec{E}_0 = -\frac{1}{c} (-i\omega \vec{B}_0) \quad \vec{k} \times \vec{E}_0 = \frac{\omega}{c} \vec{B}_0 \quad \vec{k} = \frac{\omega}{c} \hat{k}$$

$$\vec{B}_0 = \hat{k} \times \vec{E}_0 \quad \vec{B} \perp \vec{k}, \vec{E} \quad \text{also} \quad \vec{k} \times \vec{B}_0 = -\frac{\omega}{c} \vec{E}_0 \rightarrow \vec{E}_0 = -\hat{k} \times \vec{B}_0 \quad \vec{E} \perp \vec{k}, \vec{B}$$

plane wave, in vacuum $|\vec{E}_0| = |\vec{B}_0|$

$$\text{energy flow} \quad \vec{s} = \frac{c}{4\pi} (\vec{E} \times \vec{B}) \quad \text{energy current}$$

$$\text{time average} \quad \langle \vec{s} \rangle = \frac{1}{2} \frac{c}{4\pi} \operatorname{Re} \{ \vec{E} \times \vec{B}^* \}$$

$$\langle \vec{s} \rangle = \frac{c}{8\pi} |\vec{E}_0|^2 \hat{k} \quad u = \frac{1}{8\pi} (|\vec{E}|^2 + |\vec{B}|^2)$$

$$\langle s \rangle = c \langle u \rangle \quad \langle u \rangle = \frac{1}{2} \frac{1}{8\pi} 2 |\vec{E}_0|^2 = \frac{1}{8\pi} |\vec{E}_0|$$

energy transported at velocity c

$$\vec{E}_0 = (E_1, E_2, E_3) \quad E_i \in \mathbb{C} \quad |\vec{E}_0| = \vec{E}_0 \cdot \vec{E}_0^* = \sum_k ((E_k^r)^2 + (E_k^i)^2)$$

$$E_i = E_i^r + i E_i^i$$

$$\text{physical } |\vec{E}|^2 = |\operatorname{Re} \{ \vec{E} \}|^2 = \left[\vec{E} = \vec{E}_0 e^{i\omega t} \right] = |\operatorname{Re} \{ \vec{E}_0 \} \cos \omega t - \operatorname{Im} \{ \vec{E}_0 \} \sin \omega t|^2$$

$$\langle |\vec{E}|^2 \rangle = \frac{1}{2} |\operatorname{Re} \{ \vec{E}_0 \}|^2 + \frac{1}{2} |\operatorname{Im} \{ \vec{E}_0 \}|^2 = \frac{1}{2} |\vec{E}_0|^2 = \frac{1}{2} \vec{E}_0 \cdot \vec{E}_0^*$$

$$\text{Rule: } \vec{Q} = \vec{A} e^{i\omega t} \quad \langle \vec{Q} \rangle = \frac{1}{2} \vec{A} \cdot \vec{A}^*$$

Plane waves in medium (ϵ, μ)

$$\nabla \cdot \vec{D} = 0 \quad \nabla \times \vec{H} = \frac{1}{c} \frac{\partial \vec{D}}{\partial t} \quad \vec{D} = \epsilon \vec{E} \quad \vec{B} = \mu \vec{H} \quad \vec{P} = \chi \vec{E} \quad \text{more generally } P_i = \chi_{ik} E_k$$

$$\nabla^2 \vec{H} - \frac{\epsilon \mu}{c^2} \frac{\partial^2 \vec{H}}{\partial t^2} = 0 \quad c \rightarrow \frac{c}{n} \quad n = \sqrt{\epsilon \mu} \quad \text{index of refraction}$$

same for H, B, E, D

$$\text{in terms of } E, B: \quad \nabla \cdot \vec{E} = 0 \quad \nabla \cdot \vec{B} = 0 \quad \nabla \times \vec{E} = -\frac{1}{c} \frac{\partial \vec{B}}{\partial t} \quad \nabla \times \vec{B} = \frac{\epsilon_0}{c} \frac{\partial \vec{E}}{\partial t}$$

$$\vec{E} = \vec{E}_0 e^{i(\vec{k} \cdot \vec{x} - \omega t)} \quad |\vec{k}| = n \frac{\omega}{c} \quad \vec{E} = \vec{E}_0 e^{i(n \vec{k} \cdot \vec{x} - \omega t)} \quad \text{same for } \vec{B}$$

physical path \rightarrow optical path = physical path $\times n$

$$\lambda = \frac{2\pi}{|\vec{k}|} \quad \begin{array}{c} n=1 \\ ||| \end{array} \quad \begin{array}{c} n>1 \\ ||| \end{array} \quad \begin{array}{c} \text{shorter wavelength} \\ \text{same frequency} \end{array}$$

$$i n \vec{k} \times \vec{B}_0 = \frac{n^2}{c} (-i \omega \vec{E}_0) \quad \vec{E}_0 = -\frac{1}{n} \vec{k} \times \vec{B}_0 \quad |\vec{B}_0| = n |\vec{E}_0|$$

energy transport

$$\vec{S} = \frac{c}{4\pi} (\vec{E} \times \vec{H}) \quad u = \frac{1}{8\pi} (\vec{E} \cdot \vec{D} + \vec{B} \cdot \vec{H})$$

$$\langle \vec{S} \rangle = \frac{c}{8\pi} (\vec{E} \times \vec{H}^*) = \frac{c}{8\pi} \frac{n}{\mu} |\vec{E}_0|^2 \vec{k}$$

$$\langle u \rangle = \frac{1}{8\pi} \frac{1}{2} (\vec{E} \cdot \vec{B}^* + \vec{B} \cdot \vec{H}^*) = \frac{1}{8\pi} \frac{1}{2} \left(\epsilon |\vec{E}_0|^2 + \frac{n^2}{\mu} |\vec{E}_0|^2 \right) = \frac{1}{8\pi} \epsilon |\vec{E}_0|^2$$

$$\langle \vec{S} \rangle = \langle u \rangle \text{ (velocity)} \quad \Rightarrow \frac{cn}{\mu \epsilon} = \frac{c n}{n^2} = \frac{c}{n}$$

Polarization

$$\begin{array}{l} \vec{E}_1 = \vec{E}_1 E_1 e^{i(\vec{k} \cdot \vec{x} - \omega t)} \quad \vec{E}_2 = \vec{E}_2 E_2 e^{i(\vec{k} \cdot \vec{x} - \omega t)} \\ |\vec{E}_1| = |\vec{E}_2| = 1 \quad \vec{E} \in \mathbb{R}^3 \\ \text{unit vector} \end{array} \quad \vec{E} = (\vec{E}_1 E_1 + \vec{E}_2 E_2) e^{i(\vec{k} \cdot \vec{x} - \omega t)} \quad E_1, E_2 \in \mathbb{C}$$

Case a) E_1, E_2 have same phase

$$E_1 = R_1 e^{i\varphi} \quad R_1, \varphi \in \mathbb{R}$$

$$E_2 = R_2 e^{i\varphi}$$

$$\vec{E} = (R_1 \vec{E}_1 + R_2 \vec{E}_2) e^{i(\vec{k} \cdot \vec{x} - \omega t)}$$

$$\text{amplitude} = (R_1^2 + R_2^2)^{\frac{1}{2}}$$

Linear Polarization in direction $R_1 \vec{E}_1 + R_2 \vec{E}_2$

$$e^{i\varphi_2} = i$$

Case b) E_1, E_2 are 90° out of phase, $R_1 = R_2 = R$

$$E_1 = R e^{i\varphi} \quad E_2 = R e^{i(\varphi \pm \frac{\pi}{2})}$$

$$\vec{E} = (\vec{E}_1 \pm i \vec{E}_2) R e^{i(\vec{k} \cdot \vec{x} - \omega t + \varphi)}$$


Circular Polarization

$$\vec{x} = 0$$

$$\varphi = 0$$


$$R \Re \{\vec{E}\} = \vec{E}_1 R \cos(\omega t) \pm \vec{E}_2 R \sin(\omega t)$$

$$i(-i)$$



Bases: $\{\vec{E}_1, \vec{E}_2\}$ or $\{\vec{E}_1 \pm i \vec{E}_2\}$

Case c) Elliptical Polarization

Reflection and Refraction

$$n_i = \sqrt{\epsilon_i \mu_i}$$

$$\vec{E}_i = \vec{E}_o^i e^{i(n_i \vec{k}_i \cdot \vec{x} - \omega t)} \quad \vec{B}_i = n_i \hat{k}_i \times \vec{E}_o^i \quad \vec{E}_r = \vec{E}_o^r e^{i(n_i \vec{k}_i \cdot \vec{x} - \omega t)} \quad \vec{E}_t = \vec{E}_o^t e^{i(n_i \vec{k}_i \cdot \vec{x} - \omega t)}$$

B.C.s: D_n, B_n cont. E_t, H_t cont.

frequency is same $\omega_i = \omega_r = \omega_t = \omega$ to satisfy BC + t @ z=0

$n_i k_{ix} = n_r k_{rx} = n_t k_{tx}$ to satisfy BC + x, y @ z=0

$$n_i |\vec{k}_i| \sin \theta_i = n_r |\vec{k}_r| \sin \theta_r = n_t |\vec{k}_t| \sin \theta_t \quad |\vec{k}_i| = |\vec{k}_r| = |\vec{k}_t| = \frac{\omega}{c}$$

$$\Rightarrow \theta_i = \theta_r$$

$$n_i \sin \theta_i = n_t \sin \theta_t$$

Reflection

Snell's law

Friday Dec 8 ~3pm class


Final Dec 12 3-6pm

No Tues. class


11/30/2006

HW 6: 5.6, 5.13, 5.17, 7.2, 7.28

Polarization || to plane of incidence



Plane is
polarized



Polarization ⊥ to plane



$(\text{since } n_2 > n_1) \rightarrow \text{look at } E_x \text{ component}$

$$E_r = E_i + E_t \quad \text{no phase change}$$

phase difference if $n_2 > n_1 \rightarrow \pi$ phase change in reflected wave

$$E_t \text{ cont: } E_o^i \cos \theta_i - E_o^r \cos \theta_i = E_o^t \cos \theta_t$$

$$H_t \text{ cont: } \frac{n_1}{\mu_1} E_o^i + \frac{n_1}{\mu_1} E_o^r = \frac{n_2}{\mu_2} E_o^t$$

$$R = \frac{E_o^r}{E_o^i} \quad T = \frac{E_o^t}{E_o^i}$$

reflection and transmission coefficients

$$+ R \cos \theta_i + T \cos \theta_t = + \cos \theta_i$$

$$\mu_1 = \mu_2 = 1 \quad R n_1 - T n_2 = -n_1$$

$$R_H = \frac{n_2 \cos \theta_i - n_1 \cos \theta_t}{n_2 \cos \theta_i + n_1 \cos \theta_t}$$

$$T_H = \frac{2n_1 \cos \theta_i}{n_2 \cos \theta_i + n_1 \cos \theta_t}$$

Brewster Angle

$$R_{||} = 0 \quad (\text{not } R_{\perp})$$


\vec{E} polarized \perp to plane

$$n_2 \cos \theta_i = n_1 \cos \theta_t \Rightarrow \tan \theta_B = \frac{n_2}{n_1}$$

$$\theta_i + \theta_t = \frac{\pi}{2}$$

$$\sin \theta_t = \cos \theta_i$$

$$\sin \theta_i = \cos \theta_t \quad \text{Snell's law} \rightarrow n_2 \cos \theta_i = n_1 \cos \theta_t$$



light produced from
oscillations in materials

Total Internal Reflection

$$\theta = \theta_i \quad \theta' = \theta_t \quad n_1 > n_2$$

$$\sin \theta' = \frac{n_1}{n_2} \sin \theta$$

$$\text{critical angle } \theta_c \rightarrow \sin \theta_c = \frac{n_2}{n_1} < 1$$

such that $\theta' = \pi/2$ no transmission

$$\theta > \theta_c \quad \sin \theta' > 1 \quad \theta' \text{ is complex}$$


total internal reflection

$$\sin \theta' \in \mathbb{R} \quad \cos \theta' = \sqrt{1 - \sin^2 \theta'} = i \sqrt{\sin^2 \theta' - 1} \quad \cos \theta' \text{ is pure imaginary}$$

$$\sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i}$$

$$\vec{E}_t = \vec{E}_0^t \exp \left\{ i(n_2 k x \sin \theta + n_2 k z \cos \theta - wt) \right\}$$

$$\vec{E}_t = \vec{E}_0^t e^{i(n_2 k x \sin \theta - wt)} e^{-zk \sqrt{n_1^2 \sin^2 \theta - n_2^2}}$$




$$E_t \sim e^{-\frac{z}{2\delta}}$$


$$2\delta = \frac{1}{k \sqrt{n_1^2 \sin^2 \theta - n_2^2}}$$

$$k = \frac{2\pi}{\lambda}$$

$$\delta = \frac{\lambda}{4\pi \sqrt{n_1^2 \sin^2 \theta - n_2^2}}$$




1 way to focus wave in a plane



wave radiated by material to cancel incident wave

3 ways materials radiate: a, b, -1



$$b \propto a \quad \left. \begin{array}{l} \text{prop. constant} \\ \text{is the same} \end{array} \right\} \Rightarrow \frac{b}{a} = \frac{B}{A \cos(\theta + \theta')}$$

$$B \propto A \cos(\theta + \theta')$$

real component in dir. of B

$$-1 \propto a \quad -1 \propto A \cos(\theta - \theta') \quad \left. \begin{array}{l} \text{-1} \\ \text{a} \end{array} \right\} \frac{a}{A \cos(\theta - \theta')} = 1$$

$$\text{Energy conservation} \quad 1 = \epsilon a^2 + b^2$$

$$1 = \epsilon A^2 + B^2$$

$$b^2 = \frac{\sin^2(\theta - \theta')}{\sin^2(\theta + \theta')} = R_\perp^2$$

Dispersion

model for $\epsilon(\omega)$: harmonically bound charge

$$\ddot{x} + \gamma \dot{x} + \omega_0^2 x = -\frac{e}{m} E(x, t)$$

$$E(x, t) = E_0 e^{-i\omega t} \quad (\text{range of variation of } x \ll \lambda)$$

damping to keep resonance finite

$$x \sim e^{i\omega t}$$

$$x_0 (-\omega^2 - i\gamma\omega + \omega_0^2) = -\frac{e E_0}{m}$$

$$x_0 = \frac{e E_0}{m} \frac{1}{(\omega_0^2 - \omega^2 - i\gamma\omega)}$$

dipole moment

$$p = -ex_0 = \frac{e^2}{m} \frac{E_0}{(\omega_0^2 - \omega^2 - i\gamma\omega)}$$

$$\vec{P} = \chi \vec{E}$$

$$\chi(\omega) = N \frac{e^2}{m} \frac{1}{(\omega_0^2 - \omega^2 - i\gamma\omega)}$$

charges per unit volume

$$\text{more generally: } \chi(\omega) = \sum_j \frac{N e^2 / m f_j}{\omega_j^2 - \omega^2 - i\gamma\omega}$$

$$\sum_j f_j = Z$$

$\approx e^-$ in atom
strength


$$E = 1 + 4\pi \chi$$

$$\epsilon = 1 + 4\pi \chi = 1 + \sum_j \frac{4\pi N e^2 / m}{\omega_j^2 - \omega^2 - i\gamma\omega} f_j$$

$\gamma \ll \omega_j$ typically

$\therefore \epsilon$ is almost real except at

$\omega \ll \omega_j$ $\text{Re}(\epsilon) > 1$
 $\omega \gg \omega_j$ $\text{Re}(\epsilon) < 1$



12/7/2006

Final Exam comprehensive up to reflection/refraction
 no dispersion

Dispersion

$$\epsilon(\omega) = 1 + \sum_j \frac{4\pi N e^2 / m}{\omega_j^2 - i\gamma\omega - \omega^2} f_j$$

* charges/unit volume

$$\sum_j f_j = Z$$

e⁻ in atom oscillator strength

large $\text{Im}\{\epsilon\}$ \rightarrow absorption

$$n = \sqrt{\epsilon} \quad E \sim e^{i(nkx - \omega t)}$$

$$n = a + ib \quad n^2 = a^2 - b^2 + 2iab$$

$$\text{Re}\{\epsilon\} = a^2 - b^2 \quad \text{Im}\{\epsilon\} = 2ab$$

$$E \sim e^{i(akx - \omega t)} e^{-bkx} \quad \text{attenuation length } l = \frac{1}{bk} = \frac{2\pi 2\sqrt{\text{Re}\{\epsilon\}}}{2\pi \text{Im}\{\epsilon\}}$$

$$\text{because } \text{Re}\{\epsilon\} \approx a^2 \quad \text{Im}\{\epsilon\} \approx 2\sqrt{\text{Re}\{\epsilon\}}b$$

Relation to DC conductivity:

$$\omega \rightarrow 0 \quad a) \quad \omega_j \neq 0 \quad \forall j \quad \epsilon \rightarrow 1 + \sum_j \frac{4\pi N e^2 / m}{\omega_j^2} f_j \quad \text{real, } > 1$$

no free e⁻

$$b) \text{ free electrons } \omega_0 = 0 \quad \epsilon = 1 + \sum_{\omega_j \neq 0} \frac{4\pi N e^2 / m f_j}{\omega_j^2} - \frac{4\pi N e^2 / m f_0}{\omega(\omega + i\gamma)}$$

$$\epsilon = \epsilon_0 + i \frac{4\pi N e^2 / m f_0}{\omega(\gamma - i\omega)}$$

singular for $\omega \rightarrow 0$

$$1) \quad \nabla \times \vec{H} = \frac{4\pi}{c} \vec{J} + \frac{1}{c} \frac{\partial \vec{D}}{\partial t}$$

$$\vec{J} = \sigma \vec{E} \quad \vec{D} = \epsilon_0 \vec{E}$$

$$\text{or : 2) } \nabla \times \vec{H} = \frac{1}{c} \frac{\partial \vec{D}}{\partial t} \quad \vec{D} = \epsilon \vec{E} \quad \text{includes singular part}$$

$$\Rightarrow \frac{4\pi}{c} \sigma \vec{E} + \frac{1}{c} \epsilon_0 (-i\omega) \vec{E} = -\frac{i\omega}{c} (\epsilon_0 + i \frac{4\pi \sigma}{\omega}) \vec{E} = -\frac{i\omega}{c} \epsilon \vec{E}$$

$$\epsilon = \epsilon_0 + i \frac{4\pi}{\omega} \sigma$$

$$\therefore \sigma = \frac{N e^2 f_0}{m(\gamma - i\omega)}$$

singularity for $\text{Im}\{\epsilon\}$ for $\omega \rightarrow 0$
 has strength $4\pi\sigma$

No damping ($\gamma=0$) ex: Plasma

$$\epsilon = \epsilon_0 - \frac{4\pi Ne^2}{m\omega^2} = \epsilon_0 - \frac{\omega_p^2}{\omega^2}$$

Plasma frequency $\omega_{\text{pl}}^2 = \frac{4\pi Ne^2}{m}$

$\omega \ll \omega_{\text{pl}}$ $\epsilon < 0$ $n = \sqrt{\epsilon} \rightarrow$ imaginary no wave propagation: reflected

$\omega > \omega_{\text{pl}}$ wave can propagate ex: UV light through metals

Plasma + External Constant \vec{B}

special case: $\hat{k} \parallel \vec{B}$


$$m\ddot{x} = q(\vec{E} + \frac{\vec{k}}{c} \times \vec{B})$$

↑ wave ↑ external

Ansatz $\vec{E} = E_0 (\hat{\epsilon}_1 \pm i\hat{\epsilon}_2) e^{i(\vec{k} \cdot \vec{x} - \omega t)}$ circularly polarized wave

$$\simeq E_0 (\hat{\epsilon}_1 \pm i\hat{\epsilon}_2) e^{-i\omega t} (\lambda \gg x)$$

$$\vec{x} = x_0 (\hat{\epsilon}_1 \pm i\hat{\epsilon}_2) e^{-i\omega t}$$



$$m\ddot{x} - \frac{q}{c}(\dot{x} \times \vec{B}) = q E_0 (\hat{\epsilon}_1 \pm i\hat{\epsilon}_2) e^{-i\omega t}$$

$$(\hat{\epsilon}_1 \pm i\hat{\epsilon}_2) \times \vec{B} = \pm i (\hat{\epsilon}_1 \pm i\hat{\epsilon}_2) B$$

$$x_0 (-m\omega^2 \mp q \frac{\omega}{c} B) = q E_0$$

$$x_0 = \frac{q E_0}{-m\omega^2 \mp q \frac{\omega}{c} B}$$

$$\vec{x} = \frac{-q}{m\omega (\omega \pm \frac{qB}{mc})} \vec{E}$$

$$q = -e$$

$$\frac{eB}{mc} = \omega_L$$
 Larmor frequency

$$\vec{x} = \frac{e}{m\omega (\omega \mp \omega_L)} \vec{E}$$

$$\epsilon = 1 - 4\pi \frac{e^2 N}{m\omega (\omega \mp \omega_L)}$$

Birefringent


2 indices of refraction
(for positive and negative helicity)

$$\omega \rightarrow 0 \quad \epsilon_{\pm} \rightarrow 1 \pm \frac{\omega_p^2}{\omega \omega_L}$$

$$\epsilon_{\pm} = 1 - \frac{\omega_p^2}{\omega(\omega \mp \omega_L)}$$

$$\omega \rightarrow \omega_L \quad \epsilon_- = 1 + \frac{\omega_p^2}{2\omega \omega_L}$$

$$\epsilon_+ = 1 + \frac{\omega_p^2}{\omega(\omega - \omega_L)}$$



upper sign: ϵ_+ positive helicity

lower sign: ϵ_- negative helicity

$$\omega \rightarrow 0 \quad \epsilon_+ \approx \frac{\omega_p^2}{\omega \omega_L}$$

$$n = \sqrt{\epsilon} \approx \frac{\omega_p}{\sqrt{\omega \omega_L}}$$

$$v_{\text{phase}} = \frac{1}{n} \frac{\omega}{c} = \frac{\omega}{c} \frac{\sqrt{\omega \omega_L}}{\omega_p}$$

$$v_g = \frac{dw}{dk}$$

$$n \frac{\omega}{c} = k$$

$$k = \frac{\omega}{c} \frac{\omega_p}{\sqrt{\omega \omega_L}}$$

$$\frac{1}{v_g} = \frac{dk}{d\omega} = \frac{1}{2} \frac{\omega_p}{\sqrt{\omega \omega_L}} \frac{1}{c}$$

$$v_g \rightarrow 0 \text{ as } \omega \rightarrow 0$$

Page 39 and 40 are the original division in my notes for Phys 210A and 210B.
They contain no information and are not included here.

Page 39 and 40 are the original division in my notes for Phys 210A and 210B.
They contain no information and are not included here.

1/9/2007

Dispersion

HW Chpt 7: 12, 13, 22, 23

$$\epsilon = \epsilon(\omega)$$

Model: harmonically bound charges

$$\epsilon = 1 + \frac{\omega_p^2}{\omega_0^2 - \omega^2 - i\gamma\omega} \quad \omega_p^2 = \frac{4\pi Ne^2}{m}$$

a) no free charges $\omega_0 \neq 0$ resonances $\omega \rightarrow \omega_0$ absorption (large $\text{Im}(\epsilon)$)b) free charges $\omega_0 = 0$ no damping (plasma) $\epsilon \approx 1 - \frac{\omega_p^2}{\omega^2}$
for $\omega < \omega_p$ wave is excluded ($E_{\text{co}} \propto \epsilon \rightarrow \text{pure imaginary}$)with damping $\gamma \neq 0$ $\epsilon = 1 - \frac{\omega_p^2}{\omega^2 + i\gamma\omega}$ has $\frac{1}{\omega}$ singularityfor $\omega \rightarrow 0$, related to conductivity σ $\bar{\sigma} = \sigma \bar{\epsilon}$

$$\epsilon = 1 + i \frac{4\pi\sigma}{\omega} \quad \sigma = \sigma(\omega) \quad \sigma \rightarrow \text{constant as } \omega \rightarrow 0 \quad (\text{DC conductivity})$$

EM wave propagation in a plasma with \vec{B}

$$\epsilon_{\pm} = 1 - \frac{\omega_p^2}{\omega(\omega \mp \omega_L)} \quad \omega_L = \frac{eB}{mc} \quad \text{Larmor freq.}$$

ionosphere is birefringent $E_+ \neq E_-$ positive helicity
negative helicity

EM wave in conductive medium

$$\epsilon = \epsilon_0 + i \frac{4\pi\sigma}{\omega} \quad n = \sqrt{\epsilon} \quad E \sim e^{i(nkx - \omega t)} \quad k = \frac{\omega}{c}$$

→ attenuation length good conductor $\frac{\sigma}{\omega} \gg 1$ poor conductor $\frac{\sigma}{\omega} \ll 1$

$$\hookrightarrow \text{skin depth} \quad \delta = \frac{c}{\sqrt{2\pi\sigma\omega}}$$


$$\text{mag. field} \gg \text{electric field} \quad \frac{|\vec{H}_0|}{|E_0|} = |n| \sim \sqrt{\frac{\sigma}{\omega}} \gg 1 \quad (\text{for good conductors})$$

$$n = n(\omega) \quad E \sim e^{i(n\omega k_0 x - \omega t)} \quad k_0 = \frac{\omega}{c} \quad n(\omega) k_0 = k(\omega)$$

$$E \sim e^{i(kx - \omega(t))} \quad k = \frac{n\omega}{c} \quad \omega(t) \quad \text{dispersion relation}$$


$$\text{phase velocity} \quad v_p = \frac{\omega}{k}$$

$$\text{group velocity} \quad v_g = \frac{dk}{d\omega} \Big|_{k_0}$$



E or B

$$u(x, t) = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} A(k) e^{i(kx - \omega(k)t)} dk$$



$\boxed{\Delta x \Delta k \sim 1}$ fundamental relation
width of pulse

weights $|u|^2, |A|^2$

$$\langle x \rangle = \int x |u|^2 dx \quad \langle k \rangle = \int k |A|^2 dk$$

$$\Delta x = (\langle (x - \langle x \rangle)^2 \rangle)^{1/2}$$

similar for Δk

like in QM

 $\Delta x \Delta k = \frac{1}{2}$ for Gaussian packets

$$u \sim e^{-\frac{x^2}{2\sigma^2}} e^{ik_0 x} \dots$$

taking $\Delta x = \sigma$

$$A(k) = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-ikx} \underbrace{e^{ik_0 x}}_{e^{i(k-k_0)x}} e^{-\frac{x^2}{2\sigma^2}} dx \sim e^{-2\sigma^2(k-k_0)^2}$$

$$\mathcal{F}\left\{e^{-\frac{x^2}{2\sigma^2}}\right\} \sim e^{-2\sigma^2 k^2} \quad (\text{Fourier Transform})$$

$$2\sigma^2 = \frac{4\sigma^2}{2} \quad \therefore \Delta k = \frac{1}{2\sigma}$$

$$\Delta x \Delta k = \sigma \frac{1}{2\sigma} = \frac{1}{2}$$

Kramers-Kronig Relations

$$C(\omega) = a(\omega) b(\omega)$$

$$C(t) = \int_{-\infty}^{\infty} \frac{1}{2\pi} C(\omega) e^{-i\omega t} d\omega \quad A(t), B(t)$$

convolution

$$C(t) = \int_{-\infty}^{\infty} A(\tau) B(t-\tau) d\tau \quad \leftarrow \dots$$

$$c(\omega) = \int C(t) e^{i\omega t} dt$$

$$a(\omega) b(\omega) = \int A(t) e^{i\omega t} dt \int B(t') e^{i\omega t'} dt'$$

$$C(t) = \int dt' \int dt' A(t) B(t') \underbrace{\int \frac{dw}{2\pi} e^{i\omega(t+t'-t')}}_{\delta(t+t'-t')} = \int_{-\infty}^{\infty} dt A(t) B(t-t')$$

$$\delta(t+t'-t'')$$

$$\vec{D}(x, \omega) = \vec{E}(\omega) \vec{E}(x, \omega)$$

$$\epsilon = 1 + 4\pi \chi(\omega)$$

$$\vec{D}(x, \omega) = \vec{E}(x, \omega) + 4\pi \chi(\omega) \vec{E}(x, \omega)$$

$$\vec{D}(x, \omega) = \int \vec{D}(x, t) e^{i\omega t} dt$$

$$\vec{D}(x, t) = \vec{E}(x, t) + \int_{-\infty}^{\infty} G(\tau) \vec{E}(x, t-\tau) d\tau$$

$$G = \mathcal{F}^{-1}\{4\pi \chi(\omega)\}$$

causality $G(\tau) = 0$ for $\tau < 0$ so that $D(\text{now})$ doesn't depend on $E(\text{future})$

$$G(\tau) = \int \frac{1}{2\pi} (\epsilon(\omega) - 1) e^{-i\omega\tau} d\omega$$

$$\epsilon(\omega) = 1 + \int G(\tau) e^{i\omega\tau} d\tau$$

G is finite $G \rightarrow 0$ for $\tau \rightarrow \infty$

$\rightarrow E$ analytic function in $\text{Im}(\omega) \geq 0$


1/11/2007

Kramers-Kronig

$$E(\omega) = 1 + \int_{-\infty}^{\infty} G(\tau) e^{i\omega\tau} d\tau \quad G(\tau) = \frac{1}{2\pi} \int_{-\infty}^{\infty} (E(\omega) - 1) e^{-i\omega\tau} d\omega$$

$$G(\tau) = 0 \text{ for } \tau < 0 \quad \vec{D}(\omega) = E(\omega) \vec{E}(\omega)$$

$$\vec{D}(t) = \vec{E}(t) + \int_{-\infty}^{\infty} G(\tau) \vec{E}(t-\tau) d\tau$$



$E(\omega)$ is analytic for $\text{Im}(\omega) \geq 0$

$$\text{Analytic Function } W(z) = U(x, y) + iV(x, y) \quad z = x + iy$$

derivative exists in any direction

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

$$\frac{dW}{dz} = \lim_{\Delta x \rightarrow 0} \frac{W(z+\Delta x) - W(z)}{\Delta x} = \lim_{\Delta y \rightarrow 0} \frac{W(z+i\Delta y) - W(z)}{i\Delta y}$$

$$\xrightarrow{\Delta x} \frac{1}{\Delta x} \left(\frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} \right) \Delta x = \xrightarrow{i\Delta y} \frac{1}{i\Delta y} \left(\frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right) \Delta y$$

$$\text{Real: } \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$$

$$\text{Im: } \frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}$$

Cauchy: $f(z)$ analytic (no singularities)

$$\oint_{\Gamma} \frac{f(z)}{z-a} dz = 2\pi i f(a)$$

inside and on Γ

$$\oint_{\Gamma} \frac{f(z)}{z-a} dz = \begin{cases} 2\pi i f(a) & a \text{ inside } \Gamma \\ \pi i f(a) & a \text{ on } \Gamma \\ 0 & a \text{ outside } \Gamma \end{cases} \quad (\text{Principal Part})$$

f analytic

$$\oint_{\Gamma} f(z) dz = 0 \quad \text{Or}$$

$$\xrightarrow{\text{no singularities}} \int_A^B f(z) dz = \int_A^B f(z) dz$$

$$\xrightarrow{\text{circle radius constant}} \int_{\Gamma} \frac{f(z)}{z-a} dz = \int_{\gamma} \frac{f(z)}{z-a} dz = \int_0^{2\pi} \frac{f(a+r e^{i\theta})}{r e^{i\theta}} r i e^{i\theta} d\theta$$

Taking the Principal Part

$$= \lim_{R \rightarrow \infty} \left(\int_{z_0}^{a-\delta} (\) dz + \int_{a+\delta}^{z_0} (\) dz \right)$$

$$\xrightarrow{\text{half circle } \gamma} \int_{U_\gamma}^\Gamma (\) dz + \int_\Gamma^\Gamma (\) dz = 0$$

$$\oint_{\Gamma} \frac{f(z)}{z-a} dz = \pi i f(a)$$

$$= i \int_0^{2\pi} f(a+r e^{i\theta}) d\theta \quad \text{take } \lim_{R \rightarrow \infty} \rightarrow 2\pi i f(a)$$

$$\oint_{\Gamma} (\) dz = 2\pi i f(a) = \int_{\Gamma} (\) dz + \int_{\gamma} (\) dz$$

$\int_{\gamma} (\) dz$: principal part $\rightarrow \pi i f(a)$

$\therefore \pi i f(a)$

$$\int_{\Gamma} \frac{E(\omega) - 1}{\omega - a} d\omega = 2\pi i' (E(a) - 1)$$

big circle

ω

$E(\omega) = 1 + \frac{1}{\alpha} \text{ for large } \alpha$

R : radius of circle

$\frac{E(\omega) - 1}{\omega} \sim \frac{1}{R^2}$ as $R \rightarrow \infty$

$E(\omega) - 1 = \int_0^\infty G(\tau) e^{i\omega\tau} d\tau \sim \int_0^\infty G(\tau) e^{-\alpha\tau} d\tau \sim G(0) \int_0^\infty e^{-\alpha t} dt \sim G(0) \frac{1}{\alpha}$

$\int_{\text{Circle}} \frac{1}{R^2} d\omega \rightarrow 0$ as $R \rightarrow \infty$

on real axis $E - 1 \rightarrow 0$ for $\omega \rightarrow \infty$ at least as $1/\omega^2$

$$\omega \in \mathbb{R} \quad E(\omega) = 1 + \frac{1}{i\pi} P \left(\int_{-\infty}^{\infty} \frac{E(\omega) - 1}{\omega - w} dw \right)$$

principal part

$$\text{Re}\{E(\omega)\} = 1 + \frac{1}{\pi} P \int_{-\infty}^{\infty} \frac{\text{Im}\{E(w)\}}{w - \omega} dw$$

$$\text{Im}\{E(\omega)\} = -\frac{1}{\pi} P \int_{-\infty}^{\infty} \frac{\text{Re}\{E(w)\} - 1}{w - \omega} dw$$

Complex conjugates

$$E^*(\omega^*) = E(-\omega) \quad \text{from} \quad E(\omega) = 1 + \int_0^\infty G(\tau) e^{i\omega\tau} d\tau$$

Kramers-Kronig Relations

$\text{Re}\{E\}$ is even in ω $\text{Im}\{E\}$ is odd


$$\frac{\text{Im}(E)(\omega^2 + \omega)}{(\omega^2 - \omega)(\omega^2 + \omega)} = \frac{\text{Im}(E)(\omega^2 + \omega)}{\omega^2 - \omega^2} \stackrel{\text{even}}{\underset{\text{even}}{=}} \Rightarrow \text{Re}\{E(\omega)\} = 1 + \frac{2}{\pi} P \int_0^\infty \frac{\omega \text{Im}\{E(\omega)\}}{\omega^2 - \omega^2} dw$$

$$\frac{(P\text{Re}(E) - 1)(\omega^2 + \omega)}{\omega^2 - \omega^2} \stackrel{\text{even}}{\underset{\text{even}}{=}} \Rightarrow \text{Im}\{E(\omega)\} = -\frac{2\omega}{\pi} P \int_0^\infty \frac{\text{Re}\{E(\omega)\} - 1}{\omega^2 - \omega^2} dw$$

1/16/2007

Generation of E.M. Waves by Oscillating Sources

Localized Sources



$$A(\vec{x}, t) = \rho(\vec{x}) e^{-i\omega t} \quad J(\vec{x}, t) = J(\vec{x}) e^{-i\omega t}$$

$$\vec{A}(\vec{x}, t) = \frac{1}{c} \int d^3x' \int dt' \frac{\vec{J}(\vec{x}', t')}{|\vec{x} - \vec{x}'|} \delta[t' - (t - \frac{|\vec{x} - \vec{x}'|}{c})]$$

$$= \frac{1}{c} \int d^3x' \frac{\vec{J}(\vec{x}')}{|\vec{x} - \vec{x}'|} e^{-i\omega t'} e^{i\vec{k}(\vec{x} - \vec{x}')} \quad \frac{\omega}{c} = k$$

$$\vec{A}(\vec{x}, t) = \vec{A}(\vec{x}) e^{-i\omega t}$$

$$\hookrightarrow = \frac{1}{c} \int d^3x' \frac{\vec{J}(\vec{x}')}{|\vec{x} - \vec{x}'|} e^{i\vec{k}(\vec{x} - \vec{x}')}$$

$$d \ll \lambda \quad \lambda = \frac{2\pi}{k} = \frac{2\pi c}{\omega} \quad \text{long wavelength limit}$$

1) $d \ll |\vec{x}| \ll \lambda$ Near field zone static fields

2) $d \ll |\vec{x}| \sim \lambda$ Intermediate zone

3) $d \ll \lambda \ll |\vec{x}|$ Radiation zone

$$\text{Near zone} \quad |\vec{x} - \vec{x}'| \ll \lambda \quad k|\vec{x} - \vec{x}'| \ll 1 \quad \therefore e^{i\vec{k}(\vec{x} - \vec{x}') \approx 1}$$

$$\vec{A}(\vec{x}, t) = \frac{1}{c} \left(\int d^3x' \frac{\vec{J}(\vec{x}')}{|\vec{x} - \vec{x}'|} \right) e^{-i\omega t}$$

static solution oscillating at ω

$$\text{Radiation zone} \quad |\vec{x}'| \gg |\vec{x}| \quad r = |\vec{x}| \quad \frac{1}{|\vec{x} - \vec{x}'|} \approx \frac{1}{r}$$

$$\vec{x} \xrightarrow{\vec{x} - \vec{x}'} \quad |\vec{x} - \vec{x}'| \approx r - \vec{x} \cdot \hat{n}$$

unit vector in \vec{x} dir : $\hat{n} = \frac{\vec{x}}{|\vec{x}|}$

$$k|\vec{x} - \vec{x}'| = kr - k\hat{n} \cdot \vec{x}'$$

$$\vec{A}(\vec{x}) = \frac{1}{c} \frac{e^{ikr}}{r} \int \vec{J}(\vec{x}') e^{-i(k\hat{n} \cdot \vec{x}')} d^3x'$$

"spherical wave" with direction-dependant amplitude

$$\vec{B} = \nabla \times \vec{A}$$

$$\nabla \times \vec{B} = \frac{1}{c} \frac{\partial \vec{E}}{\partial t} \quad (\vec{J}_{\text{heat}} = 0) \quad \vec{E} = \frac{i}{k} \nabla \times \vec{B}$$

$$= -i \frac{\omega}{c} \vec{E}$$

$$e^{-i\vec{k} \cdot \vec{x} \cdot \hat{n}} = 1 - ik \hat{n} \cdot \vec{x}$$

↓
electric dipole
radiation

→ magnetic dipole +
electric quadrupole
radiation

No monopole radiation : monopole can't 'pulse' and conserve charge

Electric Dipole $\vec{A}(\vec{x}) = \frac{1}{c} \frac{e^{ikr}}{r} \int \vec{J}(\vec{x}') d^3x'$

$$\left[\int \vec{J}(\vec{x}') d^3x' \right]_l = \int \vec{J}_l \underbrace{\frac{\delta_{ll}}{\delta x'_l}}_{\text{integration by parts}} d^3x' = - \int x_l (\nabla \cdot \vec{J}) d^3x' = -i\omega \int \rho(\vec{x}') x_l d^3x'$$

$$\therefore \int \vec{J}(\vec{x}') d^3x' = -i\omega \underbrace{\int \vec{x}' \rho(\vec{x}') d^3x'}_{\vec{p} \text{ electric dipole moment}} = -i\omega \vec{p}$$

$$\vec{A}(\vec{x}, t) = -ik \frac{e^{ikr}}{r} \vec{p} e^{-i\omega t}$$

Fields (radiation zone) $(\nabla \times \vec{A})_i \sim \epsilon_{\text{dm}} \partial_l \frac{e^{ikr}}{r} P_m$

$$\partial_l \left(\frac{e^{ikr}}{r} \right) = \frac{r i k e^{ikr} \partial_l(r) - e^{ikr} \partial_l(r)}{r^2} \quad \partial_l(r) = \frac{x_l}{r}$$

$$= \frac{e^{ikr}}{r^3} x_l (ikr - 1) \quad \begin{matrix} \text{in radiation} \\ kr \gg 1 \end{matrix} \quad \therefore ik \frac{e^{ikr}}{r} n_l = \partial_l \frac{e^{ikr}}{r}$$

$$(\nabla \times \vec{A})_i = -ik \epsilon_{\text{dm}} \frac{e^{ikr}}{r} ik n_l P_m$$

$$\nabla \times \vec{A} = \boxed{k^2 \frac{e^{ikr}}{r} (\hat{n} \times \vec{p}) e^{-i\omega t} = \vec{B}}$$

$$\vec{E} = \frac{i}{k} \nabla \times \vec{B}$$

$$\nabla \times \left(\frac{e^{ikr}}{r} (\hat{n} \times \vec{p}) \right) = ik \frac{e^{ikr}}{r} \hat{n} \times (\hat{n} \times \vec{p})$$

$$\vec{E} = -k^2 \frac{e^{ikr}}{r} \hat{n} \times (\hat{n} \times \vec{p}) = -\hat{n} \times \vec{B} = \boxed{\vec{B} \times \hat{n} = \vec{E}} \quad \vec{B} = k^2 \frac{e^{ikr}}{r} (\hat{n} \times \vec{p}) e^{-i\omega t}$$

Power radiated

$$\vec{S} = \frac{c}{4\pi} (\vec{E} \times \vec{B})$$

$$\vec{S} = \frac{1}{2} \frac{c}{4\pi} \text{Re} \{ \vec{E} \times \vec{B}^* \} \quad (\text{time averaged}) \quad \vec{E} \perp \vec{B} \quad |\vec{E}| = |\vec{B}| \quad \text{Re} \{ \vec{E} \times \vec{B}^* \} = |\vec{E} \times \vec{B}| = |\vec{E}|^2 = |\vec{B}|^2$$

$$\vec{S} = \frac{c}{8\pi} \frac{k^4}{r^2} |\hat{n} \times \vec{p}|^2 \hat{n}$$

power radiated into $d\Omega$

$$\frac{dP}{d\Omega} = \frac{c}{8\pi} k^4 |\hat{n} \times \vec{p}|^2$$

$$\frac{dP}{d\Omega} = \frac{c}{8\pi} k^4 |\vec{p}|^2 \sin^2 \theta$$


$$dP = \vec{S} \cdot \hat{n} r^2 d\Omega$$

Total power ($\int d\Omega$)

$$\int \sin^2 \theta d\Omega = \frac{8\pi}{3}$$

$$P_{\text{tot}} = \frac{c}{3} k^4 |\vec{p}|^2$$

$$P \propto k^4$$




Dipole
Radiation
Pattern

2nd term (magnetic dipole + electric quadrupole)

$$\vec{A}(\vec{x}) = -\frac{1}{c} ik \frac{e^{ikr}}{r} \left(\vec{j}(\vec{x}) (\hat{n} \cdot \vec{x}) \right) d^3x$$

$$\vec{M} = \frac{1}{2c} (\vec{x} \times \vec{j}) \quad \vec{m} = \int \vec{M} d^3x \quad \vec{A}(\vec{x}) = -\frac{ik}{2c} \frac{e^{ikr}}{r} \left((\vec{x} \times \vec{j}) \times \hat{n} \right) d^3x$$

$$\vec{A}(\vec{x}) = -ik \frac{e^{ikr}}{r} (\vec{m} \times \hat{\vec{n}})$$

$$\vec{B} = \nabla \times \vec{A} = k^2 \frac{e^{ikr}}{r} (\hat{n} \times \vec{m}) \times \hat{n} \quad \vec{E} = \vec{B} \times \hat{n}$$


polarization perpendicular
to \vec{m}, \hat{n} plane

1/18/2007

$$\int \left[(\hat{n} \cdot \vec{x}) \vec{J} + (\hat{n} \cdot \vec{J}) \vec{x} \right] d^3x = \int (n_\ell x_\ell J_i + n_\ell J_\ell x_i) d^3x$$

$$\left(n_e x_e J_m \frac{\partial x_i}{\partial x_m} + n_e J_e x_i \right) d^3x = \left(d^3x \left(-n_e \frac{\partial}{\partial x_m} (x_e J_m) x_i + n_e J_e x_i \right) \right)$$

$$\therefore = \int d^3x \left(-n_\ell x_\ell x_i \partial_m J_m \right) = \partial_\ell^A =$$

$$= -i\omega \int \rho(\hat{n} \cdot \vec{r}) x_i d^3x$$

$$\vec{A}(\vec{x}) = -\frac{1}{2} k^2 \frac{e^{ikr}}{r} \int d^3x' \delta(\vec{x}') (\hat{n} \cdot \vec{x}') \vec{d}$$

$$\vec{B} = \nabla \times \vec{A} \quad \nabla \times \left(\frac{e^{ikr}}{r} (\hat{n} \cdot \vec{x}) \vec{x} \right) = ik \hat{n} \times ()$$

$$\vec{B} = i k (\hat{n} \times \vec{A}) \quad \vec{E} = \vec{B} \times \hat{n}$$

$$\vec{B} = -\frac{i}{2} \vec{k}^3 \frac{e^{ikr}}{r} \int D(\vec{x}') (\hat{n} \cdot \vec{x}') (\hat{n} \times \vec{x}') d^3x' \quad B_e \sim \int \epsilon_{ijk} n_\alpha x'_j n_k x'_k D(x') d^3x'$$

$$\sim \frac{1}{3} \int d^3x' \underbrace{\epsilon_{\alpha\beta\gamma} n_\alpha n_\beta p^{(\gamma)}(x')}_{Q_{\delta k}} \left(3x'_\beta x'_k - r^2 \delta_{\beta k} \right)$$


$$B_l = -\frac{i}{6} k^3 \frac{e^{ikr}}{r} \epsilon_{ljk} n_a n_k Q_{jk}$$

$$\vec{B} = -\frac{i}{6} k^3 \frac{e^{ikr}}{r} (\hat{n} \times \vec{Q})$$

$$(\vec{Q})_k = \sum_E Q_{E k} n_k$$

power radiated will be k^6 for quadrupole (dipole power $\propto k^4$)

Example: Antenna



$$I(z,t) = I(z) e^{-i\omega t}$$

$$\text{Approx: } I(z) = I_0 \left(1 - \frac{2|z|}{d} \right)$$

(to satisfy continuity)

$$\text{Continuity: } \nabla \cdot \vec{J} = \text{cwA} \quad (\times \text{ cross-sectional area}) \rightarrow \frac{dI}{dz} = \text{cwA} \quad \uparrow \text{charge per unit length}$$


$$\frac{dI}{dz} = \pm I_0 \frac{2}{d} \quad z > 0$$

$$J(z) = -\frac{i}{\omega} + I_0 \frac{2}{d} = \pm \frac{2iI_0}{\omega d}$$

$$P = \int_{-\frac{d}{2}}^{\frac{d}{2}} z J(z) dz = \frac{2i I_0}{\omega d} \frac{1}{2} \frac{d^2}{4} 2 = \frac{i I_0 d}{2 \omega}$$

$$\text{Electric dipole} \quad \frac{dP}{d\Omega} = \frac{c}{8\pi} k^4 |\vec{p}|^2 \sin^2 \theta = \frac{c}{8\pi} k^4 \sin^2 \theta \quad \frac{I_0^2 d^2}{4 \omega^2} = \frac{I_0^2 d^2}{32\pi c} k^2 \sin^2 \theta$$

Scattering



$$\vec{E}_{in} = \vec{E}_{in} E_0 e^{i \vec{k}_{in} \cdot \vec{x}} e^{-i \omega t}$$

$$\vec{B}_{in} = \hat{n}_{in} \times \vec{E}_{in} \quad \hat{n} = \frac{\vec{n}}{|\vec{n}|}$$

$$\vec{E}_{\text{out}} = k^2 \frac{e^{ikr}}{r} (\hat{n} \times \vec{p}) \times \hat{n} \quad \vec{B}_{\text{out}} = \hat{n}_{\text{out}} \times \vec{E}_{\text{out}}$$

$$\frac{(\text{power radiated into } d\Omega, \hat{n} \text{ with polarization } \vec{E})}{(\text{incident current [power/m]}^2)} = d\sigma \text{ [area]}$$

$$= \frac{C}{8\pi} |\vec{E}_{out} \cdot \hat{\vec{r}}|^2 r^2 d\Omega$$

$$[\hat{n} \times \vec{p}] \times \hat{n} = \vec{p} \quad \vec{p} \parallel \hat{\epsilon}_n$$

$$= \frac{k^4 |\vec{E} \cdot \vec{p}|^2 d\Omega}{|E_0|^2} \quad |\vec{p}| \propto E_0 \Rightarrow \frac{d\sigma}{d\Omega} \text{ independent on } E_0$$

$$\frac{d\sigma}{d\Omega} = k^4 \frac{|\vec{E} \cdot \vec{p}|^2}{|E_0|^2}$$


$$\text{Dielectric Sphere} \quad \vec{p} = \frac{\epsilon - 1}{\epsilon + 2} a^3 \vec{E}_{in} \quad \vec{E}_{in} = \vec{E}_{in} E_0 e^{i \vec{k}_{in} \cdot \vec{x}} e^{-i \omega t}$$

$$\frac{d\sigma}{d\Omega} = k^4 \left(\frac{\epsilon-1}{\epsilon+2}\right)^2 a^6 |\vec{E} \cdot \vec{E}_{in}|^2$$

- need to average over incident polarization
- then sum over scattered polarization

Dielectric sphere continued

- average over incident polarization



$$a) \langle |\vec{E} \cdot \vec{E}_{in}|^2 \rangle = \langle \cos^2 \varphi \rangle = \frac{1}{2\pi} \int_0^{2\pi} \cos^2 \varphi d\varphi = \frac{1}{2}$$

 φ parametrizes direction \vec{E}_{in} in plane $\perp \vec{K}_{in}$

$$b) \langle |\vec{E} \cdot \vec{E}_{in}|^2 \rangle = \langle |\cos \theta \cos \varphi|^2 \rangle = \frac{1}{2} \cos^2 \theta$$

$$\frac{d\sigma_i}{d\Omega} = \frac{1}{2} k^4 a^6 \left(\frac{\epsilon - 1}{\epsilon + 2} \right)^2$$

$$\frac{d\sigma_{ii}}{d\Omega} = \frac{1}{2} k^4 a^6 \left(\frac{\epsilon - 1}{\epsilon + 2} \right)^2 \cos^2 \theta$$


- sum over polarization:

$$\frac{d\sigma}{d\Omega} = \frac{d\sigma_i}{d\Omega} + \frac{d\sigma_{ii}}{d\Omega} = \frac{1}{2} k^4 a^6 \left(\frac{\epsilon - 1}{\epsilon + 2} \right)^2 (1 + \cos^2 \theta)$$

$$\sigma_{tot} = \int \frac{d\sigma}{d\Omega} d\Omega = \frac{8\pi}{3} k^4 a^6 \left(\frac{\epsilon - 1}{\epsilon + 2} \right)^2 = \frac{\text{power out}}{\text{power in}} = \frac{\dot{N}_{out}}{\dot{N}_{in}}$$


1/23/2007

$$\theta = \frac{\pi}{2} \quad \frac{d\sigma_{ii}}{d\Omega} = 0$$

polarized \perp to scattering plane $\sigma \propto k^4 \sim \frac{1}{\lambda^4}$ shorter wavelengths are scattered more \rightarrow blue sky, red sunset

$$\dot{N}_{out} = \sigma_{tot} \frac{\dot{N}_{in}}{A} \quad N \rightarrow \# \text{ photons/time}$$

Collection of Scatterers (long wavelength)



$$1 \text{ scatterer: } \vec{E}_{out} = k^2 \frac{e^{ikr}}{r} (\hat{n} \times \vec{p}) \times \hat{n}$$

$$|\vec{x} - \vec{x}_j| \approx r - \vec{x}_j \cdot \hat{n}$$

$$\hat{n} = \frac{\vec{r}}{r}$$

$$\frac{1}{|\vec{x} - \vec{x}_j|} \approx \frac{1}{r} \quad e^{ik|\vec{x} - \vec{x}_j|} \approx e^{ikr} e^{-ik\vec{x}_j \cdot \hat{n}}$$

keep track of phases

$$\vec{p} \sim e^{i\vec{K}_{in} \cdot \vec{x}_j}$$

$$\vec{E}_{out} = k^2 \frac{e^{ikr}}{r} [(\hat{n} \times \vec{p}) \times \hat{n}] \sum_j e^{-ik\vec{x}_j \cdot \hat{n}} e^{i\vec{K}_{in} \cdot \vec{x}_j}$$

$$\vec{q} = \vec{K}_{out} - \vec{K}_{in}$$

$$\sum_j \rightarrow \sum_j e^{-i\vec{q} \cdot \vec{x}_j}$$

$$\frac{d\sigma}{d\Omega} = k^4 \frac{|\vec{E} \cdot \vec{p}|^2}{E_0^2} \mathcal{F}(\vec{q})$$

$$\mathcal{F}(\vec{q}) = \left| \sum_j e^{-i\vec{q} \cdot \vec{x}_j} \right|^2 \text{ form factor}$$

$$J = \int \epsilon(\vec{x}) d^3x$$

Fourier transform of distribution (squared)

$$\left| \int d^3x J e^{-i\vec{q} \cdot \vec{x}} \right|^2 = \mathcal{F}(\vec{q})$$

Scattering 2nd Diffraction 2nd

Scalar Diffraction Theory

$$\vec{E} = \vec{E}_0 e^{ikz} e^{-i\omega t}$$

Fresnel construction - each point is source of spherical waves

$$\psi = \text{component of } \vec{E}$$

$$\left[\frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \nabla^2 \right] \psi = 0 \quad \psi \sim e^{-i\omega t}$$

$$[\nabla^2 + k^2] \psi = 0$$

$$\text{Helmholtz Eq}$$


free space Green function

$$(\nabla^2 + k^2) G(\vec{x}, \vec{x}') = -4\pi \delta(\vec{x} - \vec{x}')$$

$$G(\vec{x}, \vec{x}') = \frac{e^{ikR}}{R}$$

$$R = |\vec{x} - \vec{x}'|$$

$$\tilde{R} = |\vec{x} - \vec{x}''|$$



$$G_D(\vec{x}, \vec{x}') = \frac{e^{ikR}}{R} - \frac{e^{ik\tilde{R}}}{\tilde{R}}$$

Solution of G
and is zero on screen

$$\int_V (G \nabla^2 \psi - 4 \nabla^2 G) d^3x' = \int_S \left(G \frac{\partial \psi}{\partial n} - 4 \frac{\partial G}{\partial n} \right) ds'$$

$$\text{LHS: } \int d^3x' \left(-G \psi k^2 - 4(-k^2 G - 4\pi \delta(\vec{x} - \vec{x}')) \right) = 4\pi \psi(\vec{x})$$

$$G \rightarrow \frac{e^{ikr}}{r} \quad 4 \rightarrow \frac{e^{ikr}}{r} \quad \text{for large } r \quad \int_S = \int_{\text{screen}} + \int_{\text{half-sphere}} \quad \text{for large } r \quad (\text{falls faster than } r^2)$$

RHS:

$$+ \int_{\text{Screen}} \psi(\vec{x}') \frac{\partial G_D}{\partial n'} ds'$$

inward
normal

$$R = |\vec{x} - \vec{x}'| = ((x - x')^2 + (y - y')^2 + (z - z')^2)^{1/2} \quad \tilde{R} = |\vec{x} - \vec{x}''| = ((x - x'')^2 + (y - y'')^2 + (z - z'')^2)^{1/2}$$

$$\Psi(\vec{x}) = \frac{E_0}{4\pi} \int_{\text{Hole}} dx' dy' \frac{\partial G_D}{\partial z'}$$

$$\frac{\partial}{\partial z'} \left(\frac{e^{ikR}}{R} \right) = \frac{e^{ikR}}{R} (z - z') \left(\frac{1}{R^2} - \frac{ik}{R} \right)$$

$$\frac{\partial}{\partial z'} \left(\frac{e^{ik\tilde{R}}}{\tilde{R}} \right) = \frac{e^{ik\tilde{R}}}{\tilde{R}} (z + z') \left(\frac{1}{\tilde{R}^2} - \frac{ik}{\tilde{R}} \right)$$

$$\left. \frac{\partial G_D}{\partial z'} \right|_{z'=0} = 2 \frac{e^{ikR}}{R} (-ik) \frac{z}{R}$$

$$\boxed{\Psi(\vec{x}) = \frac{-ik}{2\pi} \int_{\text{Hole}} \frac{e^{ikR}}{R} \frac{z}{R} \psi(\vec{x}', \vec{y}') dx' dy'}$$

$$\text{or } \Psi = \frac{-ik}{2\pi} E_0 \int_{\text{Hole}} \frac{e^{ikR}}{R} \frac{z}{R} dx' dy' \psi(\vec{x}', \vec{y}')$$

1/25/2007

$$\Psi(\vec{x}) = \frac{-ik}{2\pi} \iint_{\text{Hole}} \Psi(x', y') \frac{e^{ikr}}{R} \frac{z}{R} dx' dy'$$

Approx: $\frac{1}{R} \approx \frac{1}{r}$ $\frac{z}{R} \approx \cos\theta$ (looking far from hole)

$$\text{phase } kR \approx k(r - \vec{x} \cdot \hat{n}) \quad \Psi(\vec{x}) \approx \frac{-ik}{2\pi} E_0 \frac{e^{ikr}}{r} \cos\theta \int e^{-ik\vec{x} \cdot \hat{n}} dx' dy'$$

Circular Hole

$$\vec{x}' = (r' \cos\varphi, r' \sin\varphi, 0) \quad \hat{n} = (\sin\theta, 0, \cos\theta)$$

$$\vec{x}' \cdot \hat{n} = r' \sin\theta \cos\varphi$$

$$\int_0^a r' dr' \int_0^{2\pi} d\varphi e^{-ikr' \sin\theta \cos\varphi}$$


$$J_0(x) = \frac{1}{2\pi} \int_0^{2\pi} e^{-ix \cos\varphi} d\varphi$$

Bessel functions


$$\int x J_0(x) dx = x J_1(x)$$

$$\therefore \int_0^a r' 2\pi J_0(kr' \sin\theta) dr' = \frac{2\pi}{(k \sin\theta)^2} k \sin\theta J_1(k \sin\theta) = \frac{2\pi a^2}{k \sin\theta} J_1(k \sin\theta)$$

$u = kr' \sin\theta \quad du = k \sin\theta dr'$
 $r' = \frac{u}{k \sin\theta}$



$ka \gg 1$ intensity confined in forward direction
 $\psi \neq 0$ if $\theta \neq 0$




Diffraction pattern

first minimum $J_1(x) = 0$ for $x = 3.8$

$$\frac{2\pi}{\lambda} a \sin\theta = k \sin\theta \approx 3.8 \approx 1.2\pi$$

$$\Rightarrow \sin\theta = 0.6 \frac{\lambda}{a}$$

criterion for angular resolution



Problem 9.1

$\therefore \psi(x, t)$ is periodic $T = \frac{2\pi}{\omega}$

$$\psi(x, t) = \frac{1}{2} \psi_0(x) + \sum_{n=1}^{\infty} \operatorname{Re} \left\{ A_n(x) e^{-in\omega t} \right\}$$

Fourier series

$$A_n(x, \omega) = \frac{2}{T} \int_0^T \psi(x, t) e^{in\omega t} dt$$

$$\operatorname{Re} \left\{ \frac{2}{T} \int_0^T \psi(x, t) e^{in\omega t} e^{-in\omega t} dt \right\} = \frac{2}{T} \int_0^T dt \psi(x, t) \cos(n\omega t) \cos(n\omega t) + \frac{2}{T} \int_0^T dt \psi(x, t) \sin(n\omega t) \sin(n\omega t)$$


$A_n \cos(n\omega t) = A_n \cos\left(\frac{2\pi n t}{T}\right)$

$$f(t) = \frac{1}{2} \psi_0 + \sum_{n=1}^{\infty} A_n \cos\left(\frac{2\pi n t}{T}\right) + B_n \sin\left(\frac{2\pi n t}{T}\right)$$

Fourier decomposition of $f(t)$

calculate multipole moments using A_n

one charge



$$\vec{x}_q = (R \cos \omega t, R \sin \omega t, 0)$$

$$\omega = \omega t$$

$$\rho(\vec{r}, t) = q \delta(\vec{r} - \vec{x}_q)$$

$$\rho(r, \varphi, t) = q \delta(r - R) \delta(\varphi - \omega t) \frac{1}{r} S(z) \quad \text{Cylindrical coords}$$

$$A_n(r, \varphi, z) = \frac{q}{r} S(r - R) S(z) \frac{2\omega}{2\pi} \int_0^{2\pi} \delta(\varphi - \omega t) e^{in\varphi} dt = \frac{1}{\pi} \frac{q}{r} S(r - R) S(z) e^{in\varphi}$$

$$Q_{lm} = \sum_{n=0}^{\infty} r^l \int Y_{lm}^* d^3x \quad Q_{00} = \frac{1}{\sqrt{4\pi}} \frac{1}{2} \frac{q}{\pi} \int r dr \left(\int dz \int_0^{2\pi} d\varphi \frac{S(r-R)}{r} S(z) e^{in\varphi} \right)$$

$n=0$ otherwise $\int e^{in\varphi} = 0$

$$Q_{00} = \frac{1}{\sqrt{4\pi}} q \quad \text{for } n=0 \quad Q_{00} = 0 \quad \text{for } n \neq 0$$

$n=0 \rightarrow$ no radiation

$l=1 \quad e^{-i\varphi} \quad m=1 \rightarrow$ for $n=1$ this is only nonzero term

$$\begin{matrix} Y_{1m}^* & e^{-i\varphi} \\ 1 & m=-1 \\ 1 & m=0 \end{matrix}$$

$$Q_{11} = \int dr r \int dz \int d\varphi \sqrt{\frac{3}{8\pi}} \sin \theta e^{-i\varphi} r e^{\frac{S(r-R)}{r}} S(z) e^{i\varphi} \frac{1}{\pi}$$

$$Q_{11} = -\sqrt{\frac{3}{8\pi}} 2qR \quad \text{dipole radiates at freq. } \omega \quad (n=1)$$

$$Q_n = -\sqrt{\frac{3}{8\pi}} (P_x - i P_y) \quad Q_{1-1} = \sqrt{\frac{3}{8\pi}} (P_x + i P_y) \quad P_x = qR \quad P_y = iqR$$

Problems concerning Diffraction Gratings

$$\frac{1}{d} N \text{ slits} \cdot P \quad \psi \sim \sum_{m=0}^{N-1} e^{im\theta} = \frac{1 - e^{i(N-1)\theta}}{1 - e^{i\theta}} \quad \text{ignoring single slit contribution}$$

distance large compared to size

$$d \left\{ \int_{-\frac{\lambda}{2} \sin \theta}^{\frac{\lambda}{2} \sin \theta} e^{ikr} \right\} \rightarrow e^{ikds \sin \theta} \quad kds \sin \theta = \delta$$

Diffraction from 1 slit

$$\int e^{-ikx \sin \theta} dx dy \cos \theta \sim \cos \theta \int_{-\frac{\lambda}{2} \sin \theta}^{\frac{\lambda}{2} \sin \theta} dy e^{-iky \sin \theta} = \cos \theta \frac{i \frac{\lambda}{2}}{\frac{\lambda}{2} \sin \theta} 2 \sin(k \frac{\lambda}{2} \sin \theta)$$

single dimension

translational invariance along x (?)

$$\sim \cos \theta \frac{\sin \alpha}{\alpha} \quad \alpha = k \frac{a}{2} \sin \theta$$

Grating with thermal noise spacing average d but Gaussian dist'n

$$\int_{-\frac{\lambda}{2} + \epsilon}^{\frac{\lambda}{2} + \epsilon} dy e^{-iky \sin \theta} \quad P(\epsilon) \propto e^{-\frac{\epsilon^2}{2\sigma^2}}$$

$$\langle e^{iky \sin \theta (E_m - E_n)} \rangle$$

Special Relativity

1/30/2007

Invariance

Newton's laws are invariant under Galilean transformation

 \leadsto can't tell if moving

$$\frac{dp}{dt} = -\frac{dU}{dx} \quad p=p(t) \quad U=U(x) \quad p=m \frac{d}{dt} x_p(t)$$

position of particle

$$x' = x - vt \quad t' = t \quad \Rightarrow \quad x = x' + vt' \quad t = t' \quad x = x(x', t') \quad t = t(x', t')$$

$$x'_p = x_p - vt \quad \frac{d}{dt'} = \frac{d}{dt} \frac{dt}{dt'} + \frac{d}{dx} \frac{dx}{dt} = \frac{d}{dt'}$$

$$\frac{d}{dx'} = \frac{d}{dx} \frac{dx}{dx'} + \frac{d}{dt'} \frac{dt}{dx} = \frac{d}{dx}$$

in this case (particle position not fit of position)
not a field

$$U(x(x')) = U'(x')$$

Ex: $U = \frac{1}{2} k x^2$
 $U'(x') = \frac{1}{2} k (x' + vt)^2$

U invariant

$$p' = p - mv \quad \frac{d}{dt'}(p' + mv) = -\frac{d}{dx'} U'(x') \quad \frac{dp'}{dt'} = -\frac{dU(x)}{dx'}$$

speed of trans = constant

$$m \ddot{x}_p = -k x_p$$

$$\frac{d^2}{dt'^2}(x'_p + vt) = -\frac{k}{m}(x'_p + vt) \quad \frac{d^2 x'_p}{dt'^2} = -\frac{k}{m} x'_p - \frac{k}{m} vt$$

the differential equation changes

→ choosing a frame with spring stationary makes eqn's easier

Wave Eq: → not invariant, not a problem - not fundamental eq.

$$\frac{1}{c^2} \frac{\partial^2}{\partial t'^2} \psi(x', t') - \frac{\partial^2}{\partial x'^2} \psi(x', t') = 0 \quad \psi \text{ pressure field (sound waves)}$$

$$\psi \text{ invariant} \quad \psi(x(x'), t(t')) = \psi'(x', t') \quad \text{same function regardless of frame}$$

$$\frac{\partial}{\partial x} = \frac{\partial}{\partial x'} \quad \frac{\partial}{\partial t} = \frac{\partial}{\partial t'} - v \frac{\partial}{\partial x'}$$

$$\frac{1}{c^2} \left(\frac{\partial^2}{\partial t'^2} - v \frac{\partial^2}{\partial x'^2} \right) \left(\frac{\partial}{\partial t'} - v \frac{\partial}{\partial x'} \right) \psi' - \frac{\partial^2}{\partial x'^2} \psi' = 0$$

→ there is a preferred frame ($v=0$)

$$\left(\frac{1}{c^2} \frac{\partial^2}{\partial t'^2} - \left(1 - \frac{v^2}{c^2} \right) \frac{\partial^2}{\partial x'^2} - \frac{2v}{c^2} \frac{\partial^2}{\partial t' \partial x'} \right) \psi' = 0$$

Form of wave eq not invariant

Schrödinger Eq:

$$i \frac{\partial \psi}{\partial t} = -\frac{1}{2m} \frac{\partial^2}{\partial x'^2} \psi + V(x') \psi \quad \psi' = \psi e^{i \theta(\psi) \varphi(x')}$$

→ invariant

$$\frac{d\vec{p}}{dt} = -\nabla U \quad \text{invariant under rotations}$$

$$U \rightarrow \text{scalar} \quad U(x(x')) = U'(x') \quad \vec{p} \rightarrow \text{vector} \quad P'_1 = P_1 \cos \theta + P_2 \sin \theta \quad \text{not true that } P'_1 = P_1$$

(ex.)

$$\text{Lagrangian} \quad \mathcal{L} = \sum_i \left\{ \frac{1}{2} m \dot{x}_i^2 \right\} - \frac{1}{2} \sum_{i \neq j} \left\{ V(|\vec{x}_i - \vec{x}_j|) \right\}$$

invariant under translation $x' = x - \lambda \quad t' = t$

$$V((x'_i + \lambda) - (x'_j + \lambda)) = V(\vec{x}'_i - \vec{x}'_j) \quad \mathcal{L}(\dot{x}, x) = \mathcal{L}(\dot{x}, x - \lambda) \quad \text{invariant under translation}$$

corresponds to total momentum conservation

$$\frac{dp_{\text{tot}}}{dt} = 0 \quad \vec{P}_{\text{tot}} = \sum_i \vec{p}_i \quad p_i = \frac{\partial \mathcal{L}}{\partial \dot{x}_i}$$

$$\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{x}_i} - \frac{\partial \mathcal{L}}{\partial x_i} = 0 \quad \frac{dp_i}{dt} = \frac{\partial \mathcal{L}}{\partial x_i} \quad \frac{dp_{\text{tot}}}{dt} = \frac{d}{dt} \sum p_i = \sum_i \frac{\partial \mathcal{L}}{\partial x_i} = \sum_i (-\frac{1}{2}) \frac{\partial}{\partial x_i} \sum_{l \neq i} V(|\vec{x}_i - \vec{x}_l|)$$

$$V(|\vec{x}_i - \vec{x}_l|) = V(z) \quad z = |\vec{x}_i - \vec{x}_l|$$

$= 0 \quad \sum_i (\text{force on particle } i \text{ due to all others})$

by Newton's 3rd law all cancel

$$\frac{\partial V}{\partial x_2} = \frac{\partial V}{\partial z} \frac{\partial z}{\partial x_2} = -\frac{\partial V}{\partial z} \quad \frac{\partial V}{\partial x_1} = \frac{\partial V}{\partial z} \quad \sum_i \frac{\partial V}{\partial x_i} = 0$$

$\sum_i \frac{\partial V}{\partial x_i} = 0$


If $\mathcal{L}(x) = \mathcal{L}(x - \lambda) + \lambda$

$$\mathcal{L}(x - \lambda) \approx \mathcal{L}(x) - \underbrace{\lambda \sum_i \frac{\partial \mathcal{L}}{\partial x_i}}_{\text{is zero}}$$

Postulates

1) Physics is same in different inertial systems

2) Speed of light independent of relative motion of source and observer



Seek transformation so that $(ct)^2 - |\vec{x}|^2 = s^2$ is invariant

$$x' = \alpha x + \beta t \quad t' = \gamma x + \delta t$$

$$x' = \alpha(x - vt) \quad t' = \gamma(t - \frac{x}{v}) \quad (ct')^2 - x'^2 = (c\gamma(t - \frac{x}{v}))^2 - (\alpha(x - vt))^2 = ct^2 - x^2$$

$$\alpha = \gamma \quad \beta = \gamma \quad \gamma = \frac{1}{\sqrt{1-\beta^2}}$$

$$\gamma = \frac{v}{c} \quad \gamma = \frac{1}{\sqrt{1-\beta^2}} \quad x' = \gamma(x - \beta ct) \quad ct' = \gamma(ct - \beta x)$$


$$x_0 = ct \quad x_1 = x$$

$$x'_0 = \gamma(x_0 - \beta x_1) \quad x'_1 = \gamma(x_1 - \beta x_0)$$

$$A^\alpha = (A^0, \vec{A}) \quad A^0 - |\vec{A}|^2 = A \cdot A = A_\alpha A^\alpha \quad \text{modulus of this vector is invariant}$$

\rightarrow length of 4-vector

$$A \cdot B = A_\alpha B^\alpha = A^0 B^0 - \vec{A} \cdot \vec{B}$$



Event = point in space time

$$s^2 = c^2(t_1 - t_2)^2 - |\vec{x}_1 - \vec{x}_2|^2$$

- can be causally connected $\rightarrow s^2 > 0$ "time-like" separation $\rightarrow \exists$ Lorentz transformation such that events occur at same place (future/past cone)
- not causally connected $\rightarrow s^2 < 0$ "space-like" separation $\rightarrow \exists$ Lorentz transformation such that events occur at same time (elsewhere)
- $s^2 = 0$ "light-like" separation (on light cone)

2/1/2007

Maxwell Eqs. — not invariant under Galilean \rightarrow these are fundamental eqs.

\rightarrow can find out if you are moving ... but you can't \downarrow

Source Receiver

$\xrightarrow{\text{At rest}}$)

M.E.: speed of light = c

Mechanics: speed of sound = c_s

Source moves at vel. v

M.E.: speed of light = c

Mechanics: speed of sound = c_s

same as before


Receiver moves at v

M.E.: measure: c

Mechanics: measure: $c_s - v$

\rightarrow can't tell if you are moving

\rightarrow consistent with principle of relativity
but not with Galilean addition of velocities



$$K: d\vec{x} = \vec{u} dt \quad ds^2 = c^2 dt^2 - dx^2 = c^2 dt^2 - u^2 dt^2 = (c^2 - u^2) dt^2 = c^2 dt^2 (1 - \beta^2)$$

$$K': dx'^2 = 0 \quad ds'^2 = c^2 dz'^2 \quad ds^2 = ds'^2 \quad c^2 dz'^2 = \gamma^2 dt^2 (1 - \beta^2) \quad dz = dt \sqrt{1 - \beta^2}$$

$$d\tau = \frac{dt}{\gamma} \quad \text{proper time - time in rest frame of particle}$$

\hookrightarrow is the shortest

$$\tau_2 - \tau_1 = \int_{t_1}^{t_2} \frac{dt}{\gamma(t)} < t_2 - t_1 \quad \text{since } \gamma \geq 1$$

Doppler Shift

phase is invariant — measure of how many waves \rightarrow # does not change

$$E \sim e^{i(E \cdot \vec{x} - \omega t)}$$

$$\vec{R} \cdot \vec{x} - \omega t = \vec{R} \cdot \vec{x}' - \omega t'$$

$$4\text{-vector: } (\frac{\omega}{c}, \vec{k})$$

\therefore product of 2 4-vectors

$$\frac{\omega'}{c} = \gamma \left(\frac{\omega}{c} - \beta \cdot \vec{k} \right)$$

(if $\beta \parallel \vec{k}$: $\beta \cdot \vec{k} = \beta k_z$)

$$k_{||}' = \gamma (k_{||} - \beta \frac{\omega}{c})$$

$\parallel \rightarrow \beta$

$$\vec{k}_{\perp}' = \vec{k}_{\perp}$$

$\perp \rightarrow \beta$

$$|\vec{k}| = \frac{\omega}{c} \quad |\vec{k}'| = \frac{\omega'}{c} \quad \text{for light}$$

$$\omega' = \gamma (\omega - v \frac{\omega}{c}) \quad \text{... for } \beta \parallel \vec{k} \quad \omega' = \gamma (1 - \frac{v}{c}) \omega = \frac{1 - \frac{v}{c}}{\sqrt{1 - \frac{v^2}{c^2}}} \omega = \boxed{\sqrt{\frac{1 - \frac{v}{c}}{1 + \frac{v}{c}}} \omega} \quad \omega' = \omega'$$

$$\vec{\beta} \perp \vec{k} \quad \omega' = \gamma \omega$$

Energy & Momentum

$(\frac{E}{c}, \vec{p})$ is a 4-vector

$$E = \gamma mc^2 \quad m - \text{rest mass} \quad \vec{p} = \gamma m\vec{v} \quad \vec{v} = \frac{d\vec{x}}{dt}$$

$\frac{E^2}{c^2} - |\vec{p}|^2$ is invariant

$$\hookrightarrow \text{in rest frame} = \frac{(mc^2)^2}{c^2} = m^2 c^2 \quad E^2 = p^2 c^2 + m^2 c^4$$

ultrarelativistic γ is large $\gamma \gg 1 \quad v \approx c \quad p \approx \gamma mc$

$$\frac{E^2}{c^2} - \gamma^2 m^2 c^2 = m^2 c^2 \quad \frac{E^2}{c^2} = p^2 + m^2 c^2 \approx p^2 \Rightarrow E \approx pc \quad \text{at high energies}$$

$$\text{At low energies, } \gamma \approx 1 \quad \beta \ll 1 \quad E = c \sqrt{p^2 + m^2 c^2} = mc^2 \sqrt{1 + \frac{p^2}{m^2 c^2}} \approx mc^2 \left(1 + \frac{1}{2} \frac{p^2}{m^2 c^2} \right)$$

$$E \approx mc^2 + \frac{p^2}{2m} \quad \text{at low energies}$$

Different dispersion relations: $E = E(p)$

$$\frac{d\vec{p}}{dt} = q(\vec{E} + \vec{B} \times \vec{v})$$

Space-Time Geometry

$$x^\alpha \quad \alpha = 0, 1, 2, 3 \quad x^\alpha = x^\alpha(x^0, x^1, x^2, x^3) \quad s = s \quad \text{scalar}$$

$$A^\alpha = \frac{\partial x^\alpha}{\partial x^\beta} A^\beta \quad \text{contravariant} \quad A_\alpha = \frac{\partial x^\beta}{\partial x^\alpha} A_\beta \quad \text{covariant}$$

$$A^\alpha = (A^0, \vec{A}) \quad B^\alpha = (B^0, \vec{B}) \quad A \cdot B = A^0 B^0 - \vec{A} \cdot \vec{B} \quad A_\alpha = (A^0, -\vec{A})$$

$$\text{metric tensor: } g_{\alpha\beta} = \begin{pmatrix} 1 & -1 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \quad A \cdot B = g_{\alpha\beta} A^\alpha B^\beta \quad A_\alpha = g_{\alpha\beta} A^\beta \quad \therefore A \cdot B = A_\alpha B^\alpha$$

$$ds^2 = g_{\alpha\beta} dx^\alpha dx^\beta \quad \text{distance} \quad T_{\alpha\beta} = g_{\beta\gamma} g_{\alpha\delta} T^{\gamma\delta}$$

2/6/2007

$$|A|^2 = A \cdot A = A_\alpha A^\alpha = g_{\alpha\beta} A^\alpha A^\beta = g^{00} A_0 A_0 = A^0 - |\vec{A}|^2 \quad g_{\alpha\beta} = g^{\alpha\beta}$$

$$g_{\alpha\beta} g^{\beta\gamma} = \delta_\alpha^\gamma$$

Operator corresponding to ∇

$$\frac{\partial}{\partial x^\alpha} \equiv \partial_\alpha \quad \frac{\partial}{\partial x^\alpha} = \frac{\partial x^\beta}{\partial x^\alpha} \frac{\partial}{\partial x^\beta} \quad \text{transforms like } A_\alpha \quad \partial_\alpha = \left(\frac{\partial}{\partial x^0}, \nabla \right) \quad \partial^\alpha = \left(\frac{\partial}{\partial x^0}, -\nabla \right)$$

$$\partial^\alpha \partial_\alpha = \frac{\partial^2}{\partial x^2} - \nabla^2 \equiv \square \quad \text{operator for wave equation:} \quad \square \psi = 0$$

$$\text{"Divergence": } \partial^\alpha A_\alpha = \frac{\partial A^0}{\partial x^0} + \nabla \cdot \vec{A} \quad \text{Lorentz invariant} \quad \partial^\alpha A_\alpha = \partial_\alpha A^\alpha$$

Current 4-vector

$$J^\alpha = (c\rho, \vec{J}) \quad \partial_\alpha J^\alpha = \frac{\partial \rho}{\partial t} + \nabla \cdot \vec{J} = 0 \quad \text{continuity Eq. } \partial_\alpha J^\alpha = 0$$

$$\frac{c \partial \rho}{c \partial t}$$

$$\text{In Lorentz gauge: } \frac{1}{c} \frac{\partial \phi}{\partial t} + \nabla \cdot \vec{A} = 0 \rightarrow \partial_\alpha A^\alpha = 0 \quad J^\alpha = (\phi, \vec{J})$$

$$\left(\frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \nabla^2 \right) \frac{\phi}{c} = 4\pi \frac{J^\alpha}{c} \quad \therefore \quad A^\alpha = (\phi, \vec{A}) \quad \square A^\alpha = \frac{4\pi}{c} J^\alpha$$

A Lorentz transformation

 $T^{\alpha\beta}$ tensor

$$T' = AT^{-1}A$$

$${}^T A_{ij} = A_{ji} \quad \text{transpose}$$

$$A_{ij}^+ = A_{ji}^* \quad \text{hermitian conjugate}$$

$$\text{symmetric } \vec{A} = \vec{A} \quad \text{hermitian } A^+ = A$$

real numbers: orthogonal matrix ${}^T A = A^{-1}$ preserve scalar products

$$\text{antisymmetric } TA = -A$$

$$a = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}$$

$$a \cdot b = (a) \begin{pmatrix} b \end{pmatrix} = {}^T ab$$

$$a' = Aa \quad b' = Ab$$

$$a' \cdot b' = {}^T a' \cdot b' = {}^T a^T A A b = {}^T ab \quad \text{if } {}^T A A = I \quad {}^T A = A^{-1}$$

$$\text{unitary matrix } A^+ = A^{-1} \quad {}^T(Aa) = {}^T a$$

$$a \cdot b = a^+ b$$

Tensor T_{ik} under orthogonal transf.

$$b = Ta \quad b' = T' a \quad T' = ?$$

$$b' = T' a' \Rightarrow Ab = T' Aa$$

$$a' = Aa \quad b' = Ab$$

$$b = A^{-1} {}^T A a \quad T = A^{-1} {}^T A \quad T' = A T A^{-1}$$

$$\text{if } A \text{ is orthogonal: } T' = AT^{-1}A$$

$$A \text{ is Lorentz transformation} \quad a' = Aa \quad b' = Ab \quad a = () \quad 4\text{-vector}$$

$$a \cdot b = {}^T(ga) b = a' \cdot b' = {}^T(gAa) Ab = {}^T a^T(gA) Ab = {}^T a^T Ag A b \quad {}^T g = g$$

$$\hookrightarrow {}^T a^T Ag A b = {}^T a^T Ag A b \quad \therefore {}^T Ag A = g \quad \text{for a Lorentz transformation}$$

$$g g = I$$

$$g^T Ag A = I \quad \therefore [g^T Ag = A^{-1}]$$

$$b^\alpha = T^{\alpha\beta} a_\beta \quad b'^\alpha = {}^T a^\beta a'_\beta \quad a_\beta = g_{\beta\gamma} a^\gamma$$

$$b = Tg a \quad a = () \quad b = () \quad a' = Aa \quad b' = Ab$$

$$b' = {}^T g a' \quad Ab = {}^T g Aa \quad b = A^{-1} {}^T g Aa \quad A^{-1} {}^T g A = Tg$$

$$T' g = ATg A^{-1} \quad T' = ATg A^{-1} g = AT^{-1}A \quad \therefore \quad T' = AT^{-1}A \quad \text{proved}$$

x left A
right A⁻¹ both sides

$$\vec{E} = -\nabla \phi - \frac{1}{c} \frac{\partial \vec{A}}{\partial t} \quad \vec{B} = \nabla \times \vec{A}$$

$$\partial^\alpha = \left(\frac{1}{c} \frac{\partial}{\partial t}, -\nabla \right)$$

$$E_x = -\frac{\partial \phi}{\partial x} - \frac{1}{c} \frac{\partial A_x}{\partial t} = +\partial^1 A^0 - \partial^0 A^1 = -(\partial^0 A^1 - \partial^1 A^0) \quad E_y = -(\partial^0 A^2 - \partial^2 A^0)$$

$$B_x = \frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z} = -\partial^2 A^3 + \partial^3 A^2 = -(\partial^2 A^3 - \partial^3 A^2) \quad B_y = \partial^1 A^3 - \partial^3 A^1$$

$$F^{\alpha\beta} = \partial^\alpha A^\beta - \partial^\beta A^\alpha = \begin{pmatrix} 0 & -E_x & -E_y & -E_z \\ E_x & 0 & -B_z & B_y \\ E_y & B_z & 0 & -B_x \\ E_z & -B_y & B_x & 0 \end{pmatrix}$$

antisymmetric

$$F' = A F^T A$$

Lorentz transf. with v along x

$$\tilde{x}^0 = \gamma(x^0 - \beta x_1) \quad \tilde{x}^1 = \gamma(x^1 - \beta x^0) \quad \tilde{x}^2 = x^2 \quad \tilde{x}^3 = x^3 \quad (\tilde{x}) = A(x)$$

$$A = \begin{pmatrix} \gamma & -\gamma\beta & 0 & 0 \\ -\gamma\beta & \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

2/8/2007

$$F' = \begin{pmatrix} \gamma & -\gamma\beta & 0 & 0 \\ -\gamma\beta & \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & -E_x & E_y & 0 \\ . & . & . & 0 \\ . & . & . & 0 \end{pmatrix} \begin{pmatrix} \gamma & -\gamma\beta & 0 & 0 \\ -\gamma\beta & \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$


$x \rightarrow 1$
 $y \rightarrow 2$
 $z \rightarrow 3$

$$F' = \begin{pmatrix} 0 & -E_1 & -\gamma E_2 + \gamma B B_3 & -\gamma E_3 - \gamma B B_2 \\ 0 & \gamma B E_2 - \gamma B_3 & \gamma B E_3 + \gamma B_2 & 0 \\ \text{antisymmetric} & 0 & 0 & -B_1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$E'_1 = E_1, \quad E'_2 = \gamma(E_2 - \beta B_3), \quad E'_3 = \gamma(E_3 + \beta B_2)$$

$$B'_1 = B_1, \quad B'_2 = \gamma(B_2 + \beta E_3), \quad B'_3 = \gamma(B_3 - \beta E_2)$$

Fields of a charge in uniform motion:

K  $\vec{v} = v \hat{x}_1$, P: observation point in K $P = (0, b, 0)$

K': rest frame of charge

$$x'_1 = \gamma(c_0 - \beta c t)$$

$$ct' = \gamma(ct - (c_0))$$

$$P' = (-\gamma v t, b, 0) = (-v t', b, 0)$$

fields in K': $\vec{E} = q \frac{\hat{r}_1}{r'^2}$ $r'^2 = (v t')^2 + b^2$

$$E'_1 = q \frac{-v t'}{((v t')^2 + b^2)^{3/2}} \quad E'_2 = q \frac{b}{((v t')^2 + b^2)^{3/2}} \quad E'_3 = 0 \quad \vec{B}' = 0$$

in K:

$$E_1 = E'_1 \quad E_2 = \gamma E'_2 \quad E_3 = \gamma E'_3 = 0 \quad B_1 = B_2 = 0 \quad B_3 = \gamma B E'_2 = \beta E_2$$

$$E_1 = \frac{-q \gamma v t}{((\gamma v t)^2 + b^2)^{3/2}} \quad E_2 = \frac{q \gamma b}{((\gamma v t)^2 + b^2)^{3/2}} \quad B_3 = \frac{q \gamma \beta b}{((\gamma v t)^2 + b^2)^{3/2}}$$

$$t=0 \quad E_x = E_z = 0 \quad E_y = \gamma \frac{q}{b^2} \quad \text{for } \gamma \gg 1 \quad \beta \approx 1 \quad B_2 \approx E_y \quad \text{looks like flash of a plane wave}$$

short times $t \ll \frac{b}{\gamma v}$ $E_x \approx \frac{q}{b^2} \left(\frac{-\gamma vt}{b} \right) = \gamma \frac{q}{b^2} \left(\frac{-vt}{b} \right)$ $E_y \approx \gamma \frac{q}{b^2}$


→ see Coulomb field enhanced by γ

enhanced Coulomb $\sim \cos \theta$




long times $t \gg \frac{b}{\gamma v}$ $E_x \approx \frac{1}{\gamma^2} \frac{q}{(vt)^2}$

→ see Coulomb field suppressed by $\frac{1}{\gamma^2}$



E_x averages out since it changes sign



$$t' = \gamma(t - \frac{v}{c^2}x) \quad x' = \gamma(x - vt)$$

$$dx = \gamma(dx' + vdt') \quad dt = \gamma(dt' + \frac{v}{c^2}dx')$$

$$\frac{dx}{dt} = \frac{\frac{dx'}{dt'} + v}{1 + \frac{v}{c^2} \frac{dx'}{dt'}}$$

$$u = \frac{u' + v}{1 + \frac{vu}{c^2}}$$

$$\frac{d}{dt} \frac{dx}{dt} = \frac{dt'}{dt} \frac{d}{dt'} \left(\frac{dx}{dt'} \right) + \frac{dx'}{dt} \frac{d}{dx'} \left(\frac{dx}{dt'} \right)$$

$$\frac{dt'}{dt} = \gamma \left(1 - \frac{v}{c^2} \frac{dx}{dt} \right)$$

$$\text{if } \frac{dx'}{dt'} = 0 \quad (\text{frame of moving particle}) \quad \frac{dx}{dt} = v$$

$$\frac{dt'}{dt} = \frac{1}{\gamma} \quad dt' = \frac{dt}{\gamma}$$

$$\frac{d}{dt} \frac{dx}{dt} = \gamma \left(1 - \frac{v}{c^2} \frac{dx}{dt} \right) \frac{\left[(1 + \frac{v}{c^2} \frac{dx}{dt}) \frac{d^2x'}{dt'^2} - (\frac{dx'}{dt'} + v) \frac{v}{c^2} \frac{d^2x'}{dt'^2} \right]}{(1 + \frac{v}{c^2} \frac{dx}{dt})^2} = \frac{1}{\gamma^3} \frac{\frac{d^2x}{dt^2}}{(1 + \frac{v}{c^2} \frac{dx}{dt})^3}$$

Moving Charges (Chpt 14)

2/15/2007

Potential of a moving charge $\phi(\vec{x}, t) = \int d^3x' \frac{\rho(\vec{x}', t')}{|\vec{x} - \vec{x}'|} \quad t' = t - \frac{|\vec{x} - \vec{x}'|}{c}$

charge q with vel. \vec{v}

$$\text{trajectory } \vec{x}_0(t) \text{ of charge} \quad \rho(\vec{x}', t') = q \delta(\vec{x}' - \vec{x}_0(t'))$$

det(Jacobian)

$$\phi(\vec{x}, t) = \int q \frac{\delta(\vec{x} - \vec{x}_0(t'))}{|\vec{x} - \vec{x}'|} d^3x' \quad \begin{matrix} \text{change} \\ \text{of variables} \end{matrix} \quad \vec{y} = \vec{x}' - \vec{x}_0(t')$$

$$\phi = \int d^3y \det(J) \frac{q \delta(\vec{y})}{|\vec{x} - \vec{y} - \vec{x}_0(t')|}$$

$$\phi = \det(J)(\vec{x}' = \vec{x}_0(t')) \frac{q}{|\vec{x} - \vec{x}_0(t')|} = \int \det(J)(\vec{x}') \frac{q}{|\vec{x} - \vec{x}'|} \delta(\vec{x}' - \vec{x}_0(t')) d^3x'$$

$$J_{ij} = \left(\frac{\partial y_i}{\partial x_j} \right)^{-1} \quad \text{inverse of matrix}$$

$$\det \left(\left(\frac{\partial y_i}{\partial x_j} \right)^{-1} \right) = \frac{1}{\det \left(\frac{\partial y_i}{\partial x_j} \right)}$$

$$\frac{\partial y_i}{\partial x_j} = \delta_{ij} + \frac{\partial x_{0i}(t')}{\partial x_j}$$

$$\frac{\partial x_{0i}(t')}{\partial x_j} = \frac{d x_{0i}(t')}{dt'} \frac{\partial t'}{\partial x_j} = C \beta_i(t') \frac{1}{c} n_j = \beta_i n_j$$

jet of particle at retarded time

$$\frac{\partial y_i}{\partial x_j} = \delta_{ij} - \beta_i n_j$$

$$\frac{\partial t'}{\partial x_j} = -\frac{1}{c} \frac{\partial}{\partial x_j} \left((x_1 - x'_1)^2 + (x_2 - x'_2)^2 + \dots \right)^{\frac{1}{2}} = +\frac{1}{c} \frac{1}{2} \frac{2(x_j - x'_j)}{|\vec{x} - \vec{x}'|}$$

$$n_j = \frac{x_j - x'_j}{|\vec{x} - \vec{x}'|} \quad \vec{n} = \frac{\vec{x} - \vec{x}'}{|\vec{x} - \vec{x}'|}$$

$$\frac{\partial y_i}{\partial x_j} = \begin{pmatrix} 1 - \beta_1 n_1 & -\beta_1 n_2 & -\beta_1 n_3 \\ -\beta_2 n_1 & 1 - \beta_2 n_2 & -\beta_2 n_3 \\ -\beta_3 n_1 & -\beta_3 n_2 & 1 - \beta_3 n_3 \end{pmatrix}$$

$$\det(\vec{A}) = 1 - \beta_1 n_1 - \beta_2 n_2 - \beta_3 n_3 = 1 - \vec{\beta} \cdot \vec{n}$$

$$det(\mathcal{T}) = \frac{1}{1 - \vec{\beta} \cdot \vec{n}}$$

$$\therefore \phi(\vec{x}, t) = \left[\frac{q}{|\vec{x} - \vec{x}_0|} \frac{1}{1 - \beta \cdot \hat{n}} \right]_{\text{ret}}$$

Liénard-Wiechert potentials

$$\vec{n} = \frac{\vec{x} - \vec{x}_0}{|\vec{x} - \vec{x}_0|}$$

$$t' = t - \frac{|\vec{x} - \vec{x}_0(t)|}{c}$$

defined implicitly

$$\vec{A} = \left[\frac{e\vec{\beta}}{(\vec{x} - \vec{x}_0)(1 - \vec{\beta} \cdot \vec{n})} \right]_{ret}$$

$$\vec{E} = -\nabla\phi - \frac{1}{c}\frac{\partial \vec{A}}{\partial t} \quad \vec{B} = \nabla \times \vec{A}$$

$$\vec{A} = \frac{1}{c} \left\{ \frac{\vec{J}(x, t')}{|x - x'|} d^3 x' \quad t' = t - \frac{|x - x'|}{c} \quad \vec{J} = e^{-i\vec{k} \cdot \vec{x}} J(x - x'(t')) \right.$$

$$\frac{\partial x_0(t')}{\partial x_1} = \frac{dx_0}{dt'} \frac{\partial t'}{\partial x}$$

$$\text{Complicated} \quad \text{Ex. } \frac{\partial \Phi}{\partial x_1} \sim \frac{\partial}{\partial x_1} \frac{1}{(\bar{x} - \bar{x}_0)} = -\frac{1}{2} \frac{\partial}{\partial x_1} \frac{(x_1 - \bar{x}_0)^2 + \dots}{(\bar{x} - \bar{x}_0)^2}$$

$$\vec{E} = \frac{q}{c} \left[\frac{\vec{n} - \vec{B}}{\gamma^2 (1 - \vec{B} \cdot \vec{n})^2 R^2} \right]_{\text{ret}} + \frac{q}{c} \left[\frac{\vec{n} \times \{(\vec{n} - \vec{B}) \times \vec{B}\}}{(1 - \vec{B} \cdot \vec{n})^3 R} \right]_{\text{ret}}$$

$$\vec{B} = [\vec{n} \times \vec{E}]_{\text{ret}}$$

static field

radiation field (\hat{a})

$$\vec{\beta} \parallel \hat{x}$$

$$(R(1-\beta \cdot \vec{n}))^2 = (R - R\beta \cos\theta)^2 = r^2 - a^2 \quad r^2 = b^2 + (vt)^2$$

$$a = AB \sin\theta = Ab$$

$$r^2 - a^2 = b^2 + (vt)^2 - \beta^2 b^2 = \frac{b^2}{x^2} + (vt)^2$$

$$\beta_2 = 0 \quad n_2 = \sin\theta$$

$$E_2 = \frac{qb}{\gamma^2 \left(\frac{b^2}{\gamma^2} + (vt)^2 \right)^{3/2}} = \frac{qb\gamma}{\left(b^2 + (\gamma vt)^2 \right)^{3/2}} \quad \text{as before}$$

Radiation by Charges

2/20/2007

radiation part:

$$\vec{E} = \frac{e}{c} \left[\frac{\vec{n} \times (\vec{n} \times \vec{B}) \times \vec{i}}{(1 - \vec{B} \cdot \vec{n})^3 R} \right]_{\text{ret}}$$

$$\vec{B} = [\vec{n} \times \vec{E}]_{\text{ret}}$$

$$t' = t - \frac{R(t)}{c}$$

retarded time

Power Radiated

non relativistic $\beta \ll 1$

$$\vec{E} \approx \frac{e}{c} \left[\frac{\vec{n} \times (\vec{n} \times \vec{B})}{R} \right]_{\text{ret}}$$

$$\vec{S} = \frac{c}{4\pi} (\vec{E} \times \vec{B})$$

$$\vec{S} \cdot \vec{n} = \frac{c}{4\pi} |\vec{E}|^2 = \frac{c}{4\pi} \frac{e^2}{c^2} \frac{1}{R^2} |\vec{n} \times (\vec{n} \times \vec{B})|^2$$

$$dP = \vec{S} \cdot \vec{n} R^2 d\Omega$$

$$\frac{dP}{d\Omega} = \frac{e^2}{4\pi c} |\vec{n} \times (\vec{n} \times \vec{B})|^2$$



$$|\vec{n} \times (\vec{n} \times \vec{B})| = \sin^2 \theta \cdot \vec{B}^2 = \frac{v^2}{c^2} \sin^2 \theta$$

$$\frac{dP}{d\Omega} = \frac{e^2}{4\pi c^3} |v|^2 \sin^2 \theta$$

$$\int \sin^2 \theta d\Omega = \frac{8\pi}{3}$$

$$\therefore P_{\text{tot}} = \frac{2}{3} \frac{e^2}{c^3} |v|^2$$

Larmor result

 β not small:

$$P_{\text{tot}} = -\frac{2}{3} \frac{e^2}{m^2 c^3} \left(\frac{dp_u}{dz} \frac{dp^u}{dz} \right)$$

 z - proper time $dz = \frac{dt}{\gamma}$

$$p^u = \left(\frac{E}{c} \right) \vec{p}$$

$$\frac{dp_u}{dz} \frac{dp^u}{dz} = \frac{1}{c^2} \left(\frac{dE}{dz} \right)^2 - \left| \frac{d\vec{p}}{dz} \right|^2$$

$$\frac{E^2}{c^2} - |\vec{p}|^2 = m^2 c^2$$

$$\Rightarrow \left(\frac{d\vec{p}}{dz} \right)^2 - \left| \frac{d\vec{p}}{dz} \right|^2 = \frac{dp_u}{dz} \frac{dp^u}{dz}$$

$$\frac{2}{c^2} E \frac{dE}{dz} - 2 \vec{p} \cdot \frac{d\vec{p}}{dz} = 0 \quad \frac{dE}{dz} = c^2 \frac{\vec{p}}{E} \cdot \frac{d\vec{p}}{dz}$$

$$\vec{p} = \gamma m \vec{v} \quad E = \gamma m c^2 \quad \frac{dE}{dz} = \vec{v} \cdot \frac{d\vec{p}}{dz}$$

$$\text{for } \beta \gg 1 \rightarrow -\left| \frac{d\vec{p}}{dz} \right|^2 = -\left| \frac{d\vec{p}}{dt} \right|^2 = -m^2 |v|^2 \rightarrow \text{get Larmor result}$$

Total power radiated by relativistic particles:

$$\text{a) } \vec{B} \parallel \vec{B} \quad (\text{linear accelerator}): \quad \frac{dp_u}{dz} \frac{dp^u}{dz} = (\beta^2 - 1) \left| \frac{d\vec{p}}{dz} \right|^2$$

$$P_{\text{tot}} = \frac{2}{3} \frac{e^2}{m^2 c^3} \frac{1}{\gamma^2} \left| \frac{d\vec{p}}{dz} \right|^2 = \frac{2}{3} \frac{e^2}{m^2 c^3} \left| \frac{dp}{dt} \right|^2$$

$$\text{in terms of energy gain per length: } \frac{dE}{dx}$$

$$\frac{E^2}{c^2} - p^2 = mc^2 \quad 2 \frac{1}{c^2} E \frac{dE}{dt} - p^2 \frac{dp}{dt} = 0$$

$$\frac{dE}{dt} = \frac{dx}{dt} \frac{dE}{dx} = v \frac{dE}{dx} \quad v \frac{dE}{dx} = c^2 v \frac{dp}{dt} \quad \therefore \frac{dE}{dx} = \frac{dp}{dt}$$

$$\therefore P_{\text{tot}} = \frac{2}{3} \frac{e^2}{m^2 c^3} \left(\frac{dE}{dx} \right)^2$$

$$\text{power supplied: } \frac{dE}{dt} = v \frac{dE}{dx}$$

ratio of radiated power to supplied:

$$\frac{P}{dE/dt} = \frac{2}{3} \frac{e^2}{m^2 c^3} \frac{1}{v} \frac{dE}{dx}$$

(power lost)

$$\frac{P}{dE/dt} = \frac{2}{3} \frac{e^2/mc^2}{mc^2} \frac{dE}{dx} \ll 1 \rightarrow \text{in practice } \frac{dE}{dx} \ll \frac{mc^2}{e^2/mc^2} \sim \frac{0.5 \text{ MeV}}{10^{13} \text{ cm}} \sim 0.5 \times 10^{13} \text{ MeV/cm}$$

∴ radiation losses are negligible in linear accelerators

 $\frac{e^2}{mc^2}$ - "classical radius of e^- "

b) $\hat{\beta} \perp \vec{B}$ (circular accelerator): $\frac{dp_x dp^u}{d\tau d\tau} = - \left| \frac{d\vec{p}}{d\tau} \right|^2$

$$P_{tot} = \frac{2}{3} \frac{e^2}{m^2 c^3} \left| \frac{d\vec{p}}{d\tau} \right|^2 \quad \text{in lab} \quad \frac{d\vec{p}}{dt} = \vec{\omega} \times \vec{p} = \frac{1}{\gamma} \frac{d\vec{p}}{d\tau} \quad \therefore \left| \frac{d\vec{p}}{d\tau} \right| = \gamma \omega p$$

$$\therefore P_{tot} = \frac{2}{3} \frac{e^2}{m^2 c^3} \gamma^2 \omega^2 p^2$$



$$\text{energy loss per turn: } P_{tot} \frac{2\pi}{\omega} = SE = \frac{4\pi}{3} \frac{e^2}{m^2 c^3} \omega p^2 \gamma^2$$

$$wR = v \approx c \quad \omega = \frac{c}{R}$$

$$p = \gamma mv \approx \gamma mc \quad E = \gamma mc^2 \quad \gamma = \frac{E}{mc^2}$$

$$p^2 = \gamma^2 m^2 c^2 = \frac{E^2}{c^2}$$

$$\therefore SE = \frac{4\pi}{3} \frac{e^2}{m^2 c^3} \frac{c}{R} \frac{E^2}{c^2} \gamma^2 = \frac{4\pi}{3} \frac{e^2}{m^2 c^3} \frac{c}{R} \frac{E^4}{c^2 m^2 c^4}$$

$$SE = \frac{4\pi}{3} \frac{e^2}{R} \frac{E^4}{(mc^2)^4}$$

1 GeV e^- in $R=100\text{m}$: $SE \sim 10\text{ eV}$ can get larger with $E\gamma$ or $R\downarrow$

Angular Distribution of Radiation by Fast Particle

$$\vec{S} \cdot \vec{n}(t) = \frac{c}{4\pi} \frac{e^2}{c^2} \frac{1}{R^2} \left[\frac{\vec{n} \times ((\vec{n} - \vec{\beta}) \times \vec{\beta})}{(1 - \vec{\beta} \cdot \vec{n})^3} \right]_{ret}^2$$

"instantaneous" power $\frac{d\vec{p}}{d\Omega} = \frac{e^2}{4\pi c} []^2_{ret}$
 (not quite: $\int d\Omega \frac{d\vec{p}}{d\Omega} \neq P_{tot}$)

actual instantaneous power: $\frac{dP}{d\Omega} = \frac{d\tilde{p}}{d\Omega} \frac{dt}{dt'} \quad t' = t - \frac{R(t)}{c}$

$$t = t' + \frac{R(t)}{c} \quad \frac{dt}{dt'} = 1 + \frac{1}{c} \frac{dR(t)}{dt'}$$

$$R(t') = |\vec{x} - \vec{x}(t')| = \sqrt{(x_1 - x_{10})^2 + \dots}$$

$$\frac{dR(t')}{dt'} = \frac{1}{|\vec{x} - \vec{x}_0|} (-1) \frac{d\vec{x}_0}{dt'} \cdot (\vec{x} - \vec{x}_0) = -\vec{v} \cdot \vec{n}$$

$$\therefore \frac{dt}{dt'} = 1 - \vec{\beta} \cdot \vec{n}(t')$$

$$\therefore \frac{dP}{d\Omega} = \frac{e^2}{4\pi c} \frac{|\vec{n} \times ((\vec{n} - \vec{\beta}) \times \vec{\beta})|^2}{(1 - \vec{\beta} \cdot \vec{n})^5}$$

Check: $\int \frac{dP}{d\Omega} d\Omega = P_{tot}$ special case: $\hat{\beta} \parallel \vec{B}$ $\vec{\beta} \cdot \vec{n} = 0$

$$\frac{dP}{d\Omega} = \frac{e^2}{4\pi c} |\vec{\beta}|^2 \frac{\sin^2 \theta}{(1 - \beta \cos \theta)^5}$$

$$2\pi \int_{-1}^1 d(\cos \theta) \frac{\sin^2 \theta}{(1 - \beta \cos \theta)^5} = 2\pi \frac{4}{3} \frac{1}{(1 - \beta^2)^3} \quad \therefore P = \frac{2}{3} \frac{e^2}{c^3} \frac{|\vec{\beta}|^2}{(1 - \beta^2)^3} = \frac{2}{3} \frac{e^2}{c^3} \beta^6 v^2$$

$$\frac{dP}{dt} = \gamma m v + mv \frac{d\vec{v}}{dt} = \gamma m c \vec{\beta} + \gamma^3 m c \beta^2 \vec{\beta} = \gamma m c \vec{\beta} (1 + \beta^2 \gamma^2) = \gamma^3 m c \vec{\beta} \quad \therefore P = \frac{2}{3} \frac{e^2}{c^3} \gamma^6 v^2$$

$$1 + \frac{\beta^2}{1 - \beta^2} = \frac{1 - \beta^2 + \beta^2}{1 - \beta^2} = \frac{1}{1 - \beta^2} = \gamma^2$$

$$\vec{\beta} \cdot \vec{n} \sim \frac{1}{\gamma}$$

2/22/2007

$$\vec{E} = \frac{e}{c} \left(\hat{n} \times (\hat{n} - \vec{\beta}) \times \dot{\vec{B}} \right)_{\text{ret}}$$

$$\vec{B} = \hat{n} \times \vec{E}$$

$$T_2 + \frac{R(T_2)}{c}$$

Ayşe
Eşrefendil
Elenion
Ancalima

Energy radiated between $t' = T_1, T_2$

$$E = \int_{T_1}^{T_2 + \frac{R(T_2)}{c}} \frac{d\tilde{P}(t)}{dt} dt$$

$$\frac{d\tilde{P}}{dt} = \frac{e^2}{4\pi c} [\gamma]^2$$

$$dt = \frac{dt'}{dt}, dt' = (1 - \vec{\beta} \cdot \hat{n}) dt$$

$$t = t' + \frac{R(t')}{c}$$


$$E = \int_{T_1}^{T_2} \frac{e^2}{4\pi c} [\gamma]^2 dt' (1 - \vec{\beta} \cdot \hat{n})$$

$$\frac{dP}{d\Omega} = (1 - \vec{\beta} \cdot \hat{n}) \frac{d\tilde{P}}{d\Omega}$$

$$\frac{dP}{d\Omega} = \frac{e^2}{4\pi c} \frac{|\hat{n} \times ((\hat{n} - \vec{\beta}) \times \dot{\vec{B}})|^2}{(1 - \vec{\beta} \cdot \hat{n})^5}$$

Pattern of Radiation

Simple Cases



$$\frac{dP}{d\Omega} = \frac{e^2}{4\pi c^3} |\dot{\vec{v}}|^2 \frac{\sin^2 \theta}{(1 - \beta \cos \theta)^5}$$

$$\max \text{ of } \frac{dP}{d\Omega}: \frac{d}{d\theta} \frac{\sin^2 \theta}{(1 - \beta \cos \theta)^5} = 0$$

$$\rightarrow \cos \theta_m = \frac{1}{3\beta} (-1 + \sqrt{1 + 15\beta^2}) \quad \beta \approx 1 \quad \beta = 1 - \epsilon \quad \beta^2 \approx 1 - 2\epsilon = 1 - 2(1 - \epsilon)$$

$$1 - \beta \approx \frac{1}{2}(1 - \beta^2) = \frac{1}{2\gamma^2}$$

$$\cos \theta_m = 1 - \frac{\theta^2}{2} = \frac{1}{3(1-\epsilon)} \left[-1 + \sqrt{1 + 15 - 30\epsilon} \right]$$

$$1 - \frac{\theta^2}{2} \approx \frac{1}{3}(1 + \epsilon)(3 - \frac{15}{4}\epsilon) = (1 + \epsilon)(1 - \frac{5}{4}\epsilon) \approx 1 - \frac{1}{4}\epsilon$$

$$4\sqrt{1 - \frac{15}{8}\epsilon} \approx 4(1 - \frac{15}{16}\epsilon)$$

$$\frac{\theta^2}{2} \approx \frac{1}{4}\epsilon \quad \theta^2 \approx \frac{\epsilon}{2} = \frac{1 - \beta}{2} = \frac{1}{4\gamma^2}$$

$$\theta_m \approx \frac{1}{2\gamma}$$



10 GeV e⁻ $\gamma = 10^4$ almost parallel beam with high intensity

$$\text{peak power } \frac{dP}{d\Omega}|_{\theta_m} \sim \left(\frac{1}{2\gamma}\right)^2 \left(\frac{8}{5}\right)^5 \delta^{10} \frac{e^2}{4\pi c^3} |\dot{\vec{v}}|^2 \sim \frac{e^2}{4\pi c^3} |\dot{\vec{v}}|^2 \gamma^8 \propto \gamma^8$$

$$1 - \beta \left(1 - \frac{\theta^2}{2}\right) = 1 - \beta + \frac{1}{2}\left(\frac{1}{2\gamma}\right)^2 \approx \frac{1}{2\gamma^2} + \frac{1}{8\gamma^2} = \frac{5}{8} \frac{1}{\gamma^2}$$


$$P \sim \left(\frac{dP}{d\Omega}\right) \Delta\Omega \quad \Delta\Omega \propto \theta_m^2 \quad \therefore P \sim \gamma^8 \frac{1}{\gamma^2} \sim \gamma^6$$

$$\vec{\beta} \perp \vec{B}: \quad \vec{\beta} = (0, 0, \beta) \quad \dot{\vec{B}} = (\dot{\beta}, 0, 0) \quad \hat{n} = (\sin \theta \cos \phi, \sin \theta \sin \phi, \cos \theta)$$

$$|\dot{\vec{v}}|^2 = \beta^2 \left\{ (1 - \beta \cos \theta)^2 - (1 - \beta^2) \sin^2 \theta \cos^2 \phi \right\}$$

$$\frac{dP}{d\Omega} = \frac{e^2}{4\pi c^3} |\dot{\vec{v}}|^2 \frac{1}{(1 - \beta \cos \theta)^5} \left(1 - \frac{\sin^2 \theta \cos^2 \phi}{\theta^2 (1 - \beta \cos \theta)^2} \right)$$

$$\theta_m = 0 \quad \text{peak power } \frac{dP}{d\Omega}|_{\theta_m} = \frac{e^2 |\dot{\vec{v}}|^2}{4\pi c^3} 8\gamma^6 \propto \gamma^6$$



$$\text{peak } \frac{1}{(1 - \beta \cos \theta)^3} \approx \frac{1}{(1 - \beta + \beta \frac{\theta^2}{2})} = \frac{1}{\frac{1}{2}\gamma^2 + \frac{\theta^2}{2}} = \frac{2\gamma^2}{1 + \theta^2 \gamma^2} \quad \text{opening angle } \frac{\theta^2}{2} \sim \frac{1}{2\gamma^2} \quad \theta \sim \frac{1}{\gamma}$$

$$\text{total power at peak } P \sim \left(\frac{dP}{d\Omega}\right) \Delta\Omega \sim \gamma^6 \frac{1}{\gamma^2} \sim \gamma^4$$

$$\vec{\beta} \perp \vec{B}: P = \frac{2}{3} \frac{e^2}{c^3} |\dot{\vec{v}}|^2 \gamma^4 \quad \vec{\beta} \parallel \vec{B}: P = \frac{2}{3} \frac{e^2}{c^3} |\dot{\vec{v}}|^2 \gamma^6$$


in terms of applied force

$$\text{circular motion } \vec{F} = \frac{d\vec{p}}{dt} = \frac{d}{dt}(\gamma m \vec{v}) = \gamma m \frac{d\vec{v}}{dt} \quad |\dot{\vec{v}}|^2 = \frac{F^2}{\gamma^2 m^2}$$

$$\begin{aligned} \text{linear motion } \vec{F} &= \frac{d}{dt}(\gamma m \vec{v}) = \gamma m \frac{dv}{dt} + mv \frac{d\vec{v}}{dt} \quad \frac{d\vec{v}}{dt} = \gamma^2 \frac{1}{c} \vec{v} \quad \therefore |\dot{\vec{v}}|^2 = \frac{F^2}{\gamma^2 m^2} \\ &= \gamma m \dot{v} + m v^2 \vec{v} \cdot \vec{v} = \gamma m \dot{v} (1 + \gamma^2 v^2) = \gamma^3 m \dot{v} \\ &\quad 1 + \frac{1}{1-\beta^2} = \frac{1-\beta^2+\beta^2}{1-\beta^2} = \frac{1}{1-\beta^2} = \gamma^2 \end{aligned}$$

$$\therefore P_{\perp} = \frac{2}{3} \frac{e^2}{c^3} \frac{F^2}{m^2} \gamma^2 \quad P_{\parallel} = \frac{2}{3} \frac{e^2}{c^3} \frac{F^2}{m^2}$$

\rightarrow power radiated comes from \vec{P}_{\perp} ($\gamma \gg 1$)




$$v_{\perp} = \frac{v^2}{R} \approx \frac{c^2}{R}$$

2/27/2007

HW 6: 14. 4, 5, 7ab, 12

Power radiated in large impact parameter collision



$$\text{a) Non-relativistic } \beta \ll 1 \quad \text{Larmor formula: } P = \frac{2}{3} \frac{e^2}{c^3} |\dot{\vec{v}}|^2$$

$$\vec{F} = \frac{qQ}{r^2} \quad r^2 = (vt)^2 + b^2 \quad \vec{F} = m\dot{\vec{v}} \quad \dot{v} = \frac{F}{m} = \frac{qQ}{mr^2} = \frac{qQ}{m((vt)^2 + b^2)}$$

$$E = \int_{-\infty}^{\infty} P dt = \frac{2}{3} \frac{Q^2 e^4}{m^2 c^3} \int_{-\infty}^{\infty} \frac{dt}{((vt)^2 + b^2)^2}$$

$$\text{Integral} = \frac{1}{b^4} \frac{b}{v} \int_{-\infty}^{\infty} \frac{d(vt)}{(1 + (\frac{vt}{b})^2)^2} = \frac{1}{vb^3} \int_{-\infty}^{\infty} \frac{dx}{(1+x^2)^2} = \frac{\alpha}{vb^3}$$

$$E = \alpha \frac{2}{3} \frac{Q^2 e^4}{m^2 c^3 b^3 v}$$

$$\int_{\Gamma} \frac{dz}{1+z^2}$$

$$\int_{\Gamma} \frac{dz}{z-i}$$

$$f(z) \downarrow$$

important part of a number: $\alpha \omega \nu$

$$\int_{\Gamma} \frac{dz}{1+z^2} = 2\pi i \frac{1}{2i} = \pi$$

2*pi*i (Residue at pole)

$$f(\text{pole}) = -\frac{1}{2i}$$



small deflection

$$P_{\parallel} \gg P_{\perp} \quad \frac{P_{\perp}}{P_{\parallel}} \ll 1$$

$$\sin \theta = \frac{b}{r}$$

b) Ultrarelativistic $\gamma \gg 1$ use \dot{v}_\perp $P = \frac{2}{3} \frac{e^2}{c^3} \dot{v}_\perp^2 \gamma^4$ for $\vec{\dot{B}} \perp \vec{B}$

$$\frac{d\vec{p}}{dt} = \frac{qQ}{r^2} \quad \vec{p} = qm\vec{v}$$

$$\frac{d}{dt}(qmv) = qm\dot{v}_\perp$$

considering only perpendicular motion

$$\dot{v}_\perp^2 = \left(\frac{qQ}{r^2} \frac{b}{r} \right)^2 \frac{1}{8^2 m^2} \quad \text{at } r = b \quad \sin\theta = \frac{b}{r}$$

$$\dot{v}_\perp^2 \propto \frac{1}{r^6} \quad r^2 = (vt)^2 + b^2$$

$$E = \frac{2}{3} \frac{e^4 Q^2}{m^2 c^4} \frac{\gamma^2}{b^3} A$$

$$E = \frac{2}{3} \frac{q^2}{c^3} \gamma^4 \frac{q^2 Q^2}{8^2 m^2} b^2 \int_{-\infty}^{\infty} \frac{dt}{((vt)^2 + b^2)^3}$$

$$\frac{1}{b^6} \frac{b}{v} \int_{-\infty}^{\infty} \frac{dx}{(x^2 + 1)^3} \quad v = c \quad A = \frac{3\pi}{8}$$

Contribution from $\vec{\dot{B}} \parallel \vec{B}$

$$P = \frac{2}{3} \frac{q^2}{c^3} \dot{v}_\parallel^2 \gamma^2 \gamma^6$$

$$\frac{dp_\parallel}{dt} = qm\dot{v}_\parallel + mv\dot{v}$$

$$P = \frac{2}{3} \frac{q^2}{c^3} \frac{F_\parallel^2}{m^2} \quad \text{small in ultrarelativistic limit} \quad F_\parallel = \frac{qQ}{r^2} \frac{vt}{r}$$

Frequency Components



pulse width Δt $\Delta E \Delta t \sim h$ $\Delta E = h \Delta \nu$ $\Delta \nu \sim \frac{1}{\Delta t}$

light contains frequencies at least up to $\frac{1}{\Delta t}$

one would say (wrong): $R \frac{1}{\delta} = c \Delta t \quad \Delta t = \frac{R}{c \delta}$ wrong

$$\begin{aligned} & \text{Diagram showing light travel from source at } (0,0) \text{ to observer at } (t_2, R) \\ & t_1 = 0 \quad t_2 = \frac{R}{c \delta} \quad \tilde{t}_1 = 0 + \frac{r}{c} = \frac{r}{c} \quad \text{light gets to observer} \\ & \text{if } \delta = \frac{R}{\delta} \quad \Delta \tilde{t} = \tilde{t}_2 - \tilde{t}_1 = \frac{R}{c \delta} + \frac{r}{c} - \frac{R}{c \delta} + \frac{R}{2 \delta^2 c} - \frac{r}{c} = 0 \quad \text{wrong} \end{aligned}$$

$$\delta = \frac{R}{\delta} \cos(\frac{1}{\delta}) \approx \frac{R}{\delta} \left(1 - \frac{1}{2 \delta^2} \right) \quad \Delta \tilde{t} = \tilde{t}_2 - \tilde{t}_1 = \frac{R}{c \delta} + \frac{r}{c} - \frac{R}{c \delta} + \frac{R}{2 \delta^2 c} - \frac{r}{c} = \frac{R}{2 \delta^2 c}$$

$$\Delta t \sim \frac{R}{\delta^2 c} \quad \Delta \nu \sim \frac{c}{R} \delta^3 \quad \text{high freq. can get x-rays from synchrotron radiation}$$

Thomson Scattering

3/1/2007

EM wave scattering off a free charge

nonrelativistic: ok if $\frac{hv}{c} \ll mc$ ($hv \ll mc^2$)

$$\frac{dP}{d\Omega} = \frac{e^2}{4\pi c^3} \left| \hat{n} \times (\hat{n} \times \vec{v}) \right|^2 \quad \text{incoming wave: } \vec{E} = \vec{E}_{in} E_0 e^{i(\vec{k}_0 \cdot \vec{x} - \omega t)}$$

$$\vec{v} = \frac{e \vec{E}}{m} = \frac{e}{m} E_0 \vec{E}_{in} e^{i(\vec{k}_0 \cdot \vec{x} - \omega t)} \quad \text{if amplitude of motion of charge is } \ll \lambda : \vec{v} \approx \frac{e E_0}{m} \vec{E}_{in} e^{i\omega t} e^{i\text{const}}$$

Power radiated (in \hat{n}) with polarization \vec{E}_{out}

$$\frac{dP}{d\Omega}(\vec{E}_{out}) = \frac{e^2}{4\pi c^3} \left| \vec{E}_{out} \cdot [\hat{n} \times (\hat{n} \times \vec{v})] \right|^2 \quad \vec{E}_{out} \cdot \{ (\hat{n} \cdot \vec{v}) \hat{n} - \vec{v} \} = \vec{E}_{out} \cdot \vec{v}$$

$$\frac{dP}{d\Omega} = \frac{e^2}{4\pi c^3} \left| \vec{E}_{out} \cdot \vec{v} \right|^2$$

$$\frac{dP}{d\Omega}(\vec{E}_{in}, \vec{E}_{out}) = \frac{e^4 E^2}{4\pi m^2 c^3} \frac{1}{2} \left| \vec{E}_{out} \cdot \vec{E}_{in}^* \right|^2$$

time averaged $\rightarrow \frac{1}{2} |\vec{A} \cdot \vec{B}|^2$

$$\frac{d\sigma}{d\Omega} = \frac{dP}{d\Omega} \frac{8\pi}{cE^2}$$

\downarrow (Flux in)⁻¹ flux in = $S = \frac{c}{8\pi} E^2$

$$\frac{e^2}{mc^2} = \text{"classical e- radius"}$$

average over initial polarizations (\vec{E}_{in}) then sum over \vec{E}_{out}

a) $\vec{E}_{out} \perp$ scattering plane

$$\vec{E}_{out} \cdot \vec{E}_{in} = \cos \varphi$$

$$\langle |\vec{E}_{out} \cdot \vec{E}_{in}| \rangle = \frac{1}{2}$$

"top down" (k_out of page)

$$\langle \cos^2 \varphi \rangle = \frac{1}{2\pi} \int_0^{2\pi} \cos^2 \varphi d\varphi$$

b) $\vec{E}_{out} \parallel$ scattering plane

$$\vec{E}_{out} \cdot \vec{E}_{in} = \cos \theta$$

$$\vec{E}_{out} \cos \theta \cdot \vec{E}_{in} = \cos \theta \cos \varphi$$

$$\langle \cos^2 \theta \cos^2 \varphi \rangle = \cos^2 \theta \langle \cos^2 \varphi \rangle = \frac{1}{2} \cos^2 \theta$$

$$\frac{d\sigma_I}{d\Omega} = \left(\frac{e^2}{mc^2} \right)^2 \frac{1}{2} \quad \frac{d\sigma_{II}}{d\Omega} = \left(\frac{e^2}{mc^2} \right)^2 \frac{1}{2} \cos^2 \theta \quad \frac{d\sigma}{d\Omega} = \left(\frac{e^2}{mc^2} \right)^2 \frac{1}{2} (1 + \cos^2 \theta)$$

If relativistic \rightarrow get Compton scattering

$$\begin{array}{ccc} \left(\frac{h\nu}{c}, h\vec{k}_{in} \right) & \left(mc, 0 \right) & \left(\frac{h\nu'}{c}, h\vec{k}_{out} \right) \\ \xrightarrow{\text{e-}} & \downarrow & \xrightarrow{\text{e-} (\gamma mc, \gamma m\vec{v})} \end{array}$$

Motion of Particles in EM Fields (Chpt 12)

$$\frac{d\vec{v}}{dt} = e(\vec{E} + \frac{\vec{v}}{c} \times \vec{B}) \quad \vec{p} = \gamma m \vec{v} \quad \frac{d\vec{E}}{dt} = e \vec{E} \cdot \vec{v} \quad P = \vec{F} \cdot \vec{v}$$

$$\frac{d\vec{x}}{dt} = \gamma^3 \vec{B} \cdot \dot{\vec{B}}$$

$$\frac{dE}{dt} = mc^2 \frac{d\gamma}{dt} = \gamma^3 mc^2 \vec{B} \cdot \dot{\vec{B}} \quad \vec{v} \cdot \frac{d\vec{p}}{dt} = e \vec{E} \cdot \vec{v} \quad \frac{d\vec{p}}{dt} = \gamma m \frac{d\vec{v}}{dt} + m\vec{v} \frac{d\vec{x}}{dt} = \gamma m c \dot{\vec{B}} + mc \vec{B} (\gamma^3 \vec{B} \cdot \dot{\vec{B}})$$

$$\vec{v} \cdot \frac{d\vec{p}}{dt} = mc^2 \vec{B}^2 \gamma^3 \vec{B} \cdot \dot{\vec{B}} + \gamma m c^2 \vec{B} \cdot \dot{\vec{B}} = \gamma m c^2 \vec{B} \cdot \dot{\vec{B}} (\underbrace{\vec{B}^2 \gamma^2 + 1}_{\gamma^2}) = \gamma^3 m c^2 \vec{B} \cdot \dot{\vec{B}} = \frac{dE}{dt}$$

$$\therefore \frac{dE}{dt} = \vec{v} \cdot \frac{d\vec{p}}{dt} \quad \frac{\vec{B}^2}{1-\beta^2} + 1 = \frac{1}{1-\beta^2} = \gamma^2$$

1. Uniform, constant \vec{B}

$$\frac{d\vec{E}}{dt} = 0 \quad \gamma = \text{const.} \quad \gamma m \frac{d\vec{v}}{dt} = \frac{e}{c} \vec{v} \times \vec{B} \quad \frac{d\vec{v}}{dt} = \vec{v} \times \frac{e\vec{B}}{\gamma mc} = \vec{v} \times \vec{\omega} \quad \vec{\omega} = \frac{e\vec{B}}{\gamma mc}$$

trajectory is helix along \vec{B}


circular motion

radius of circle

$$r_\perp = \omega R \quad \frac{r_\perp}{\gamma m} = \frac{eB}{\gamma mc} R \quad R = \frac{cp_\perp}{eB}$$

2. Crossed \vec{E} & \vec{B} (uniform, constant) $\vec{E} \perp \vec{B}$

$$K: \vec{E} = E\hat{j}, \vec{B} = B\hat{z}$$



Lorentz transf. $\vec{B}' = \beta \hat{x}$

$$1) |\vec{B}| > |\vec{E}| \quad \beta = \frac{E}{B}$$

$$K': E'_1 = E_1, E'_2 = \gamma(E_2 - \beta B_3), E'_3 = \gamma(E_3 + \beta B_2)$$

$$E'_1 = 0, E'_2 = \gamma(E - \beta B) = \gamma(E - \frac{E}{B}B) = 0, E'_3 = 0$$

$$B'_1 = 0, B'_2 = 0, B'_3 = \gamma(B_3 - \beta E_2) = \gamma(B - \frac{E}{B}E) = \gamma B(1 - \frac{E^2}{B^2}) = \gamma B(1 - \beta^2) = \frac{1}{\gamma} B$$

$$\vec{B}' = \frac{\vec{E} \times \vec{B}}{B^2} \quad \text{for } |\vec{B}| > |\vec{E}| \rightarrow \text{eliminates } E \text{ field}$$

in K' trajectory is helix along \vec{B}'

in K trajectory: circle around B lines + drift \vec{B}



$$2) |\vec{E}| > |\vec{B}| \quad \beta = \frac{B}{E} \dots \vec{B} = 0, \vec{E} = \frac{1}{\gamma} E \hat{y}$$

Static, uniform \vec{E} $\vec{v}_0 \perp \vec{E}$

$$K: \vec{E} = E\hat{j}, \vec{v}_0 = v_0\hat{x}$$

$$\frac{d\vec{p}}{dt} = q\vec{E}$$

$$\frac{d}{dt}(qmv_x) = 0 \quad \frac{d}{dt}(qmv_y) = qE$$

$$qmv_x = v_0 m v_0$$

$$qmv_y = qEt$$

calculate $\gamma(t)$

$$\frac{dE}{dt} = mc^2 \frac{d\gamma}{dt} = qE v_y$$

$$mc^2 \frac{d\gamma}{dt} = qE \frac{qEt}{qm}$$

$$\gamma \frac{d\gamma}{dt} = \frac{q^2 E^2}{m^2 c^2} t$$

$$\frac{1}{2}(\gamma^2 - \gamma_0^2) = \frac{q^2 E^2}{m^2 c^2} \frac{1}{2} t^2$$

$$\gamma(t) = \sqrt{\gamma_0^2 + \frac{q^2 E^2}{m^2 c^2} t^2}$$

$$v_x = \frac{v_0 v_0}{\gamma(t)} = v_0 \left(1 + \left(\frac{qE}{v_0 m}\right)^2 t^2\right)^{-1/2} \quad \text{characteristic time } \tau = \frac{v_0 m c}{qE}$$

$$v_x(t) = \frac{v_0}{\sqrt{1 + (\frac{t}{\tau})^2}}$$

$$v_y(t) = \frac{qE}{qm} \frac{t}{\sqrt{1 + (\frac{t}{\tau})^2}}$$

3/6/2007

As $t \rightarrow \infty$ $v_x \rightarrow 0$ $v_y \rightarrow c$

short times ($t \ll \tau$)

$$v_x \approx v_0 \left(1 - \frac{1}{2} \left(\frac{t}{\tau}\right)^2\right)$$

$$v_y \approx \frac{qE}{qm} t \left(1 - \frac{1}{2} \left(\frac{t}{\tau}\right)^2\right)$$

$$v_x \approx v_0$$

$$v_y \approx \frac{qE}{qm} t$$

$$\therefore x(t) \approx v_0 t \quad y(t) \approx \frac{qE}{2qm} t^2$$

long times ($t \gg \tau$)


$$v_x \approx \frac{v_0 \tau}{t}$$

$$v_y \approx \frac{qE}{qm} t \frac{\tau}{t} = c$$

$$x(t) \approx v_0 \tau \ln\left(\frac{t}{\tau}\right)$$

$$y(t) \approx ct \quad \text{parabola (as non-rel)}$$

$$\text{trajectory: } y = c\tau e^{\frac{x}{v_0 \tau}}$$



locally: B lines are parallel with curvature $\sim \frac{1}{R}$

cylindrical coord. at origin
(r, ϕ, z)

$$\vec{B} = B_\phi \hat{\phi}$$

$$\nabla \times \vec{B} = 0 \quad (\text{no currents}) \Rightarrow \frac{1}{r} \frac{\partial}{\partial r} (r B_\phi) = 0 \quad \therefore B_\phi = B_0 \frac{R}{r} \quad \nabla \cdot \vec{B} = 0$$

$\gamma = \text{const.}$ $\frac{d}{dt}(\gamma m c^2) = 0$ no electric fields

$$\frac{d\vec{p}}{dt} = \gamma m \frac{d\vec{v}}{dt} = \frac{e}{c} \vec{v} \times \vec{B} \quad \vec{v} = \rho \hat{r} + \rho \dot{\phi} \hat{\phi} + \dot{z} \hat{z} \quad \dot{\vec{v}} = (\ddot{\rho} - \rho \dot{\phi}^2) \hat{r} + (\rho \ddot{\phi} + 2\rho \dot{\phi}) \hat{\phi} + \dot{z} \hat{z}$$

$$\vec{v} \times \vec{B} = B_\phi \rho \dot{z} \hat{z} - B_\phi \dot{z} \hat{r}$$

$$\hat{r}: \quad \ddot{\rho} - \rho \dot{\phi}^2 = -\frac{e B_0}{\gamma m c} \frac{R}{\rho} \dot{z}$$

$$\omega_B = \frac{e B_0}{\gamma m c}$$

$$\hat{\phi}: \quad \rho \ddot{\phi} + 2\rho \dot{\phi} = 0 \quad \hat{z}: \quad \ddot{z} = \frac{e B_0}{\gamma m c} \frac{R}{\rho} \dot{\rho}$$

$$\frac{d}{dt}(\rho^2 \dot{\phi}) = 2\rho \dot{\rho} \dot{\phi} + \rho^2 \ddot{\phi} = \rho(\rho \ddot{\phi} + 2\rho \dot{\phi}) = 0 \quad \therefore \rho^2 \dot{\phi} = \text{constant} = R v_{||}$$

$$\frac{d}{dt}(\ln \rho) = \frac{\dot{\rho}}{\rho} \propto \hat{z} \quad \therefore \hat{z} = \omega_B R \ln\left(\frac{\rho}{R}\right) + v_0 \quad \text{--- integration constant}$$

$$\frac{d}{dt}(\ln\left(\frac{\rho}{R}\right)) = \frac{R}{\rho} \frac{d}{dt}\left(\frac{\rho}{R}\right) = \frac{\dot{\rho}}{\rho}$$

$$\text{ansatz: } \rho(t) = R + y(t) \quad y \ll R$$

→ small curvature

probable motion:

this should be y

$$\ln\left(\frac{\rho}{R}\right) = \ln\left(1 + \frac{y}{R}\right) \simeq \frac{y}{R} \quad \therefore \hat{z} \simeq \omega_B y + v_0$$

$$\hat{r}: \quad \ddot{y} - \frac{R^2 v_{||}^2}{\rho^3} = -\omega_B \frac{R}{\rho} (\omega_B y + v_0) \quad \frac{1}{\rho^3} = \frac{1}{(R+y)^3} = \frac{1}{R^3} \left(1 + \frac{y}{R}\right)^3 \simeq \frac{1}{R^3} \left(1 - 3 \frac{y}{R}\right) \quad \frac{1}{\rho} \simeq \frac{1}{R} \left(1 - \frac{y}{R}\right)$$

$$\ddot{y} - \frac{v_{||}^2}{R} \left(1 - 3 \frac{y}{R}\right) = -\omega_B \left(1 - \frac{y}{R}\right) (\omega_B y + v_0) \quad \ddot{y} + y \left(3 \frac{v_{||}^2}{R^2} - \omega_B \frac{v_0}{R} + \omega_B^2\right) = \frac{v_{||}^2}{R} - \omega_B v_0 \quad \text{oscillator}$$

Drop y^2 terms

$$\ddot{x} + \omega^2 x = 0 \quad x = A \sin(\omega t) + \frac{\dot{x}_0}{\omega^2} \quad \text{simple harmonic oscillator around displaced equilibrium}$$

$$\frac{\dot{x}_0}{\omega^2} = \langle y \rangle = \frac{\frac{v_{||}^2}{R} - \omega_B v_0}{3 \frac{v_{||}^2}{R^2} + \omega_B^2 - \omega_B \frac{v_0}{R}} \quad \therefore \langle \hat{z} \rangle = \omega_B \langle y \rangle + v_0 \quad \text{drift motion}$$

$$\text{say: } \omega_B R \gg v_{||}, v_0 \Rightarrow \langle y \rangle \simeq \frac{v_{||}^2}{\omega_B^2 R} - \frac{v_0}{\omega_B} \quad \langle \hat{z} \rangle \simeq \frac{v_{||}^2}{R \omega_B} \quad \text{curvature drift}$$

$$\text{If } v_{||} = 0, \quad \langle \hat{z} \rangle = \frac{v_0^2}{R \omega_B} \quad \text{still a drift} \quad \text{no drift if } v_{||} = v_0 = 0$$

Another case: \vec{B} has a gradient $\nabla_x \vec{B} \perp \vec{B}$

$$\vec{B} \parallel \hat{x} \quad \nabla \cdot \vec{B} \parallel \hat{x} \quad \gamma = \text{const.}$$

$$\gamma m \frac{d\vec{v}}{dt} = \frac{e}{c} \vec{v} \times \vec{B}$$

$$\vec{v} = \frac{e}{\gamma mc} \vec{v} \times \vec{B}$$

$$\vec{B} = \vec{B}_0 + \vec{B}_1(x) = \left(\vec{B}_0 + \frac{\partial \vec{B}}{\partial x} dx \right) \hat{x}$$

$$\dot{\vec{v}} = \vec{v} \times \vec{\omega}(x)$$

$$\vec{\omega} = \frac{e}{\gamma mc} (\vec{B}_0 + \vec{B}_1) = \vec{\omega}_0 + \vec{\omega}_1$$

uniform

first order
perturbation

location of particle

$$\vec{v} = \vec{v}_0(t) + \vec{v}_1(t)$$

unperturbed
(we know sol'n)

$$\vec{v}_1 = \vec{v}_0 \times \vec{\omega}_1 + \vec{v}_1 \times \vec{\omega}_0$$

$$\vec{\omega}_0 = \frac{e \vec{B}_0}{\gamma mc} \quad \vec{\omega}_1 = \frac{e}{\gamma mc} \frac{\partial \vec{B}}{\partial x} dx$$

look for a drift: $\langle \dot{\vec{v}}_1 \rangle = 0 \quad \langle \vec{v}_1 \rangle \neq 0 \quad \langle \vec{v}_0 \times \vec{\omega}_1 \rangle = -\langle \vec{v}_1 \times \vec{\omega}_0 \rangle = -\langle \vec{v}_1 \rangle \times \vec{\omega}_0$

Solution to unperturbed problem: $\vec{v}_0(t) = (a \cos(\omega_0 t), -a \sin(\omega_0 t), 0)$

$$\vec{v}_0 = (-a \omega_0 \sin(\omega_0 t), -a \omega_0 \cos(\omega_0 t), 0) \quad dx \rightarrow a \cos(\omega_0 t)$$

$$\vec{v}_0 \times \vec{\omega}_1 = -\frac{e}{\gamma mc} \frac{\partial \vec{B}}{\partial x} a \cos(\omega_0 t) (a \omega_0 \cos(\omega_0 t) \hat{x} - a \omega_0 \sin(\omega_0 t) \hat{y})$$

$$\langle \vec{v}_0 \times \vec{\omega}_1 \rangle = \hat{x} \frac{1}{2} \frac{e}{\gamma mc} \frac{\partial \vec{B}}{\partial x} a^2 \omega_0 = -\hat{x} \frac{1}{2} a^2 \omega_0^2 \frac{1}{B_0} \frac{\partial \vec{B}}{\partial x}$$

$$-\langle \vec{v}_1 \rangle \times \vec{\omega}_0 \Rightarrow \boxed{\langle \vec{v}_1 \rangle = \hat{y} \frac{1}{2} a^2 \omega_0 \frac{1}{B_0} \frac{\partial \vec{B}}{\partial x}}$$

3/8/2007

Lagrangian of Particles

Action: non rel.: $S = \int_{t_1}^{t_2} L(\dot{x}, x) dt$ $\vec{x}(t)$ trajectory

$$S = 0 \rightarrow \frac{d}{dt} \frac{\partial L}{\partial \dot{x}} - \frac{\partial L}{\partial x} = 0 \quad \text{equation of motion}$$

free particle: $\mathcal{L} = \frac{1}{2} m v^2$

free particle, relativistic

ask that S is a Lorentz scalar

$$dt = \gamma d\tau \quad \text{in frame of particle (rest frame)}$$

$$S = \int_{x_1}^{x_2} \gamma \mathcal{L} d\tau \quad \gamma \mathcal{L} \text{ is } \downarrow \text{L-scalar}$$

$$P^\mu = \left(\frac{E}{c}, \vec{p} \right)$$

$$P_\mu P^\mu = m^2 c^2$$

$$\gamma \mathcal{L} \propto m^2 c^2 \dots$$

$$\mathcal{L} \propto \frac{1}{8} = \sqrt{1 - \beta^2}$$

$$\text{want } \beta \rightarrow 0 \quad \mathcal{L} \rightarrow \frac{1}{2} m v^2$$

$$\sqrt{1 - \beta^2} \approx 1 - \frac{1}{2} \frac{v^2}{c^2}$$

$$\boxed{\mathcal{L} = -mc^2 \sqrt{1 - \beta^2}}$$

relativistic \mathcal{L}
for free particle

$$\frac{\partial \mathcal{L}}{\partial v} = \frac{\partial}{\partial v} \left(-mc^2 \sqrt{1 - \frac{v^2}{c^2}} \right) = -mc^2 \left(\frac{1}{2} \right) \left(-\frac{2v}{c^2} \right) \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} = \gamma m v$$

$$\frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial v} \right) = \frac{d}{dt} (\gamma m v) = 0 \quad \text{true}$$

interaction part of \mathcal{L} : \mathcal{L}_{int} in EM field

non-rel: $\mathcal{L}_{\text{int}} = -q\phi$ rel: $\delta\mathcal{L}_{\text{int}}$ is L-scalar

\mathcal{L}_{int} will depend on fields (linear), \vec{v} , and \vec{x} (through fields $\vec{A}(\vec{x})$, $\phi(\vec{x})$)

$$p^\mu = \left(\frac{E}{c}, \vec{p} \right) \quad A^\mu = (\phi, \vec{A}) \quad p_\mu A^\mu \Rightarrow \delta\mathcal{L}_{\text{int}} \propto p_\mu A^\mu = \gamma mc \phi - \vec{p} \cdot \vec{A}$$

$$\mathcal{L}_{\text{int}} \propto mc\phi - \frac{1}{8}\vec{p} \cdot \vec{A} \rightarrow -q\phi \text{ non-rel}$$

$$\mathcal{L}_{\text{int}} = -q\phi + \frac{q}{\gamma mc} \vec{p} \cdot \vec{A} = -q\phi + \frac{q}{c} \vec{v} \cdot \vec{A} \quad \mathcal{L}_{\text{int}} = -q\phi + q \frac{\vec{v}}{c} \cdot \vec{A} \quad \text{relativistic } \mathcal{L}_{\text{int}}$$

$$\boxed{\mathcal{L} = -mc^2 \sqrt{1-\beta^2} - q\phi + q \frac{\vec{v}}{c} \cdot \vec{A}} \quad \text{relativistic } \mathcal{L} \text{ with EM fields}$$

$$\frac{\partial \mathcal{L}}{\partial v_i} = \gamma mv_i + \frac{q}{c} A_i = p_i + \frac{q}{c} A_i \quad \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial v_i} \right) = \frac{dp_i}{dt} + \frac{q}{c} \frac{dA_i}{dt}$$

$$\frac{\partial \mathcal{L}}{\partial x_i} = -q \frac{\partial \phi}{\partial x_i} + \frac{q}{c} v_j \frac{\partial A_j}{\partial x_i} \quad \frac{dp_i}{dt} = -\frac{q}{c} \frac{dA_i}{dt} + \frac{q}{c} v_j \frac{\partial A_j}{\partial x_i} - q \frac{\partial \phi}{\partial x_i}$$

$$\vec{E} = -\nabla\phi - \frac{1}{c} \frac{\partial \vec{A}}{\partial t} \quad \vec{B} = \nabla \times \vec{A}$$

$$\vec{F} = q \left(\vec{E} + \frac{q}{c} \vec{v} \times (\nabla \times \vec{A}) \right) \quad \frac{dA_i}{dt} = \frac{\partial A_i}{\partial t} + v_j \frac{\partial A_i}{\partial x_j}$$

$$\frac{dp_i}{dt} = q \left(-\underbrace{\frac{\partial \phi_i}{\partial x_i}}_{E_i} - \frac{1}{c} \underbrace{\frac{\partial A_i}{\partial t}}_{\frac{1}{c}(\nabla \times \vec{B})_i} + \frac{v_j}{c} \left(\frac{\partial A_i}{\partial x_i} - \frac{\partial A_i}{\partial x_j} \right) \right)$$

$$\begin{aligned} (\vec{v} \times (\nabla \times \vec{A}))_i &= \epsilon_{ijk} v_j \epsilon_{klm} \frac{\partial A_m}{\partial x_l} \\ &= (\epsilon_{ik} \epsilon_{jm} - \delta_{im} \delta_{jk}) v_j \frac{\partial A_m}{\partial x_k} \\ &= v_j \frac{\partial A_i}{\partial x_j} - v_j \frac{\partial A_i}{\partial x_j} \end{aligned}$$

$$\therefore \frac{d\vec{p}}{dt} = q \left(\vec{E} + \frac{\vec{v}}{c} \times \vec{B} \right) \quad \text{true}$$

Hamiltonian of Particles

$$\mathcal{L} = \mathcal{L}(x, v) \rightarrow H = H(x, p) \quad p_i = \frac{\partial \mathcal{L}}{\partial v_i} \quad H = \sum_i p_i v_i - \mathcal{L}$$

$$p_i = p_i + \frac{q}{c} A_i \quad \vec{P} = \gamma m \vec{v} + \frac{q}{c} \vec{A}$$

$$\text{free particle: } H = \sqrt{m^2 c^4 + p^2 c^2}$$

with EM fields

$$\boxed{H = \sqrt{m^2 c^4 + c^2 (\vec{p} - \frac{q}{c} \vec{A})^2} + q\phi}$$

$$\text{non-rel: } \mathcal{L} = \frac{1}{2} mv^2 - q\phi + q \frac{\vec{v}}{c} \cdot \vec{A} \quad H = \frac{1}{2m} (\vec{p} - \frac{q}{c} \vec{A})^2 + q\phi \quad \left(= \frac{p^2}{2m} + q\phi \right)$$

connection with Quantum Mechanics: $\hat{P} = -i\nabla$

$$\text{non-rel: } \hat{H} = \frac{1}{2m} (-i\nabla - \frac{q}{c} \vec{A})^2 + q\phi$$

Example: charge in uniform magnetic field

$$\vec{B} = B_0 \hat{z}$$

$$\vec{A} = A_\varphi \hat{\varphi} \quad \text{in cyl. coords}$$

$$\nabla \times \vec{A} = B_0 \hat{z} = \frac{1}{\rho} \frac{\partial}{\partial \rho} (\rho A_\varphi) \hat{z}$$

$$\frac{\partial}{\partial \rho} (\rho A_\varphi) = \rho B_0$$

$$\rho A_\varphi = \frac{1}{2} B_0 \rho^2 + \text{const}$$

$\hookrightarrow \text{set} = 0$

$$A_\varphi = \frac{1}{2} B_0 \rho$$

stationary states

$$\hat{H} \Psi = E \Psi$$

$$q = -e, e > 0$$

$$(-\nabla^2 + \frac{e^2}{c^2} A^2 - i \frac{2e}{c} \vec{A} \cdot \nabla) \Psi = 2mE \Psi \quad \Psi_{(l, \varphi, z)} = \Psi_{(l, \varphi)} e^{ikz}$$

$$\left(-\frac{1}{\rho} \frac{\partial}{\partial \rho} \left(\rho \frac{\partial}{\partial \rho} \right) - \frac{1}{\rho^2} \frac{\partial^2}{\partial \varphi^2} + \frac{e^2}{c^2} A_\varphi^2 - i \frac{2e}{c} \frac{1}{2} B_0 \rho \frac{1}{\rho} \frac{\partial}{\partial \varphi} \right) \Psi = 2mE \Psi$$

circular motion

so use:

$$\Psi_{(l, \varphi)} = R_{(l)} e^{il\varphi} \quad l: \text{integer}$$

$$\left(-\frac{1}{\rho} \frac{\partial}{\partial \rho} \left(\rho \frac{\partial}{\partial \rho} \right) - \frac{l^2}{\rho^2} + \frac{e^2 B_0^2}{4c^2} \rho^2 + \frac{e B_0}{c} l \right) R = 2mE R$$

$$\text{Ansatz } R_{(l)} \sim \rho^n e^{-\frac{l^2}{2\rho^2}} \quad \text{similar to S.H.O.} \quad (\text{want find all levels})$$

$$-\rho^{n-2} (-n^2 + l^2) = 0 \Rightarrow n = l$$

$$\rho^n \left(\frac{n+2}{\alpha^2} \rho^n + \frac{n}{\alpha^2} \rho^n - \frac{1}{\alpha^2} \rho^{n+2} + \left(\frac{l^2}{\rho^2} + \frac{e B_0}{c} l + \frac{e^2 B_0^2}{4c^2} \rho^2 \right) \rho^n \right) = 2mE \rho^n$$

$$\rho^{n+2} \left(-\frac{1}{\alpha^4} + \frac{e^2 B_0^2}{4c^2} \right) = 0 \Rightarrow \alpha^2 = \frac{2c}{e B_0} \quad \frac{2}{\alpha^2} = \frac{e B_0}{c}$$

$$2mE = \frac{2e B_0}{c} n + \frac{e B_0}{c}$$

$$E = \underbrace{\frac{e B_0}{mc}}_{w_B} (n + \frac{1}{2})$$

$$E = w_B (n + \frac{1}{2})$$

have missed some solutions

should have used $R \sim (\sum a_n \rho^n) e^{-\frac{l^2}{2\rho^2}}$

Waveguides

3/13/2007

② vacuum
 a) conductor $E_1 = 0, H_1 = 0$ $E_2^{(N)} = 4\pi \Sigma$ $\oint \vec{H} \cdot d\vec{l} = \frac{4\pi}{c} j \Delta l \Delta s \Rightarrow H_2 = \frac{4\pi}{c} K$
 ① $\sigma = \infty$ $H_2 \Delta l$ $K = \frac{f}{\Delta s} j \Delta s$

E_2 is normal, H_2 is tangential

b) if $\sigma = \text{finite}$ $E = E_0 + i \frac{4\pi\sigma}{\omega}$ there is a penetration depth

$$n = \sqrt{\epsilon} \quad \text{good conductor} \quad \frac{4\pi\sigma}{\epsilon_0 \omega} \gg 1 \quad n = \sqrt{\frac{4\pi\sigma}{\omega}} + i \sqrt{\frac{4\pi\sigma}{2\omega}}$$

$$E \sim e^{i(kx - \omega t)} \quad \text{Im}\{\epsilon\} k = \frac{1}{\delta} = \frac{\omega}{c} \sqrt{\frac{2\pi\sigma}{\omega}} = \sqrt{\frac{2\pi\omega\sigma}{c^2}} \quad \delta = \frac{c}{\sqrt{2\pi\omega\sigma}} \quad \text{skin depth}$$

fields drop to zero in region of size δ

waveguide

metal pipe
(hollow)

\hat{z} (translational sym.)

in cavity: $\nabla \times \vec{E} = -\frac{i}{c} \frac{\partial \vec{B}}{\partial t}$ $\nabla \times \vec{B} = \frac{1}{c} \frac{\partial \vec{E}}{\partial t}$

B.C. $E_{||}|_S = 0$ $B_{\perp}|_S = 0$ $\nabla \cdot \vec{E} = 0$ $\nabla \cdot \vec{B} = 0$

tangential
at surface

Ansatz: $\vec{E}(x, y, t) = \vec{E}(x, y) e^{i(kz - \omega t)}$ $(\nabla^2 - \frac{1}{c^2} \frac{\partial^2}{\partial t^2}) \vec{E} = 0$ same for \vec{B}

$$\nabla^2 = \nabla_t^2 + \frac{\partial^2}{\partial z^2} \quad : \quad (\nabla_t^2 - k^2 + \frac{\omega^2}{c^2}) \vec{E}(x, y) = 0$$

$\vec{E}(x, y) = \vec{E}_t(x, y) + E_z(x, y) \hat{z}$ B.C. $E_z|_S = 0$ ($E \perp$ surface) $\vec{B}_t \cdot \hat{n}|_S = 0$ ($B \parallel$ surface)

reduce problem to eqs + b.c. for E_z, B_z only

$$\vec{B} = \vec{B}_t + B_z \hat{z} \quad \vec{B}_t = B_x \hat{x} + B_y \hat{y} \quad \nabla \times \vec{B} = \begin{vmatrix} \hat{e}_1 & \hat{e}_2 & \hat{e}_3 \\ \partial_x & \partial_y & \partial_z \\ B_x \hat{e}_1 & B_y \hat{e}_2 & B_z \hat{e}_3 \end{vmatrix} = \left(\begin{array}{l} \left(\frac{\partial B_z}{\partial y} - i k B_y \right) \hat{e}_1 \\ + \left(i k B_x - \frac{\partial B_z}{\partial x} \right) \hat{e}_2 \\ \left(\frac{\partial B_y}{\partial x} - \frac{\partial B_x}{\partial y} \right) \hat{e}_3 \end{array} \right) e^i$$

$$\nabla \times \vec{B} = (\nabla_t \times (B_z \hat{e}_3) + i k \hat{e}_3 \times \vec{B}_t + \nabla_t \times \vec{B}_t) e^i$$

$$\nabla_t \times (B_z \hat{e}_3) + i k \hat{e}_3 \times \vec{B}_t + \nabla_t \times \vec{B}_t = -i \frac{\omega}{c} (E_t + E_z \hat{e}_3)$$

$$\Rightarrow \nabla_t \times (B_z \hat{e}_3) + i k \hat{e}_3 \times \vec{B}_t = -i \frac{\omega}{c} \vec{E}_t \quad \nabla_t \times \vec{B}_t = -i \frac{\omega}{c} E_z \hat{e}_3$$

$$(\nabla_t \times \hat{e}_3) \rightarrow \nabla_t B_z = i \frac{\omega}{c} \vec{E}_t \times \hat{e}_3 + i k \vec{B}_t \quad (*)$$

Doing the same with $\nabla \times \vec{E} = -\frac{i}{c} \frac{\partial \vec{B}}{\partial t}$: $\nabla_t E_z = -i \frac{\omega}{c} \vec{B}_t \times \hat{e}_3 + i k \vec{E}_t$

(*) $\cdot \hat{n}|_S$ $\nabla_t B_z \cdot \hat{n}|_S = \frac{\partial B_z}{\partial n}|_S$ $(\vec{E}_t \times \hat{e}_3) \cdot \hat{n}|_S = E_{||}|_S = 0$ (b.c.)

$\vec{B}_t \cdot \hat{n}|_S = 0$ (b.c.)

$$\therefore \frac{\partial B_z}{\partial n}|_S = 0 \quad E_z|_S = 0$$

$(\nabla_t^2 - k^2 + \frac{\omega^2}{c^2}) E_z(x, y) = 0$ (same for B_z)


$$-k^2 + \frac{\omega^2}{c^2} = \gamma^2 \rightarrow \text{solutions only for certain eigenvalues } \gamma_n$$

To simultaneously satisfy eigenvalue problem:

TM mode: $B_z = 0$ TE mode: $E_z = 0$ expect $\gamma^2 > 0$

TM waves: $\vec{B}_t = \frac{\omega}{ck} (\vec{E}_t \times \hat{e}_3)$ $\nabla_t E_z = -i \frac{\omega^2}{c^2 k} \vec{E}_t + i k \vec{E}_t = i \left(\frac{-\omega^2}{c^2 k} + k \right) \vec{E}_t \Rightarrow \vec{E}_t = i \frac{i k}{\gamma^2} \nabla_t E_z$


$k^2 = \frac{\omega^2}{c^2} - \gamma^2$ given mode, γ : there is a cutoff freq $\omega_c = c\gamma$ if $\omega < \omega_c \rightarrow$ no propagation





phase vel $v_p = \frac{\omega}{k} = \frac{1}{k} c \sqrt{\gamma^2 + k^2} = c \sqrt{1 + \frac{\gamma^2}{k^2}} > c \rightarrow \infty$ at cutoff

group vel $v_g = \frac{dw}{dk} = \frac{2\omega dw}{c^2} = 2k dk \quad \frac{dw}{dk} = c^2 \frac{k}{\omega} = \frac{c}{\sqrt{1 + \frac{\gamma^2}{k^2}}} < c \rightarrow 0$ at cutoff

Ex:




constructive interference $\frac{\lambda}{2L}$



only special angles give constructive interference

Example: rectangular waveguide



TE waves: $E_z = 0$ $\vec{B}_t = \frac{i k}{\delta^2} \nabla_t B_z$

$$B_z = \Psi(x, y) e^{i(kz - \omega t)}$$

$$\frac{\partial \Psi}{\partial n} \Big|_S = 0$$

$$\nabla^2 \Psi + \gamma^2 \Psi = 0$$

$$\frac{\partial \Psi}{\partial x} = 0 \quad \text{for } x=0 \quad x=a$$

$$\frac{\partial \Psi}{\partial y} = 0 \quad \text{for } y=0 \quad y=b$$

$$B_x = i \frac{k}{\delta^2} \frac{\partial \Psi}{\partial x} e^{i(kz - \omega t)} \quad B_y = i \frac{k}{\delta^2} \frac{\partial \Psi}{\partial y} e^{i(kz - \omega t)}$$

$$\vec{E}_t = \frac{\omega}{ck} \vec{B}_t \times \hat{e}_z$$

$$E_x = \frac{\omega}{ck} B_y$$

$$E_y = -\frac{\omega}{ck} B_x \quad E_z = 0$$

$$\Psi(x, y) = A \cos\left(\frac{n\pi}{a}x\right) \cos\left(\frac{m\pi}{b}y\right)$$

$$\left(\frac{n\pi}{a}\right)^2 + \left(\frac{m\pi}{b}\right)^2 = \gamma_{nm}^2 \quad n, m = 0, 1, 2, \dots$$

Dispersion relation:

$$-k^2 + \frac{\omega^2}{c^2} = \left(\frac{n\pi}{a}\right)^2 + \left(\frac{m\pi}{b}\right)^2$$

Cutoff Freq (k=0)

$$\omega_{nm} = c \gamma_{nm}$$

lowest freq.
possible

$$n=1 \quad m=0 : \quad \gamma^2 = \left(\frac{\pi}{a}\right)^2$$

$$k^2 = \frac{\omega^2}{c^2} - \left(\frac{\pi}{a}\right)^2$$

$$\text{Cutoff: } \omega_0 = c \frac{\pi}{a}$$

$$k^2 = \frac{1}{c^2} (\omega^2 - \omega_0^2) \quad \frac{\pi}{a} \frac{k}{\gamma} = \sqrt{\frac{\omega^2}{\omega_0^2} - 1}$$

$$B_z = A \cos\left(\frac{\pi}{a}x\right) e^{i(kz - \omega t)}$$


$$B_x = -i A \sqrt{\frac{\omega^2}{\omega_0^2} - 1} \sin\left(\frac{\pi}{a}x\right) e^{-i\omega t} \quad B_y = 0$$

$$= \frac{\omega^2}{c^2 k^2} (1 - \frac{\omega_0^2}{\omega^2})$$

$$E_z = 0$$

$$E_x = 0$$

$$E_y = \frac{-i}{\sqrt{1 - \frac{\omega_0^2}{\omega^2}}} B_x$$



$$B_x + E_y$$

plane wave
traveling along z

Coaxial cable: TEM mode: $E_z = B_z = 0$ (only in coaxial cable)

$$\nabla_t \times \vec{B}_t = 0$$

$$\nabla_t \times \vec{E}_t = 0$$

$$\nabla_t \cdot \vec{E}_t = 0$$

$$\nabla_t^2 \vec{E}_t = 0$$

$$(\nabla_t^2 - k^2 + \frac{\omega^2}{c^2}) \vec{E}_t = 0 \Rightarrow k = \frac{\omega}{c}$$

no cutoff freq.

Problem:



TE waves $E_z = 0$

$$B_z = \Psi(x, y) e^{i(kz - \omega t)}$$

$$(\nabla^2 + \gamma^2) \Psi = 0$$

$$B.C.$$

$$\frac{\partial \Psi}{\partial n} \Big|_S = 0$$

$$\frac{\partial \Psi}{\partial x} = 0 \quad @ x=a$$

$$\frac{\partial \Psi}{\partial y} = 0 \quad @ y=0$$

$$\nabla \Psi \cdot (-1, 1) \Big|_S = 0$$

$$\left(\frac{-\partial \Psi}{\partial x} + \frac{\partial \Psi}{\partial y} \right) \Big|_S = 0 \quad @ x=y$$

$$\Psi = XY$$

$$X = \cos\left(\frac{n\pi}{a}x\right) + \sin\left(\frac{n\pi}{a}x\right)$$

$$Y = \cos\left(\frac{m\pi}{b}y\right) + \sin\left(\frac{m\pi}{b}y\right)$$

$$\gamma^2 = \left(\frac{\pi}{a}\right)^2 (n^2 + m^2)$$

$$\omega = \frac{c\pi}{a} \sqrt{n^2 + m^2} \quad \text{Cutoff}$$