

Homework 5

Jack Brolin, Abhiram Nallamalli

July 2023

P. 1 Consider the linear programming problem:

$$\begin{aligned} \min \quad & x_1 - x_2 \\ \text{s.t.} \quad & 2x_1 + 3x_2 - x_3 + x_4 \leq 0 \\ & 3x_1 + x_2 + 4x_3 - 2x_4 \geq 3 \\ & -x_1 - x_2 + 2x_3 + x_4 = 6 \\ & x_1 \leq 0 \\ & x_2, x_3 \geq 0. \end{aligned}$$

Write down the corresponding dual problem.

Following the table from section 4.2 of the text, the corresponding dual problem is as follows:

$$\begin{aligned} \text{Max} \quad & 3p_2 + 6p_3 \\ \text{s.t.} \quad & 2p_1 + 3p_2 - p_3 \geq 1 \\ & p_1 + p_2 - p_3 \leq -1 \\ & -p_1 + 4p_2 + 2p_3 \leq 0 \\ & p_1 - 2p_2 - p_3 = 0 \\ & p_1 \leq 0 \\ & p_2 \geq 0 \\ & p_3 \text{ is free} \end{aligned}$$

P. 2 Consider a linear programming problem in standard form and assume that the rows of A are linearly independent. For each one of the following statements, provide either a proof or a counterexample.

- a. Let x^* be a basic feasible solution. Suppose that for every basis corresponding to x^* , the associated basic solution to the dual is infeasible. Then, the optimal cost must be strictly less than $c'x^*$.
- b. The dual of the auxiliary primal problem considered in Phase I of the simplex method is always feasible.
- c. Let p_i be the dual variable associated with the i^{th} equality constraint in the primal. Eliminating the i^{th} primal equality constraint is equivalent to introducing the additional constraint $p_i = 0$ in the dual problem.
- d. If the unboundedness criterion is satisfied when we solve the primal with the simplex method, then the dual problem is infeasible.

- a. True. We know that the optimal cost of the dual is less than or equal to the optimal cost of the primal. If x^* is a basic feasible solution of the primal but infeasible for the dual, then its cost must be greater than the cost of the dual.
- b. True. Since the primal auxiliary problem is always feasible, the dual is feasible too.
- c. True. Constraints are of the form $\sum p_i a_i$, so setting $p_i = 0$ effectively eliminates it.
- d. True. If the primal cost goes to $-\infty$ then the dual is infeasible.

P. 3 Let A be a symmetric square matrix. Consider the linear programming problem:

$$\begin{array}{ll}\text{minimize} & c'x \\ \text{subject to} & Ax \geq c \\ & x \geq 0\end{array}$$

Prove that if x^* satisfies $Ax^* = c$ and $x^* \geq 0$, then x^* is an optimal solution.

P. 7 Consider the following pair of linear programming problems:

$$\begin{array}{ll}\text{minimize } c'x & \text{maximize } p'b \\ \text{subject to } Ax \geq b & \text{subject to } p'A \leq c' \\ x \geq 0, & p \geq 0.\end{array}$$

Suppose at least one of these two problems has a feasible solution. Prove that the set of feasible solutions to at least one of the two problems is unbounded.

P. 10 Consider the problem:

$$\begin{aligned} \min \quad & \max_{i=1,\dots,m} (a'_i x - b_i) \\ \text{s.t.} \quad & x \in \mathbb{R}^n. \end{aligned}$$

Let v be the optimal value of the above optimization problem, let A be the $m \times n$ matrix whose rows are a'_1, \dots, a'_m , and let b be the vector whose components are b_1, \dots, b_m .

- a. Consider any vector $p \in \mathbb{R}^m$ such that $p'A = 0'$, $p \geq 0$, and $\sum_{i=1}^m p_i = 1$. Show that $-p'b \leq v$.
- b. In order to obtain the best possible lower bound of the form obtained in part (a), we consider the linear program:

$$\begin{aligned} \max \quad & -p'b \\ \text{s.t.} \quad & p'A = 0' \\ & \sum_{i=1}^m p_i = 1 \\ & p \geq 0. \end{aligned}$$

Show that the optimal cost of the above linear program is v .