




Jiangxi University of Science and Technology

# Discrete Mathematics and Its Applications

Section 10.4 Connectivity

Section 10.5


Euler and Hamiltonian Graphs



# Acknowledgement

---

Most of these slides are adapted from ones created by  
Professor Bart Selman at Cornell University ,  
and Dr Johnnie Baker and **Discrete Mathematics and Its  
Applications** (Seventh Edition) **Kenneth H. Rosen**



# Connectivity Section 10.4

## Section Summary

---

- Paths
- Connectedness in Undirected Graphs
- Vertex Connectivity and Edge Connectivity (*not currently included in overheads*)
- Connectedness in Directed Graphs
- Paths and Isomorphism (*not currently included in overheads*)
- Counting Paths between Vertices

# Paths

---

**Informal Definition:** A *path* is a sequence of edges that begins at a vertex of a graph and travels from vertex to vertex along edges of the graph. As the path travels along its edges, it visits the vertices along this path, that is, the endpoints of these.

**Applications:** Numerous problems can be modeled with paths formed by traveling along edges of graphs such as:

- determining whether a message can be sent between two computers.
- efficiently planning routes for mail delivery.

# Paths

**Definition:** Let  $n$  be a nonnegative integer and  $G$  an undirected graph. A *path* of length  $n$  from  $u$  to  $v$  in  $G$  is a sequence of  $n$  edges  $e_1, \dots, e_n$  of  $G$  for which there exists a sequence  $x_0 = u, x_1, \dots, x_{n-1}, x_n = v$  of vertices such that  $e_i$  has, for  $i = 1, \dots, n$ , the endpoints  $x_{i-1}$  and  $x_i$ .


- When the graph is simple, we denote this path by its vertex sequence  $x_0, x_1, \dots, x_n$  (since listing the vertices uniquely determines the path).
- The path is a *circuit* if it begins and ends at the same vertex ( $u = v$ ) and has length greater than zero.
- The path or circuit is said to *pass through* the vertices  $x_1, x_2, \dots, x_{n-1}$  and *traverse* the edges  $e_1, \dots, e_n$ .
- A path or circuit is *simple* if it does not contain the same edge more than once.

This terminology is readily extended to directed graphs.  
(see text)

## Paths (*continued*)


**Example:** In the simple graph here:

- $a, d, c, f, e$  is a simple path of length 4.
- $d, e, c, a$  is not a path because  $e$  is not connected to  $c$ .
- $b, c, f, e, b$  is a circuit of length 4.
- $a, b, e, d, a, b$  is a path of length 5, but it is not a simple path.




# Degrees of Separation

**Example: *Paths in Acquaintanceship Graphs.*** In an acquaintanceship graph there is a path between two people if there is a chain of people linking these people, where two people adjacent in the chain know one another. In this graph there is a chain of six people linking Kamini and Ching.



Some have speculated that almost every pair of people in the world are linked by a small chain of no more than six, or maybe even, five people. The play *Six Degrees of Separation* by John Guare is based on this notion.

# Erdős numbers



Paul Erdős

## Example: *Erdős numbers*.

In a collaboration graph, two people  $a$  and  $b$  are connected by a path when there is a sequence of people starting with  $a$  and ending with  $b$  such that the endpoints of each edge in the path are people who have collaborated.

• In the academic collaboration graph of people who have written papers in mathematics, the *Erdős number* of a person  $m$  is the length of the shortest path between  $m$  and the prolific mathematician Paul Erdős.

— To learn more about Erdős numbers, visit

<http://www.ams.org/mathscinet/collaborationDistance.html>

**TABLE 1** The Number of Mathematicians with a Given Erdős Number (as of early 2006).

<i>Erdős Number</i>	<i>Number of People</i>
0	1
1	504
2	6,593
3	33,605
4	83,642
5	87,760
6	40,014
7	11,591
8	3,146
9	819
10	244
11	68
12	23
13	5



# Bacon Numbrers

In the Hollywood graph, two actors  $a$  and  $b$  are linked when there is a chain of actors linking  $a$  and  $b$ , where every two actors adjacent in the chain have acted in the same movie.

The *Bacon number* of an actor  $c$  is defined to be the length of the shortest path connecting  $c$  and the well-known actor Kevin Bacon. (Note that we can define a similar number by replacing Kevin Bacon by a different actor.)

- The *oracle of Bacon* web site <http://oracleofbacon.org/how.php> provides a tool for finding Bacon numbers.


**TABLE 2** The Number of Actors with a Given Bacon Number (as of early 2011).

<i>Bacon Number</i>	<i>Number of People</i>
0	1
1	2,367
2	242,407
3	785,389
4	200,602
5	14,048
6	1,277
7	114
8	16


# Connectedness in Undirected Graphs

**Definition:** An undirected graph is called *connected* if there is a path between every pair of vertices. An undirected graph that is not *connected* is called *disconnected*. We say that we *disconnect* a graph when we remove vertices or edges, or both, to produce a disconnected subgraph.

**Example:**  $G_1$  is connected because there is a path between any pair of its vertices, as can be easily seen. However  $G_2$  is not connected because there is no path between vertices  $a$  and  $f$ , for example.



$G_1$




$G_2$

# Connected Components

---

**Definition:** A *connected component* of a graph  $G$  is a connected subgraph of  $G$  that is not a proper subgraph of another connected subgraph of  $G$ . A graph  $G$  that is not connected has two or more connected components that are disjoint and have  $G$  as their union.

**Example:** The graph  $H$  is the union of three disjoint subgraphs  $H_1$ ,  $H_2$ , and  $H_3$ , none of which are proper subgraphs of a larger connected subgraph of  $G$ . These three subgraphs are the connected components of  $H$ .



# Connectedness in Directed Graphs


---

**Definition:** A directed graph is *strongly connected* if there is a path from  $a$  to  $b$  and a path from  $b$  to  $a$  whenever  $a$  and  $b$  are vertices in the graph.


**Definition:** A directed graph is *weakly connected* if there is a path between every two vertices in the underlying undirected graph, which is the undirected graph obtained by ignoring the directions of the edges of the directed graph.

# Connectedness in Directed Graphs (*continued*)

**Example:**  $G$  is strongly connected because there is a path between any two vertices in the directed graph. Hence,  $G$  is also weakly connected.



The graph  $H$  is not strongly connected, since there is no directed path from  $a$  to  $b$ , but it is weakly connected.



**Definition:** The subgraphs of a directed graph  $G$  that are strongly connected but not contained in larger strongly connected subgraphs, that is, the maximal strongly connected subgraphs, are called the *strongly connected components* or *strong components* of  $G$ .

**Example (continued):** The graph  $H$  has three strongly connected components, consisting of the vertex  $a$ ; the vertex  $e$ ; and the subgraph consisting of the vertices  $b, c, d$  and edges  $(b, c)$ ,  $(c, d)$ , and  $(d, b)$ .

# The Connected Components of the Web Graph

- Recall that at any particular instant the web graph provides a snapshot of the web, where vertices represent web pages and edges represent links. According to a 1999 study, the Web graph at that time had over 200 million vertices and over 1.5 billion edges. (The numbers today are several orders of magnitude larger.)
- The underlying undirected graph of this Web graph has a connected component that includes approximately 90% of the vertices.
- There is a *giant strongly connected component (GSCC)* consisting of more than 53 million vertices. A Web page in this component can be reached by following links starting in any other page of the component. There are three other categories of pages with each having about 44 million vertices:
  - pages that can be reached from a page in the GSCC, but do not link back.
  - pages that link back to the GSCC, but can not be reached by following links from pages in the GSCC.
  - pages that cannot reach pages in the GSCC and can not be reached from pages in the GSCC

# Counting Paths between Vertices

---

- We can use the adjacency matrix of a graph to find the number of paths between two vertices in the graph.

**Theorem:** Let  $G$  be a graph with adjacency matrix  $\mathbf{A}$  with respect to the ordering  $v_1, \dots, v_n$  of vertices (with directed or undirected edges, multiple edges and loops allowed). The number of different paths of length  $r$  from  $v_i$  to  $v_j$ , where  $r > 0$  is a positive integer, equals the  $(i,j)$ th entry of  $\mathbf{A}^r$ .

*Proof by mathematical induction:*

*Basis Step:* By definition of the adjacency matrix, the number of paths from  $v_i$  to  $v_j$  of length 1 is the  $(i,j)$ th entry of  $\mathbf{A}$ .

*Inductive Step:* For the inductive hypothesis, we assume that the  $(i,j)$ th entry of  $\mathbf{A}^r$  is the number of different paths of length  $r$  from  $v_i$  to  $v_j$ .


– Because  $\mathbf{A}^{r+1} = \mathbf{A}^r \mathbf{A}$ , the  $(i,j)$ th entry of  $\mathbf{A}^{r+1}$  equals  $b_{i1}a_{1j} + b_{i2}a_{2j} + \dots + b_{in}a_{nj}$ , where  $b_{ik}$  is the  $(i,k)$ th entry of  $\mathbf{A}^r$ . By the inductive hypothesis,  $b_{ik}$  is the number of paths of length  $r$  from  $v_i$  to  $v_k$ .

– A path of length  $r + 1$  from  $v_i$  to  $v_j$  is made up of a path of length  $r$  from  $v_i$  to some  $v_k$ , and an edge from  $v_k$  to  $v_j$ . By the product rule for counting, the number of such paths is the product of the number of paths of length  $r$  from  $v_i$  to  $v_k$  (i.e.,  $b_{ik}$ ) and the number of edges from  $v_k$  to  $v_j$  (i.e.,  $a_{kj}$ ). The sum over all possible intermediate vertices  $v_k$  is  $b_{i1}a_{1j} + b_{i2}a_{2j} + \dots + b_{in}a_{nj}$ .



# Counting Paths between Vertices (*continued*)

**Example:** How many paths of length four are there from  $a$  to  $d$  in the graph  $G$ .



$$A = \begin{bmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix}$$

*adjacency  
matrix of  $G$*

**Solution:** The adjacency matrix of  $G$  (ordering the vertices as  $a, b, c, d$ ) is given above. Hence the number of paths of length four from  $a$  to  $d$  is the  $(1, 4)$ th entry of  $A^4$ . The eight paths are as:

$$A^4 = \begin{bmatrix} 8 & 0 & 0 & 8 \\ 0 & 8 & 8 & 0 \\ 0 & 8 & 8 & 0 \\ 8 & 0 & 0 & 8 \end{bmatrix}$$

$a, b, a, b, d$	$a, b, a, c, d$
$a, b, d, b, d$	$a, b, d, c, d$
$a, c, a, b, d$	$a, c, a, c, d$
$a, c, d, b, d$	$a, c, d, c, d$





# Section 10.5: Euler and Hamiltonian Graphs

## Section Summary

---


- Euler Paths and Circuits
- Hamilton Paths and Circuits
- Applications of Hamilton Circuits

# Euler Paths and Circuits




Leonard Euler (1707-1783)

- The town of Königsberg, Prussia (now Kaliningrad, Russia) was divided into four sections by the branches of the Pregel river. In the 18th century seven bridges connected these regions.
- People wondered whether whether it was possible to follow a path that crosses each bridge exactly once and returns to the starting point.
- The Swiss mathematician Leonard Euler proved that no such path exists. This result is often considered to be the first theorem ever proved in graph theory.



The 7 Bridges of Königsberg



Multigraph Model  
of the Bridges of  
Königsberg


# Euler Paths and Circuits (*continued*)

**Definition:** An *Euler circuit* in a graph  $G$  is a simple circuit containing every edge of  $G$ . An *Euler path* in  $G$  is a simple path containing every edge of  $G$ .


**Example:** Which of the undirected graphs  $G_1$ ,  $G_2$ , and  $G_3$  has a Euler circuit? Of those that do not, which has an Euler path?

**Solution:**


The graph  $G_1$  has an Euler circuit (e.g.,  $a, e, c, d, e, b, a$ ). But, as can easily be verified by inspection, neither  $G_2$  nor  $G_3$  has an Euler circuit. Note that  $G_3$  has an Euler path (e.g.,  $a, c, d, e, b, d, a, b$ ), but there is no Euler path in  $G_2$ , which can be verified by inspection.



$G_1$



$G_2$



$G_3$

# Necessary Conditions for Euler Circuits and Paths

---

An Euler circuit begins with a vertex  $a$  and continues with an edge incident with  $a$ , say  $\{a, b\}$ . The edge  $\{a, b\}$  contributes one to  $\deg(a)$ .

Each time the circuit passes through a vertex it contributes two to the vertex's degree.

Finally, the circuit terminates where it started, contributing one to  $\deg(a)$ . Therefore  $\deg(a)$  must be even.

We conclude that the degree of every other vertex must also be even.


By the same reasoning, we see that the initial vertex and the final vertex of an Euler path have odd degree, while every other vertex has even degree. So, a graph with an Euler path has exactly two vertices of odd degree.

In the next slide we will show that these necessary conditions are also sufficient conditions.

# Sufficient Conditions for Euler Circuits and Paths

Suppose that  $G$  is a connected multigraph with  $\geq 2$  vertices, all of even degree. Let  $x_0 = a$  be a vertex of even degree. Choose an edge  $\{x_0, x_1\}$  incident with  $a$  and proceed to build a simple path  $\{x_0, x_1\}, \{x_1, x_2\}, \dots, \{x_{n-1}, x_n\}$  by adding edges one by one until another edge can not be added.

We illustrate this idea in the graph  $G$  here. We begin at  $a$  and choose the edges  $\{a, f\}$ ,  $\{f, c\}$ ,  $\{c, b\}$ , and  $\{b, a\}$  in succession.




- 
- The path begins at  $a$  with an edge of the form  $\{a, x\}$ ; we show that it must terminate at  $a$  with an edge of the form  $\{y, a\}$ . Since each vertex has an even degree, there must be an even number of edges incident with this vertex. Hence, every time we enter a vertex other than  $a$ , we can leave it. Therefore, the path can only end at  $a$ .
  - If all of the edges have been used, an Euler circuit has been constructed. Otherwise, consider the subgraph  $H$  obtained from  $G$  by deleting the edges already used.

In the example  $H$  consists of the vertices  $c, d, e$ .

# Sufficient Conditions for Euler Circuits and Paths (*continued*)

• Because  $G$  is connected,  $H$  must have at least one vertex in common with the circuit that has been deleted.



• Every vertex in  $H$  must have even degree because all the vertices in  $G$  have even degree and for each vertex, pairs of edges incident with this vertex have been deleted. Beginning with the shared vertex construct a path ending in the same vertex (as was done before). Then splice this new circuit into the original circuit.

In the example, the vertex is  $c$ .

- 
- Continue this process until all edges have been used. This produces an Euler circuit. Since every edge is included and no edge is included more than once.
  - Similar reasoning can be used to show that a graph with exactly two vertices of odd degree must have an Euler path connecting these two vertices of odd degree

In the example, we end up with the circuit  $a, f, c, d, e, c, b, a$ .



# Algorithm for Constructing an Euler Circuits

---

In our proof we developed this algorithms for constructing a Euler circuit in a graph with no vertices of odd degree.

```
procedure Euler( $G$ : connected multigraph with all vertices of even degree)
   $circuit :=$  a circuit in  $G$  beginning at an arbitrarily chosen vertex with edges
    successively added to form a path that returns to this vertex.
   $H := G$  with the edges of this circuit removed
  while  $H$  has edges
     $subcircuit :=$  a circuit in  $H$  beginning at a vertex in  $H$  that also is
      an endpoint of an edge in circuit.
     $H := H$  with edges of  $subcircuit$  and all isolated vertices removed
     $circuit := circuit$  with  $subcircuit$  inserted at the appropriate vertex.
  return  $circuit$ { $circuit$  is an Euler circuit}
```




# Necessary and Sufficient Conditions for Euler Circuits and Paths (*continued*)

---




**Theorem:** A connected multigraph with at least two vertices has an Euler circuit if and only if each of its vertices has an even degree and it has an Euler path if and only if it has exactly two vertices of odd degree.

**Example:** Two of the vertices in the multigraph model of the Königsberg bridge problem have odd degree. Hence, there is no Euler circuit in this multigraph and it is impossible to start at a given point, cross each bridge exactly once, and return to the starting point.



# Euler Circuits and Paths

**Example:**



$G_1$  contains exactly two vertices of odd degree ( $b$  and  $d$ ). Hence it has an Euler path, e.g.,  $d, a, b, c, d, b$ .

$G_2$  has exactly two vertices of odd degree ( $b$  and  $d$ ). Hence it has an Euler path, e.g.,  $b, a, g, f, e, d, c, g, b, c, f, d$ .

$G_3$  has six vertices of odd degree. Hence, it does not have an Euler path.

# Applications of Euler Paths and Circuits

---


Euler paths and circuits can be used to solve many practical problems such as finding a path or circuit that traverses each

- street in a neighborhood,
- road in a transportation network,
- connection in a utility grid,
- link in a communications network.

Other applications are found in the


- layout of circuits,
- network multicasting,
- molecular biology, where Euler paths are used in the sequencing of DNA.

# Hamilton Paths and Circuits



William Rowan  
Hamilton  
(1805- 1865)

- Euler paths and circuits contained every edge only once. Now we look at paths and circuits that contain every vertex exactly once.
  - William Hamilton invented the *Icosian puzzle* in 1857. It consisted of a wooden dodecahedron (with 12 regular pentagons as faces), illustrated in (a), with a peg at each vertex, labeled with the names of different cities. String was used to plot a circuit visiting 20 cities exactly once
  - The graph form of the puzzle is given in (b).
- 
- The solution (a Hamilton circuit) is given here.



# Hamilton Paths and Circuits

---

**Definition:** A simple path in a graph  $G$  that passes through every vertex exactly once is called a *Hamilton path*, and a simple circuit in a graph  $G$  that passes through every vertex exactly once is called a *Hamilton circuit*.

That is, a simple path  $x_0, x_1, \dots, x_{n-1}, x_n$  in the graph  $G = (V, E)$  is called a Hamilton path if  $V = \{x_0, x_1, \dots, x_{n-1}, x_n\}$  and  $x_i \neq x_j$  for  $0 \leq i < j \leq n$ , and the simple circuit  $x_0, x_1, \dots, x_{n-1}, x_n, x_0$  (with  $n > 0$ ) is a Hamilton circuit if  $x_0, x_1, \dots, x_{n-1}, x_n$  is a Hamilton path.


# Hamilton Paths and Circuits (*continued*)

**Example:** Which of these simple graphs has a Hamilton circuit or, if not, a Hamilton path?

**Solution:**  $G_1$  has a Hamilton circuit:  $a, b, c, d, e, a$ .

$G_2$  does not have a Hamilton circuit (Why?), but does have a Hamilton path :  $a, b, c, d$ .

$G_3$  does not have a Hamilton circuit, or a Hamilton path. Why?



# Necessary Conditions for Hamilton Circuits

---

- Unlike for an Euler circuit, no simple necessary and sufficient conditions are known for the existence of a Hamilton circuit.
- However, there are some useful necessary conditions. We describe two of these now.

**Dirac's Theorem:** If  $G$  is a simple graph with  $n \geq 3$  vertices such that the degree of every vertex in  $G$  is  $\geq n/2$ , then  $G$  has a Hamilton circuit.

**Ore's Theorem:** If  $G$  is a simple graph with  $n \geq 3$  vertices such that  $\deg(u) + \deg(v) \geq n$  for every pair of nonadjacent vertices, then  $G$  has a Hamilton circuit.



Gabriel Andrew Dirac  
(1925-1984)



Øysten Ore  
(1899-1968)




# Applications of Hamilton Paths and Circuits

---

Applications that ask for a path or a circuit that visits each intersection of a city, each place pipelines intersect in a utility grid, or each node in a communications network exactly once, can be solved by finding a Hamilton path in the appropriate graph.

The famous *traveling salesperson problem* (*TSP*) asks for the shortest route a traveling salesperson should take to visit a set of cities. This problem reduces to finding a Hamilton circuit such that the total sum of the weights of its edges is as small as possible.

A family of binary codes, known as *Gray codes*, which minimize the effect of transmission errors, correspond to Hamilton circuits in the  $n$ -cube  $Q_n$ . (*See the text for details.*)



There is no end to education. It is not  
that you read a book, pass an  
examination, and finish with education.  
The whole of life, from the moment  
you are born to the moment you die, is  
a process of learning.

— *Jiddu Krishnamurti* —

AZ QUOTES