

# Seagrass ecosystems as a significant global carbon stock

James W. Fourqurean<sup>a \*</sup>

Carlos M. Duarte<sup>b,c</sup>

Hilary Kennedy<sup>d</sup>

Núria Marbà<sup>b</sup>

Marianne Holmer<sup>e</sup>

Miguel Angel Mateo<sup>f</sup>

Eugenia T. Apostolaki<sup>g</sup>

Gary A. Kendrick<sup>c</sup>

Dorte Krause-Jensen<sup>h</sup>

Karen J. McGlathery<sup>i</sup>

Oscar Serrano<sup>f</sup>

a. Department of Biological sciences and Southeast Environmental Research Center, Marine Science Program, Florida International University, 3000 NE 151<sup>st</sup> St, North Miami, FL 33181, USA

b. Department of Global Change Research. IMEDEA (CSIC-UIB) Institut Mediterrani d'Estudis Avançats, C/ Miguel Marqués 21, 07190 Esporles (Mallorca) Spain

c. The UWA Oceans Institute, University of Western Australia, 35 Stirling Highway, Crawley 6009, Australia

d. School of Ocean Sciences, College of Natural Sciences, Bangor University, Askew Street  
Menai Bridge, Anglesey LL59 5AB, Wales, United Kingdom

e. Institute of Biology, University of Southern Denmark, Campusvej 55, DK-5230 Odense M, Denmark


f. Centre for Advanced Studies of Blanes, (CEAB-CSIC), Acceso Cala S. Francesc 14, 17300 Blanes, Spain

g. Institute of Oceanography, Hellenic Centre for Marine Research, PO Box 2214, 71003, Heraklion – Crete, Greece

h. Department of Bioscience, Aarhus University, Vejlsøvej 25, DK-8600 Silkeborg, Denmark

i. Department of Environmental Sciences, PO Box 400123, Clark Hall, University of Virginia  
Charlottesville, VA 22904, USA


# Seagrasses have a broad global distribution



Green and Short 2003




**Sediment stabilizers: seagrasses efficiently hold sediments in place, preventing resuspension and movement of sediment deposits**



(c) Thad Murdoch 2005




**Sediment stabilizers: seagrasses efficiently hold sediments in place, preventing resuspension and movement of sediment deposits**



*Thalassia testudinum*

(c) Thad Murdoch 2005

# Measuring C stored in seagrass soils: Piston corer to collect uncompressed cores





# Measuring C stored in living biomass








## Need:


- volumetric measures of Dry Bulk Density (mass of soil per volume)
- Carbon content of soil (as a fraction of mass)
  - Organic matter, or Loss on Ignition (LOI)
  - $C_{org}$





# Case study: C stores in Florida Bay seagrasses









**$C_{org}$  generally  
decreases downcore  
in Florida Bay  
seagrass soils.**

**Buried peats have  
high  $C_{org}$**




**DBD generally  
increases downcore  
in Florida Bay  
seagrass soils.**

**Buried peats have  
low DBD**






# $C_{org}$ stocks in top m of seagrass beds



**There are about 18,000 km<sup>2</sup> of seagrass beds in south Florida**





**A very rough estimate of carbon stored in the top meter of seagrass soils in south Florida:**

**18,000 km<sup>2</sup> of seagrasses**

**594 tons CO<sub>2</sub>e ha<sup>-1</sup>**

**1 x 10<sup>9</sup> tons CO<sub>2</sub>e stored in the soils!**

**A very rough estimate of carbon stored in the top meter of seagrass soils in south Florida:**

**18,000 km<sup>2</sup> of seagrasses**


**594 tons CO<sub>2</sub>e ha<sup>-1</sup>**

**1 x 10<sup>9</sup> tons CO<sub>2</sub>e stored in the soils!**

**Anthropogenic CO<sub>2</sub>e flux is about 29 x 10<sup>9</sup> tons y<sup>-1</sup>**


# Distribution of seagrass C stock data

# Global distribution of seagrass soil DBD






# LOI can be used to predict $C_{org}$



# Global distribution of soil $C_{org}$ in



# Global values for Seagrass abundance


	n	Range	Median	Mean $\pm$ 95%CI
Aboveground biomass Mg (dry weight) ha <sup>-1</sup>	251	0.002 – 15.850	0.756	2.151 $\pm$ 0.364
Belowground biomass Mg (dry weight) ha <sup>-1</sup>	251	0.002 – 50.957	1.542	5.008 $\pm$ 1.069
Total seagrass biomass Mg (dry weight) ha <sup>-1</sup>	251	0.004 – 66.807	2.854	7.159 $\pm$ 1.399
Dry Bulk Density g (dry weight) mL <sup>-1</sup>	2484	0.06 – 2.35	0.92	1.03 $\pm$ 0.02
Porosity %	1687	18.0 – 99.7	61.6	59.2 $\pm$ 0.8
Organic matter as LOI % of dry weight	2783	0.0 – 100.0	4.2	5.7 $\pm$ 0.3

# Global ranges of C<sub>org</sub> density


	n	Range	Median	Mean ± 95%CI
Aboveground biomass Mg (C <sub>org</sub> ) ha <sup>-1</sup>	251	0.001 – 5.548	0.264	0.755 ± 0.128
Belowground biomass Mg (C <sub>org</sub> ) ha <sup>-1</sup>	251	0.001 – 17.835	0.540	1.756 ± 0.375
Total seagrass biomass Mg (C <sub>org</sub> ) ha <sup>-1</sup>	251	0.001 – 23.382	1.000	2.514 ± 0.489
Soil C <sub>org</sub>	2535	0 – 48.2	1.8	2.5 ± 0.1
% of dry weight	3561	0 – 48.2	1.4	2.0 ± 0.1
Dry Bulk Density g (dry weight) mL <sup>-1</sup>	2484	0.06 – 2.35	0.92	1.03 ± 0.02



# Global estimates of seagrass soil $C_{org}$



# Some seagrass beds rival C-rich



# Seagrass bioregions

# Regional estimates of Seagrass C<sub>org</sub>


Region	Living Seagrass Biomass MgC ha <sup>-1</sup>			Soil C <sub>org</sub> MgC ha <sup>-1</sup>	
	n	Mean ± 95%CI		n	Mean ± 95%CI
Northeast Pacific	5	0.97 ± 1.02		1	64.4
Southeast Pacific	0	ND		0	ND
North Atlantic	50	0.85 ± 0.19		24	48.7 ± 14.5
Tropical Western Atlantic	44	0.84 ± 0.17		13	150.9 ± 26.3
Mediterranean	57	7.29 ± 1.52		29	372.4 ± 74.5
South Atlantic	5	1.06 ± 0.51		5	137.0 ± 56.8
Indopacific	47	0.61 ± 0.26		8	23.6 ± 8.3
Western Pacific	0	ND		0	ND
South Australia	40	2.32 ± 0.63		9	268.3 ± 101.7
Global Average	251	2.51 ± 0.49		89	194.2 ± 20.2



# How big are Global Seagrass Blue


- 300,000-600,000 km<sup>2</sup> of seagrasses
- Median estimate of seagrass biomass:  
**2.5 Mg C<sub>org</sub> ha<sup>-1</sup>**
- Median estimate of seagrass soil C<sub>org</sub> (top meter)  
**139.7 Mg C<sub>org</sub> ha<sup>-1</sup>**
- Global seagrass biomass:  
**75.5 and 151.0 Tg C**
- Global seagrass Soil C<sub>org</sub>:  
**4.2 - 8.4 Pg C**  
(earlier estimate of salt marshes and mangrove combined is 10 Pg C)

**Reports of seagrass losses  
and the rates of decline are  
increasing dramatically**



Waycott et al. 2009 *PNAS*

# Seagrass ecosystems are declining globally



Waycott et al. 2009 *PNAS*

# What are the consequences of seagrass

- Seagrass loss has averaged  $1.5\% \text{ y}^{-1}$  since the beginning of the 20<sup>th</sup> century
- Resulting loss of seagrass biomass:  
 **$11.3 - 22.7 \text{ Tg C y}^{-1}$**
- Resulting loss of seagrass soil  $\text{C}_{\text{org}}$  (top meter)  
 **$63 - 297 \text{ Tg C y}^{-1}$**
- These rates are roughly 10% of total  $\text{CO}_2$  fluxes