

Anetone cough medicine sachets

| item | Content |
|--|---|
| Pharmaceutical classification | Over-the-counter drugs |
| Medicinal efficacy classification | Antitussive expectorant |
| Approved brand name | |
| product name | Anetone cough medicine sachets |
| Product name (reading) | Aneton Sekidome Ekibunpou |
| Product features | Anetone cough medicine sachets relieve coughs caused by various causes and improve the removal of phlegm. It is a refreshing lemon tea-flavored liquid medicine, and it is a one-drink type that is convenient to carry. |
| Precautions for use | <p>■What you should not do [If you do not follow these rules, your current symptoms may get worse, or side effects and accidents may occur more easily]</p> <ol style="list-style-type: none"> The following people should not take this drug : (1) People who have had allergic symptoms due to this drug or its ingredients. (2) Children under 15 years of age. Do not use any of the following medicines while taking this drug. Other antitussive expectorants, cold medicines, sedatives, oral medicines containing antihistamines, etc. (oral medicines for rhinitis, motion sickness medicine, allergy medicines, etc.) etc.) After taking this medicine, do not drive a vehicle or operate machinery (you may experience drowsiness, etc.) If you are breastfeeding, please do not take this drug or avoid breastfeeding if you are taking this drug⁵ Do not take an overdose or use it for a long period of time (feelings of fatigue or weakness may occur.) <p>■To consult</p> <ol style="list-style-type: none"> The following people should consult a doctor, pharmacist, or registered salesperson before taking this product: (1) People receiving treatment from a doctor. (2) Pregnant women or people who appear to be pregnant. (3) Elderly people. (4) People who have had allergic symptoms due to drugs, etc. (5) People with the following symptoms. High fever, difficulty urinating (6) Persons with the following diagnosis. Heart disease, hypertension, diabetes, glaucoma, thyroid dysfunction, respiratory dysfunction, obstructive sleep apnea syndrome, |

| | <p>obesity²</p> <p>. If the following symptoms appear after taking this medicine, it may be a side effect, so please stop taking it immediately and consult your doctor, pharmacist, or registered salesperson with this document.</p> <p>[Related areas: Symptoms]</p> <p>Skin: Rash/redness, itching</p> <p>Digestive system: Nausea, vomiting, loss of appetite</p> <p>Neurological system: Dizziness</p> <p>Urinary system: Difficulty urinating</p> <p>The following serious symptoms may occur in rare cases. In that case, please see a doctor immediately.</p> <p>[Symptom name: Symptoms]</p> <p>Aplastic anemia: Bruises, nosebleeds, bleeding gums, fever, pale skin and mucous membranes, fatigue, palpitations, shortness of breath, feeling sick and dizzy, hematuria, etc. Appears.</p> <p>Agranulocytosis: Sudden high fever, chills, sore throat, etc. appear.</p> <p>Respiratory depression: Shortness of breath, difficulty breathing, etc. occur.</p> <p>3. The following symptoms may appear after taking this medicine. If these symptoms persist or worsen, stop taking this medicine and consult your doctor, pharmacist, or registered salesperson with this document.</p> <p>Constipation</p> <p>, dry mouth, drowsiness</p> <p>4. If your symptoms do not improve even after taking 5 to 6 times, stop taking this medicine and consult your doctor, pharmacist, or registered salesperson with this document.</p> | | | | | | | | | | | | |
|-------------------------------------|--|-----------|----------|---------------------------|------|----------------------------------|------|--------------------------|------|--------------------|------|----------------------|--------|
| Efficacy/effect | cough, phlegm | | | | | | | | | | | | |
| Efficacy-related precautions | | | | | | | | | | | | | |
| Usage/Dosage | <p>Please take the following amount.</p> <p>[Age: 1 dose: Number of doses per day]</p> <p>Adults (15 years and older): 1 packet (10 mL): 3 times. Depending on your symptoms, you can take this medicine up to 6 times a day at intervals of about 4 hours</p> <p>.</p> <p>Under 15 years of age: Do not take.</p> | | | | | | | | | | | | |
| Usage-related notes | <p>(1) Please strictly follow the directions and dosage.</p> <p>(2) Take one packet at a time, and do not leave any left behind.</p> <p>(3) Be careful not to spill the liquid when opening the package or taking it.</p> | | | | | | | | | | | | |
| Ingredient amount | <p>6 sachets (60mL)</p> <table border="1"> <thead> <tr> <th>component</th><th>quantity</th></tr> </thead> <tbody> <tr> <td>codeine phosphate hydrate</td><td>50mg</td></tr> <tr> <td>dl-methylephedrine hydrochloride</td><td>75mg</td></tr> <tr> <td>Chlorpheniramine maleate</td><td>12mg</td></tr> <tr> <td>anhydrous caffeine</td><td>60mg</td></tr> <tr> <td>Senega style extract</td><td>1500mg</td></tr> </tbody> </table> | component | quantity | codeine phosphate hydrate | 50mg | dl-methylephedrine hydrochloride | 75mg | Chlorpheniramine maleate | 12mg | anhydrous caffeine | 60mg | Senega style extract | 1500mg |
| component | quantity | | | | | | | | | | | | |
| codeine phosphate hydrate | 50mg | | | | | | | | | | | | |
| dl-methylephedrine hydrochloride | 75mg | | | | | | | | | | | | |
| Chlorpheniramine maleate | 12mg | | | | | | | | | | | | |
| anhydrous caffeine | 60mg | | | | | | | | | | | | |
| Senega style extract | 1500mg | | | | | | | | | | | | |

| | |
|---|---|
| Additive | Sodium citrate, citric acid, sodium benzoate, methylparaben, propylene glycol, glycerin, D-sorbitol, caramel, fragrance, ethanol, pH regulator |
| Precautions for storage and handling | <p>(1) Store in a cool place away from direct sunlight.</p> <p>(2) Please keep out of reach of children.</p> <p>(3) Do not transfer to another container. (It may cause misuse or change in quality.)</p> <p>(4) Do not take the product after the expiration date (written on the outer box).</p> <p>(5) When carrying, be careful not to bend the packet.</p> |
| Consumer consultation desk | <p>Company name: Johnson & Johnson Co., Ltd.</p> <p>Contact: Customer Service Office</p> <p>Telephone: 0120-834389 (toll-free)</p> <p>Reception hours: 9:00-17:00 (excluding Saturdays, Sundays, and holidays)</p> |
| Manufacturing and sales company | <p>Panakeia Pharmaceutical Co., Ltd. Package insert information</p> <p>Company name: Panakeia Pharmaceutical Co., Ltd.</p> <p>Address: 4576 Nakata, Takaoka City, Toyama Prefecture 939-1275</p> |
| Sales company | Johnson & Johnson Co., Ltd. |
| Dosage form | liquid agent |
| Risk classification etc. | "2" class drugs |