

# Bias-Variance Tradeoffs in Joint Spectral Embeddings


Benjamin Draves   Daniel Sussman

Boston University

Joint Statistical Meetings, 3 August 2020


# Multiplex Networks

- Multiplex networks encode multiple relationships between entities as a collection of networks.




- Application areas; International Trade, Transportation Systems, Terrorist Groups, Neuroscience (Kivelä et al. 2014).

# Individual Spectral Embeddings



# Joint Spectral Embeddings



# Analysis Framework

- Consider  $m$  graphs over a common vertex set  $\mathcal{V}$  of size  $n$
- Associate  $v \in \mathcal{V}$  with a *latent position*  $\mathbf{X}_v \in \mathbb{R}^d$

## Inner Product Distribution

Let  $F$  be a probability distribution over  $\mathbb{R}^d$ . We say  $F$  is a  *$d$ -dimensional inner product distribution* if all  $\mathbf{x}, \mathbf{y} \in \text{supp}(F)$  has the property  $\mathbf{x}^T \mathbf{y} \in [0, 1]$ .

- Assume latent positions  $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_n \stackrel{i.i.d.}{\sim} F$ . Organize in the rows of a matrix  $\mathbf{X} = [\mathbf{X}_1 \mathbf{X}_2 \dots \mathbf{X}_n]^T$ .
- Associate each network with a diagonal matrix  $\mathbf{C}^{(g)} \in \mathbb{R}_{\geq 0}^{d \times d}$  such that for all  $\mathbf{x}, \mathbf{y} \in \text{supp}(F)$  and  $g \in [m]$ ,  $\mathbf{x}^T \mathbf{C}^{(g)} \mathbf{y} \in [0, 1]$ .

# Eigen-Scaling Random Dot Product Graph

## Eigen-Scaling Random Dot Product Graph

- Suppose that for  $\mathbf{y} \sim F$ ,  $\Delta = \mathbb{E}[\mathbf{y}\mathbf{y}^T]$  is diagonal and full rank and the matrices  $\{\mathbf{C}^{(g)}\}_{g=1}^m$  satisfy  $\min_{i \in [d]} \max_{g \in [m]} \mathbf{C}_{ii}^{(g)} > 0$ .
- Then the random adjacency matrices  $\{\mathbf{A}^{(g)}\}_{g=1}^m$  are said to be jointly distributed according to the *ESRDPG* with *latent positions*  $\mathbf{X}$  iff  $\{\mathbf{A}_{ij}^{(g)}\}$  are conditionally independent with

$$\mathbb{P}(\mathbf{A}_{ij}^{(g)} = 1 | \mathbf{X}_i, \mathbf{X}_j) = \mathbf{X}_i^T \mathbf{C}^{(g)} \mathbf{X}_j$$

- In essence,  $\mathbf{A}_{ij}^{(g)} | \mathbf{X} \stackrel{\text{ind.}}{\sim} \text{Bern}(\mathbf{X}_i^T \mathbf{C}^{(g)} \mathbf{X}_j)$ .
- Goal: Given  $\{\mathbf{A}^{(g)}\}_{g=1}^m$ , estimate  $\{\mathbf{X} \sqrt{\mathbf{C}^{(g)}}\}_{g=1}^m$

# Individual Network Embedding Techniques

- First approach: ignore shared structure and individually embed networks  $\mathbf{A}^{(g)}$  for  $g \in [m]$ .

## Adjacency Spectral Embedding (Sussman et al. 2012)

Let  $\mathbf{A}^{(g)}$  have eigendecomposition

$$\mathbf{A}^{(g)} = [\mathbf{U}_{\mathbf{A}^{(g)}} | \tilde{\mathbf{U}}_{\mathbf{A}^{(g)}}][\mathbf{S}_{\mathbf{A}^{(g)}} \oplus \tilde{\mathbf{S}}_{\mathbf{A}^{(g)}}][\mathbf{U}_{\mathbf{A}^{(g)}} | \tilde{\mathbf{U}}_{\mathbf{A}^{(g)}}]^T$$

where  $\mathbf{U}_{\mathbf{A}^{(g)}} \in \mathbb{R}^{n \times d}$  and  $\mathbf{S}_{\mathbf{A}^{(g)}} \in \mathbb{R}^{d \times d}$  contains the top  $d$  eigenvalues of  $\mathbf{A}^{(g)}$ . Then the ASE of  $\mathbf{A}^{(g)}$  is defined by  $\text{ASE}(\mathbf{A}^{(g)}, d) = \mathbf{U}_{\mathbf{A}^{(g)}} \mathbf{S}_{\mathbf{A}^{(g)}}^{1/2}$ .

- Second approach: assume identical structure and embed the sample mean matrix  $\bar{\mathbf{A}} = m^{-1} \sum_{g=1}^m \mathbf{A}^{(g)}$  by  $\text{ASE}(\bar{\mathbf{A}}, d)$ .

# Joint Network Embedding Techniques

- Third approach: *jointly* embed the networks  $\{\mathbf{A}^{(g)}\}_{g=1}^m$ .

## Omnibus Embedding (Levin et al. 2017)

Let the *omnibus matrix* be defined as

$$\tilde{\mathbf{A}} = \begin{bmatrix} \mathbf{A}^{(1)} & \frac{1}{2}[\mathbf{A}^{(1)} + \mathbf{A}^{(2)}] & \dots & \frac{1}{2}[\mathbf{A}^{(1)} + \mathbf{A}^{(m)}] \\ \frac{1}{2}[\mathbf{A}^{(2)} + \mathbf{A}^{(1)}] & \mathbf{A}^{(2)} & \dots & \frac{1}{2}[\mathbf{A}^{(2)} + \mathbf{A}^{(m)}] \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{2}[\mathbf{A}^{(m)} + \mathbf{A}^{(1)}] & \frac{1}{2}[\mathbf{A}^{(m)} + \mathbf{A}^{(2)}] & \dots & \mathbf{A}^{(m)} \end{bmatrix}.$$


Then the *omnibus embedding* of  $\{\mathbf{A}^{(g)}\}_{g=1}^m$  is given by  $\hat{\mathbf{L}} = \text{ASE}(\tilde{\mathbf{A}}, d)$ .

- Notice  $\hat{\mathbf{L}} \in \mathbb{R}^{nm \times d}$  so each vertex has a latent position estimate for each graph.



# Mean Squared Error Comparison

- Suppose  $\mathbf{A}^{(1)} \sim \text{ER}(p)$  and  $\mathbf{A}^{(2)} \sim \text{ER}(c^2 p)$
- Under ESRDPG:  $\mathbf{X} = \sqrt{p}\mathbf{1}_n$ ,  $\mathbf{C}^{(1)} = \mathbf{I}$ , and  $\mathbf{C}^{(2)} = c^2\mathbf{I}$



# Main Results

- Let  $\hat{\mathbf{L}} = \text{ASE}(\tilde{\mathbf{A}}, d)$  and  $h = n(g-1) + i$  for  $i \in [n]$  and  $g \in [m]$  so that  $\hat{\mathbf{L}}_h \in \mathbb{R}^{d \times 1}$  is some row of  $\hat{\mathbf{L}}$  written as a column vector.

## Theorem

- There exists diagonal matrices  $\{\mathbf{S}^{(g)}\}_{g=1}^m$  that only depend on  $\{\mathbf{C}^{(g)}\}_{g=1}^m$  and a sequence of orthogonal matrices  $\{\tilde{\mathbf{W}}_n\}_{n=1}^\infty$  such that

$$\hat{\mathbf{L}}\tilde{\mathbf{W}}_n - \mathbf{L} = (\mathbf{S}^{(g)} - \sqrt{\mathbf{C}^{(g)}})\mathbf{X}_i + \mathbf{R}_h \quad (1)$$

where  $\mathbf{R}_h$  is a residual.

- $\mathbf{R}_h$  satisfies  $\max_{h \in [nm]} \|\mathbf{R}_h\|_2 = O_{\mathbb{P}}\left(m^{3/2} \frac{\log nm}{\sqrt{n}}\right)$  and has asymptotic distribution

$$\lim_{n \rightarrow \infty} \mathbb{P}[\sqrt{n}\mathbf{R}_h \leq \mathbf{x}] = \int_{\text{supp}(F)} \Phi(\mathbf{x}; \mathbf{0}, \Sigma_g(\mathbf{y})) dF(\mathbf{y}). \quad (2)$$

# Simulation Experiment


## Simulation Design

- 1 Draw  $\mathbf{X}_1, \dots, \mathbf{X}_n \stackrel{i.i.d}{\sim} F$  where  $F$  corresponds to a two-group SBM with parameters  $(a = 0.25, b = 0.05)$ .
- 2 For  $t \in [0, 1]$ , draw  $(\{\mathbf{A}^{(g)}\}_{g=1}^2, \mathbf{X}) \sim \text{ESRDPG}(F, n, \{\mathbf{C}^{(g)}\}_{g=1}^m)$  with


$$\mathbf{C}^{(1)} = \mathbf{I} \quad \mathbf{C}^{(2)} := \mathbf{C}(t) = \begin{bmatrix} 1+t & 0 \\ 0 & 1-t \end{bmatrix}$$

- 3 Jointly embed  $\hat{\mathbf{L}} = \text{ASE}(\tilde{\mathbf{A}}, d)$
- At  $t = 0$ ,  $\mathbf{A}^{(2)}$  is a SBM and at  $t = 1$ ,  $\mathbf{A}^{(2)}$  is an Erdős-Rényi graph with parameter  $p = 0.3$ .
  - Goal: Analyze techniques that utilize  $\hat{\mathbf{L}}$  for statistical inference.

# Community Detection







# Two Graph Hypothesis Testing



# Conclusion & Future Work

- Preprint available: <https://arxiv.org/abs/2005.02511>

# References I

-  Avanti Athreya et al. “Statistical inference on random dot product graphs: A survey”. In: *Journal of Machine Learning Research* 18 (Sept. 2017).
-  A. Athreya et al. “A Limit Theorem for Scaled Eigenvectors of Random Dot Product Graphs”. In: *Sankhya A* 78.1 (Feb. 2016), pp. 1–18.
-  Peter D Hoff, Adrian E Raftery, and Mark S Handcock. “Latent Space Approaches to Social Network Analysis”. In: *Journal of the American Statistical Association* 97.460 (2002), pp. 1090–1098.
-  Mikko Kivelä et al. “Multilayer networks”. In: *Journal of Complex Networks* 2.3 (2014), pp. 203–271.

# References II



Keith Levin et al. “A Central Limit Theorem for an Omnibus Embedding of Multiple Random Dot Product Graphs”. In: Nov. 2017, pp. 964–967.



Daniel L. Sussman et al. “A Consistent Adjacency Spectral Embedding for Stochastic Blockmodel Graphs”. In: *Journal of the American Statistical Association* 107.499 (2012), pp. 1119–1128.



Shangsi Wang et al. *Joint Embedding of Graphs*. 2017.



Stephen J. Young and Edward R. Scheinerman. “Random Dot Product Graph Models for Social Networks”. In: *Algorithms and Models for the Web-Graph*. Ed. by Anthony Bonato and Fan R. K. Chung. Berlin, Heidelberg: Springer Berlin Heidelberg, 2007, pp. 138–149.