
One-sample / one repetition per setting
(a) Reference
(b) Prevalence $\pi = 0.01$
(c) Prevalence $\pi = 0.05$
(d) Prevalence $\pi = 0.50$
Balanced accuracy
1.0
0.8
0.6
0.4
0.2
0.0
Logistic correction
nnPU logistic
Naive CatBoost
CS CatBoost
RN CatBoost
PU gradient boosting
Same-model quantile
PU-estimated BA pilot
Oracle BA diagnostic


One-sample / one repetition per setting
(a) High dimension ($p = 500$)
(b) Curved boundary
(c) Feature interaction
(d) Unequal class covariance
Balanced accuracy
1.0
0.8
0.6
0.4
0.2
0.0
Logistic correction
nnPU logistic
Naive CatBoost
CS CatBoost
RN CatBoost
PU gradient boosting
Same-model quantile
PU-estimated BA pilot
Oracle BA diagnostic


Two-sample / one repetition per setting
(a) Reference
(b) Prevalence $\pi = 0.01$
(c) Prevalence $\pi = 0.05$
(d) Prevalence $\pi = 0.50$
Balanced accuracy
1.0
0.8
0.6
0.4
0.2
0.0
Logistic correction
nnPU logistic
Naive CatBoost
CS CatBoost
RN CatBoost
PU gradient boosting
Same-model quantile
PU-estimated BA pilot
Oracle BA diagnostic


Two-sample / one repetition per setting
(a) High dimension ($p = 500$)
(b) Curved boundary
(c) Feature interaction
(d) Unequal class covariance
Balanced accuracy
1.0
0.8
0.6
0.4
0.2
0.0
Logistic correction
nnPU logistic
Naive CatBoost
CS CatBoost
RN CatBoost
PU gradient boosting
Same-model quantile
PU-estimated BA pilot
Oracle BA diagnostic